



Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

Solving and Simulating Stochastic Differential Equations

Steven R. Dunbar

<2024-11-12 Tue>



Rule of Four

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method


Mathematical concepts should be presented:

- Symbolically
- Visually
- Numerically
- Verbally



Stochastic Process Family Tree

Definition 1.4:



Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method



Brownian Motion Paths

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar


Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

Definition 1.11





Calculating Probabilities 1

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion


Stochastic
Differential
Equations

Milstein Method

Definition 1.8 (2) and Proposition 1.1 (3):

Let $B(t)$ be standard Brownian motion.

- Find the probability that $0 < B(1) < 1$ ($= 0.34134$).
- Find the probability that $0 < B(1) < 1$ and $1 < B(2) < 3$ ($= 0.10046$).
- Find the probability that $0 < B(1) < 1$ and $1 < B(2) < 3$ and $0 < B(3) < 1/2$ ($= 0.00861$).





Calculating Probabilities 2

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

Definition 1.8 (2) and Proposition 1.1 (3):

Write the joint probability density function for $B(t_1) = x_1$ and $B(t_2) = x_2$ explicitly

$$f(x_1, t_1; x_2, t_2) = \frac{1}{2\pi\sqrt{t_1 \cdot (t_2 - t_1)}} e^{\frac{-x_1^2}{2t_1} - \frac{(x_2 - x_1)^2}{2(t_2 - t_1)}}$$

and integrate (with $t_1 = 1$, $t_2 = 2$)

$$\int_0^1 \int_1^3 \frac{1}{2\pi\sqrt{t_1 \cdot (t_2 - t_1)}} e^{\frac{-x_1^2}{2t_1} - \frac{(x_2 - x_1)^2}{2(t_2 - t_1)}} dx_1 dx_2 = 0.10046$$



Properties of Brownian Motion

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background


Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

Proposition 1.1 (1) and (2)

Sample Mean and StdDev over 1000 B_t





Euler-Maruyama Method: 1

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

The sample path that the Euler-Maruyama method produces numerically is the analog of using the Euler method.

The Euler-Maruyama (EM) method for

$$dX_t = G(t, X_t) dt + H(t, X_t) dB_t, \quad X(t_0) = X_0$$

is based on the definition of the Ito stochastic integral:

$$\begin{aligned} X_{j+1} &= X_j + G(t_j, X_j) dt + H(t_j, X_j)(B(t_j + dt) - B(t_j)) \\ t_{j+1} &= t_j + dt. \end{aligned}$$

Note that the initial conditions X_0 and t_0 set the starting point.



Euler-Maruyama Method: 2

$dX = 2X dt + X dB_t$, $X(0) = 1$ with $dt = 0.1$ and
 $dB_t = B(gt_{j-1} + dt) - B(t_{j-1})$ (Proposition 1.1 (3))

j	t_j	X_j	$2X_j dt$	dB_t	$X_j dB_t$	X_{j+1}
0	0	1	0.2	0.271	0.271	1.471
1	0.1	1.471	0.294	-0.049	-0.072	1.693
2	0.2	1.693	0.339	-0.448	-0.758	1.274
3	0.3	1.274	0.255	-0.21	-0.268	1.261
4	0.4	1.261	0.252	0.333	0.420	1.933
5	0.5	1.933	0.387	-0.683	-1.320	1.000
6	0.6	1.000	0.200	-0.292	-0.292	0.908
7	0.7	0.908	0.182	0.128	0.116	1.206
8	0.8	1.206	0.241	-0.34	-0.410	1.037
9	0.9	1.037	0.207	-0.147	-0.152	1.092
10	1.0	1.092				

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method



Error Measures:

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

Let $X_t^{(h)}$ be the numerical solution on $[0, T]$ with step-size $h = dt$.

Weak Error at time t (for a Lipschitz function f) is

$$\epsilon_w(h) = \left| \mathbb{E} [f(X_T)] - \mathbb{E} [f(X_T^{(h)})] \right|.$$

(This is the $L^1(\Omega)$ distance between $f(X_T)$ and $f(X_T^{(h)})$.)

Strong Error is:

$$\epsilon_s(h) = \mathbb{E} \left[\sup_{t \in [0, T]} \|X_t - X_t^{(h)}\|^2 \right]^{1/2}$$



Error Bounds:

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

Theorem (Euler-Maruyama Error Bounds)

$$\epsilon_w(h) = |\mathbb{E}[f(X_T)] - \mathbb{E}[f(X_T)]| = O(h)$$

$$\epsilon_s(h) = \mathbb{E} \left[\sup_{t \in [0, T]} \|X_t - X_t^{(h)}\|^2 \right]^{1/2} = O(h^{1/2})$$

Proof: Uses Ito's Isometry (LN, footnote 2), Gronwall's Inequality, Jensen's Inequality and some simple inequalities.



Other Numerical Methods:

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

- Euler-Heun(Maruyama) - Adapts the Euler-Heun method. Strong Order 0.5 in the Stratonovich sense.
- Milstein (1974) - Adds a 2nd-order correction based on Ito integral definition. Various adaptations have $\epsilon_w(h) = \epsilon_s(h) = O(h)$.
- Stochastic Runge-Kutta (difficulty is iterated stochastic integrals).
- Others: Achieve $\epsilon_w(h) = O(h^2)$ and $\epsilon_s(h) = O(h^{3/2})$ but need special noise structure for systems.



Example 2.1

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar


Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

$dN_t = rN_t dt + \alpha N_t dB_t$, $N(0) = N_0$ with $r = 1$, $\alpha = 1$,
 $N_0 = 1/2$. Solve with EM, $h = dt = 1/16$





Example 2.1 Ensembles

Solving and
Simulating
Stochastic
Differential
Equations


Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method





Example 2.1 Means, Variances

Solving and
Simulating
Stochastic
Differential
Equations


Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method





Example 2.2

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

Financial Model 2.2, (4b)

Pepsi and Walmart Stock prices, 2006

$$\begin{aligned}dX_p &= -0.0545X_p \, dt + 0.5X_p \, dB_p(t) + 0.1X_w \, dB_w(t) \\dX_w &= +0.0125X_w \, dt + 0.1X_p \, dB_p(t) + 0.5X_w \, dB_w(t)\end{aligned}$$



Example 2.2

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

Euler-Maruyama Method, $dt = 0.01$





Example 2.2

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar


Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

RKMilCommute - An explicit Runge-Kutta discretization of the strong order 1.0 Milstein method, adaptive step size.





Milstein Method 1

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

The Milstein method increases the accuracy of the E-M approximation by adding a second-order “correction” term derived from the stochastic Taylor series expansion of $X(t)$ by applying Ito’s lemma to the $G()$ and $H()$ functions.

$$\begin{aligned}dX_t &= G(t, X_t) dt + H(t, X_t) dB_t, & X(t_0) &= X_0 \\X_{j+1} &= X_j + G(t_j, X_j) dt + H(t_j, X_j) dB_j + \\&\quad \frac{1}{2} H(t_j, X_j) H'(t_j, X_j) ((dB_j)^2 - dt) \\t_{j+1} &= t_j + dt.\end{aligned}$$



Milstein Derivation for Geometric Brownian Motion

Solving and
Simulating
Stochastic
Differential
Equations

Steven R.
Dunbar

Background

Brownian
Motion

Stochastic
Differential
Equations

Milstein Method

$$dX_t = rX_t dt + \alpha X_t dB_t$$

$$d(\log X_t) = \left(r - \frac{\alpha^2}{2}\right) dt + \alpha dB_t \quad \text{LN, Ex 2.1, p.3}$$

$$X_{t+dt} = X_t \exp \left(\int_t^{t+dt} \left(r - \frac{\alpha^2}{2}\right) dt + \int_t^{t+dt} \alpha dB_t \right)$$

$$X_{t+dt} \approx X_t \left(1 + r dt - \frac{\alpha^2}{2} dt + \alpha dB_t + \frac{\alpha^2}{2} (dB_t)^2 \right)$$

$$= X_t + rX_t dt + \alpha X_t dB_t +$$

$$\frac{1}{2} \alpha X_t \cdot \alpha ((dB_t)^2 - dt)$$