

# Notes on Fourier Analysis

October 4, 2023

## Contents

<b>1</b>	<b>Fourier Series</b>	<b>2</b>
1.1	It all Adds Up . . . . .	2

# 1 Fourier Series

## 1.1 It all Adds Up

$$\frac{a_0}{2} + \sum_{n=1}^N (a_n \cos(2\pi nt) + b_n \sin(2\pi nt))$$

Given that

$$e^{2\pi int} = \cos(2\pi nt) + i \sin(2\pi nt)$$

and

$$e^{-2\pi int} = \cos(2\pi nt) - i \sin(2\pi nt)$$

Since  $\cos(-x) = \cos(x)$  while  $\sin(-x) = -\sin(x)$ .

The above also mean that

$$\cos(2\pi nt) = \frac{1}{2} (e^{2\pi int} + e^{-2\pi int})$$

and

$$\sin(2\pi nt) = \frac{1}{2i} (e^{2\pi int} - e^{-2\pi int}) = -\frac{i}{2} (e^{2\pi int} - e^{-2\pi int})$$

In the last step we used the fact that  $i^{-1} = -i$ . ( $1 = (-i)i = i \cdot i^{-1}$ ).

Using these expressions to rewrite our series of sines and cosines we get

$$\begin{aligned} \frac{a_0}{2} + \sum_{n=1}^N (a_n \cos(2\pi nt) + b_n \sin(2\pi nt)) &= \frac{a_0}{2} + \sum_{n=1}^N \left[ \frac{a_n}{2} (e^{2\pi int} + e^{-2\pi int}) + \frac{b_n}{2i} (e^{2\pi int} - e^{-2\pi int}) \right] \\ &= \frac{a_0}{2} + \sum_{n=1}^N \left[ \frac{a_n}{2} (e^{2\pi int} + e^{-2\pi int}) - \frac{ib_n}{2} (e^{2\pi int} - e^{-2\pi int}) \right] \\ &= \frac{a_0}{2} + \sum_{n=1}^N \left[ \frac{a_n}{2} e^{2\pi int} + \frac{a_n}{2} e^{-2\pi int} - \frac{ib_n}{2} e^{2\pi int} + \frac{ib_n}{2} e^{-2\pi int} \right] \\ &= \frac{a_0}{2} + \sum_{n=1}^N \left[ \frac{1}{2} (a_n - ib_n) e^{2\pi int} + \frac{1}{2} (a_n + ib_n) e^{-2\pi int} \right] \end{aligned}$$

If we look at the terms within the square braces we see that we have a complex number times and exponent and its complex conjugate,

$$c_n := \frac{1}{2} (a_n - ib_n)$$

and

$$\bar{c}_n = \frac{1}{2} (a_n + ib_n)$$

So

$$\frac{a_0}{2} + \sum_{n=1}^N (a_n \cos(2\pi nt) + b_n \sin(2\pi nt)) = \frac{a_0}{2} + \sum_{n=1}^N [c_n e^{2\pi int} + \bar{c}_n e^{-2\pi int}]$$

Here is where the additional requirement comes in. If we have  $\bar{c}_n = c_{-n}$ , then  $a_{-n} = a_n$  and  $b_{-n} = -b_n$ . We can then reindex our series,

$$\begin{aligned} \frac{a_0}{2} + \sum_{n=1}^N (a_n \cos(2\pi nt) + b_n \sin(2\pi nt)) &= \frac{a_0}{2} + \sum_{n=1}^N \left[ c_n e^{2\pi i n t} + \bar{c}_n \overline{e^{2\pi i n t}} \right] \\ &= \frac{a_0}{2} + \sum_{n=1}^N c_n e^{2\pi i n t} + \sum_{n=-1}^{-N} c_{-n} e^{-2\pi i n t} \end{aligned}$$

and if we have  $c_0 = \frac{1}{2}(a_0 - ib_0) = \frac{a_0}{2}$ , ( $b_0 = 0$ ), then we can see where

$$\sum_{n=-N}^N c_n e^{2\pi i n t}$$

comes from.