The Mass-Radius Relation for 71 Exoplanets with $R_{\rm P} < 7 R_{\oplus}$

L. M. Weiss & G. W. Marcy

Received	_;	accepted
----------	----	----------

1. Introduction

The Kepler Mission has found an arbundance of planets with $R < 4R_{\oplus}$. However, in many systems, it is difficult to measure the masses of such small planets because the gravitational acceleration these planets induce on their host stars or neighboring planets is too small to detect with current telescopes and instruments. How can we determine the composition of these planets? To understand the nature of small planets, we reverse the question: how do the mass of a planet, its composition, and the incident stellar flux from its star affect the radius of a planet?

Our work (Weiss et al. 2013) has shown that for planets between ~ 20 and thousands of Earth masses, we can predict the radius of a planet from its mass and incident stellar flux, suggesting that planets of a given mass have a particular composition. However, below $\sim 20~M_{\oplus}$, the large apparent scatter makes predicting planet radius harder. At 10 Earth masses, planets can have a radius of 2-8 R_{\oplus} , which spans a range in density from less dense than water to solid iron.

There are two explanations for the large apparent scatter in planet radius: either errors in radius and/or mass are underestimated for planets below 20 M_{\oplus} , or the (real) scatter in radius indicates a diversity of planet compositions. Identifying whether this apparent scatter is real or not is one of the most pressing issues in understanding planetary compositions today. Does the mass of a planet uniquely determine a planets composition, or is there diversity of the rock to volatile ratio at a given mass? If the scatter is only apparent, then it is possible that there is a one-to-one correspondence between planet mass and composition below 20 M_{\oplus} . However, if the scatter is real, then there must be a diversity of planet compositions

at a given mass below 20 M_{\oplus} . Further refinements to the measurements of planet masses, especially below 20 M_{\oplus} , is necessary to test to what extent the apparent scatter of planetary masses at a given radius is real.

2. Justification for a Mass-Radius Relation for Low-Mass Exoplanets

Is there a correlation between planet mass and radius for small exoplanets? By eye, we might judge that there is a correlation, but what is the probability that we have fooled ourselves into believing there is a correlation when in fact there is none?

To answer this question, I calculated the probability that the data were uncorrelated. First, I calculated the correlation coefficient (Pearson R test) r = 0.61. For r = 0.61, the Student t value is 6.38. There are 71 planets with $R_P < 7R_{\oplus}$, so there are 69 degrees of freedom in this sample. What is the probability that among data with 69 degrees of freedom, we find a Student t value of 6.38 if the data are uncorrelated? According to a Wolfram Alpha calculation, the probability of obtaining $t \geq 6.38$ for 69 degrees of freedom, given that the data are really uncorrelated, is 3.56×10^{-8} . We can safely say that the masses and radii of small planets are correlated.

Does this correlation also exist for planets with $R_{\rm P} < 4R_{\oplus}$? There are 58 planets smaller than 4 R_{\oplus} , leaving 56 degrees of freedom. The Pearson R coefficient is 0.51, and the Student t value is 4.4. The probability that these data are uncorrelated is 8.4×10^{-5} . Thus, the masses and radii of planets between the sizes of Earth and Neptune are correlated.

3. The Updated Mass-Radius Relation for Small Exoplanets

Now that we have determined that a correlation between planet mass and radius exists, what is the best way to model the relation between exoplanet mass and radius? Figure 2 suggests that the data can be fit with a line; a more complicated fit is not warranted. To check this, we try a traditional power-law fit and obtain $M_{\rm P} \propto R_{\rm P}^1$ as the best result.

The best linear fit to the data for $R_P < 7$ R_{\oplus} is:

$$M_{\rm P}/M_{\oplus} = -0.15 + 3.01 R_{\rm P}/R_{\oplus}$$

There are 71 exoplanets in this sample. The reduced χ^2 =5.90, and the RMS is 9.37 M_{\oplus} .

However, we notice in Figure 1 that the weighted mean masses might follow a different trend above and below 4 R_{\oplus} . Also, there are very few planets with $4 < R_{\rm P} < 7$, and the scatter in their masses is large. This motivates us to do a linear fit between mass and radius on the subsample of exoplanets with $R_{\rm P} < 4$, obtaining:

$$M_{\rm P}/M_{\oplus} = 1.01 + 2.33 R_{\rm P}/R_{\oplus}$$

There are 58 exoplanets in this sample. The reduced $\chi^2=3.74$, and the RMS is $5.01M_{\oplus}$.

We present an update to the relation between exoplanet mass, exoplanet radius, and incident flux from the host star established in Weiss et al. (2013). In this paper, we determine a statistical relation between the masses and radii of 225 transiting exoplanets with measurements or upper limits to their masses. These exoplanets

have radii spanning $0.30 R_{\oplus}$ to $22.8 R_{\oplus}$ and orbital periods spanning 0.67 days to 229 days. The coldest exoplanet (Kepler-16b) receives only 0.35 of the incident flux that Earth does, whereas the hottest exoplanet receives 53,000 times Earth's incident flux.

This update benefits from (1) the masses of 42 low-mass planets recently characterized in ?, (2) additional planets of all masses added to exoplanets.org between 27 September 2012 and [use most recent update] 2013, and (3) refined planet and stellar parameters for several systems, including 55 Cnc e (cite), Kepler-11 (?), GJ 3470 (?), HD 97658 b (?), [... other systems].

3.1. Using Negative Planet Masses for Statistical Soundness

? allow "negative" planet masses as Keplerian orbital solutions to avoid the Lutz-Kelker bias. A "negative" planet mass arises when, given the orbital period and phase determined by the transit, the RV data have the opposite sign from what is expected. This yields a negative semi-amplitude K in the orbital solution, which results in a negative planet mass. In a traditional analysis, the uncertainties in the RVs and the number of measurements combine to determine an upper limit to the planet mass. However, upper limits to planet masses are difficult to use statistically. The artifice of a negative planet mass allows the population of small planets to be treated statistically. Since there is no bias toward large or small planet masses in our sample, we can take the weighted mean mass of planets of a given radius.

Table 1. Exoplanets with Measured Mass and Radius

Name	Planet Mass (M_{\bigoplus})	Planet Radius (R_{\oplus})	Incident Flux (F_{\bigoplus})	Period (d)	First Ref.	Orbital Ref.
55 Cnc e	8.380	2.210	2439.690	0.737	McArthur et al. (2004)	Endl et al. (2012)
CoRoT-7 b	5.021	1.679	1779.433	0.854	Queloz et al. (2009) ; Léger et al. (2009)	Queloz et al. (2009)
CoRoT-8 b	68.673	6.389	88.184	6.212	Bordé et al. (2010)	Bordé et al. (2010)
GJ 1214 b	6.260	2.800	16.631	1.580	Charbonneau et al. (2009)	Carter et al. (2011)
GJ 3470 b	13.900	4.830	38.335	3.337	Bonfils et al. (2012)	?
GJ 436 b	23.105	4.222	29.882	2.644	Butler et al. (2004)	Maness et al. (2007)
HAT-P-11 b	26.231	4.730	97.355	4.888	Bakos et al. (2010)	Bakos et al. (2010)
HAT-P-26 b	18.640	6.333	163.050	4.235	Hartman et al. (2011b)	Hartman et al. (2011b
HD 97658 b	7.862	2.340	48.052	9.491	?	?
Kepler-10 b	4.539	1.416	3572.048	0.837	Batalha et al. (2011)	Batalha et al. (2011)
Kepler-11 b	1.900	1.800	126.384	10.304	Lissauer et al. (2011)	?
Kepler-11 c	2.900	2.870	91.413	13.025	Lissauer et al. (2011)	Lissauer et al. (2011)
Kepler-11 d	7.300	3.120	43.562	22.687	Lissauer et al. (2011)	Lissauer et al. (2011)
Kepler-11 e	8.000	4.190	27.524	31.996	Lissauer et al. (2011)	Lissauer et al. (2011)
Kepler-11 f	2.000	2.490	16.745	46.689	Lissauer et al. (2011)	Lissauer et al. (2011)
Kepler-18 b	6.900	2.000	462.244	3.505	Borucki et al. (2011)	Cochran et al. (2011)
Kepler-18 c	17.299	5.490	163.493	7.642	Borucki et al. (2011)	Cochran et al. (2011)
Kepler-18 d	16.399	6.980	67.364	14.859	Borucki et al. (2011)	Cochran et al. (2011)
Kepler-20 b	8.474	1.908	343.928	3.696	Borucki et al. (2011)	Gautier et al. (2012)
Kepler-20 c	15.734	3.067	81.783	10.854	Borucki et al. (2011)	Gautier et al. (2012)
Kepler-20 d	7.528	2.748	5.937	77.612	Borucki et al. (2011)	Gautier et al. (2012)
Kepler-36 b	4.461	1.485	217.365	13.840	Borucki et al. (2011)	Carter et al. (2012)
Kepler-36 c	8.101	3.676	175.646	16.239	Carter et al. (2012)	Carter et al. (2012)
Kepler-4 b	24.544	4.002	1123.918	3.213	Borucki et al. (2010)	Borucki et al. (2010)
Kepler-68 b	8.300	2.310	409.092	5.399	Borucki et al. (2011)	Gilliland et al. (2013)
Kepler-68 c	4.377	0.952	189.764	9.605	Batalha et al. (2013)	Gilliland et al. (2013)
KOI-94 b	9.400	1.770	1155.374	3.743	Weiss et al. (2013)	Weiss et al. (2013)
KOI-94 c	8.300	4.280	295.035	10.424	Borucki et al. (2011)	Weiss et al. (2013)
KOI-94 d	105.000	11.400	106.760	22.343	Borucki et al. (2011)	Weiss et al. (2013)
KOI-94 e	38.000	6.640	32.631	54.320	Borucki et al. (2011)	Weiss et al. (2013)
KOI-41.01	0.85000000	2.2000000	1156.6217	12.815900		
KOI-41.02	7.3400000	1.3000000	65.166782	6.8870500		
KOI-41.03	-5.3100000	1.6000000	85.869041	35.333100		
KOI-69.01	2.5900000	1.5000000	172.33078	4.7267400		
KOI-82.01	8.9300000	2.2000000	806.96474	16.145700		
KOI-82.02	3.8000000	1.2000000	147.34753	10.311700		
KOI-82.03	8.1200000	0.90000000	447.82440	27.453600		
KOI-82.04	-2.4500000	0.60000000	447.27347	7.0714200		
KOI-82.05	0.41000000	0.50000000	2950.4414	5.2869600		
KOI-104.01	10.840000	3.5000000	1189.7862	2.5080600		
KOI-108.01	18.690000	3.4000000	348.15857	15.965400		




Fig. 1.— The weighted mean mass (left), density (center), and incident flux (right) in bins of $1 R_{\oplus}$. The error bars are the weighted uncertainties in the means.

Table 1—Continued

Name	Planet Mass	Planet Radius	Incident Flux	Period	First Ref.	Orbital Ref
	(M_{\oplus})	(R_{\bigoplus})	(F_{\bigoplus})	(d)		
KOI-108.02	-22.540000	5.1000000	163.05049	179.61200		
KOI-116.01	10.440000	2.5000000	308.73816	13.570800		
KOI-116.02	11.170000	2.6000000	604.88523	43.844500		
KOI-116.03	0.15000000	0.80000000	416.85849	6.1648600		
KOI-116.04	-13.340000	0.90000000	298.50217	23.980200		
KOI-122.01	13.000000	3.4000000	1191.9507	11.523100		
KOI-123.01	1.3000000	2.4000000	618.50330	6.4816300		
KOI-123.02	2.2200000	2.5000000	1914.9410	21.222700		
KOI-148.01	3.9400000	1.9000000	1903.1669	4.7780000		
KOI-148.02	14.610000	2.7000000	537.46087	9.6739500		
KOI-148.03	7.9300000	2.0000000	1030.1836	42.896100		
KOI-153.01	-5.7000000	2.2000000	1818.3791	8.9250700		
KOI-153.02	7.1000000	1.8000000	439.71133	4.7540000		
KOI-244.01	24.600000	5.2000000	226.04191	12.720400		
KOI-244.02	9.6000000	2.7000000	1560.3683	6.2385000		
KOI-245.01	-5.9800000	1.9000000	1367.9423	39.792200		
KOI-245.02	3.3500000	0.80000000	1611.6302	21.302000		
KOI-245.03	-0.42000000	0.30000000	2337.2286	13.367500		
KOI-246.01	7.8900000	2.3000000	926.85884	5.3987500		
KOI-246.02	2.1800000	1.0000000	1299.0590	9.6050400		
KOI-261.01	8.4600000	2.7000000	4070.4875	16.238500		
KOI-283.01	16.130000	2.4000000	1637.1673	16.092000		
KOI-283.02	17.020000	0.80000000	909.50346	25.516900		
KOI-292.01	3.5100000	1.5000000	584.57028	2.5866400		
KOI-299.01	3.5500000	2.0000000	1299.4980	1.5416800		
KOI-305.01	6.1500000	1.5000000	142.83632	4.6035800		
KOI-321.01	6.3500000	1.4000000	344.40060	2.4262900		
KOI-321.02	2.7100000	0.80000000	725.85388	4.6233200		
KOI-1442.01	0.060000000	1.1000000	11.621547	0.66931000		
KOI-1612.01	0.48000000	0.80000000	1420.2428	2.4650200		
KOI-1925.01	2.6900000	1.2000000	418.26336	68.958400		




Fig. 2.— Mass vs. radius for planets with $R_{\rm P} < 7R_{\oplus}$. Individual planets are represented by black dots, with error bars corresponding to 1σ uncertainties. The dashed line is the best linear fit to the data. The dotted line is a simple quadratic fit. Red squares represent the weighted mean mass in bins of $1 R_{\oplus}$, and error bars are the weighted uncertainty in the mean mass (as in Figure 1), to guide the eye.




Fig. 3.— Mass vs. radius for planets with $R_{\rm P} < 4R_{\oplus}$. Individual planets are represented by black dots, with error bars corresponding to 1σ uncertainties. The dashed line is the best linear fit to the data. The dotted line is a simple quadratic fit. Red squares represent the weighted mean mass in bins of $1 R_{\oplus}$, and error bars are the weighted uncertainty in the mean mass (as in Figure 1), to guide the eye. The solar system planets are plotted as blue triangles for reference.




Fig. 4.— Radius versus mass for exoplanets with measured masses and radii. Planets receiving lower than the median incident flux are blue; those receiving higher than the median incident flux are red. A plane was fit to the logspace points to determine a mass-radius-flux relation; the projection of that plane at the median flux is shown (black line). The Solar System planets are over plotted as black triangles for comparison, although they are not used in determining the fit.




Fig. 5.— Density versus mass for exoplanets with measured masses and radii. Planets receiving lower than the median incident flux are blue; those receiving higher than the median incident flux are red. A plane was fit to the logspace points to determine a mass-radius-flux relation; the projection of that plane at the median flux is shown (black line). The Solar System planets are over plotted as black triangles for comparison, although they are not used in determining the fit.




Fig. 6.— Radius vs. flux for exoplanets with measured masses and radii. Giant planets are orange; low-mass planets are green.

4. Discussion

REFERENCES

Aigrain, S., Collier Cameron, A., Ollivier, M., et al. 2008, A&A, 488, L43

Alonso, R., Brown, T. M., Torres, G., et al. 2004, ApJ, 613, L153

Alonso, R., Auvergne, M., Baglin, A., et al. 2008, A&A, 482, L21

Alsubai, K. A., Parley, N. R., Bramich, D. M., et al. 2011, MNRAS, 417, 709

Anderson, D. R., Gillon, M., Hellier, C., et al. 2008, MNRAS, 387, L4

Anderson, D. R., Hellier, C., Gillon, M., et al. 2010, ApJ, 709, 159

Anderson, D. R., Collier Cameron, A., Hellier, C., et al. 2011a, A&A, 531, A60

Anderson, D. R., Barros, S. C. C., Boisse, I., et al. 2011b, PASP, 123, 555

Bakos, G. A., Noyes, R. W., Kovács, G., et al. 2007a, ApJ, 656, 552

Bakos, G. Á., Shporer, A., Pál, A., et al. 2007b, ApJ, 671, L173

Bakos, G. Á., Kovács, G., Torres, G., et al. 2007c, ApJ, 670, 826

Bakos, G. Á., Pál, A., Torres, G., et al. 2009a, ApJ, 696, 1950

Bakos, G. A., Howard, A. W., Noyes, R. W., et al. 2009b, ApJ, 707, 446

Bakos, G. Á., Torres, G., Pál, A., et al. 2010, ApJ, 710, 1724

Bakos, G. Á., Hartman, J., Torres, G., et al. 2011, ApJ, 742, 116

Bakos, G. Á., Hartman, J. D., Torres, G., et al. 2012, AJ, 144, 19

Barbieri, M., Alonso, R., Desidera, S., et al. 2009, A&A, 503, 601

Barge, P., Baglin, A., Auvergne, M., et al. 2008, A&A, 482, L17

Barros, S. C. C., Faedi, F., Collier Cameron, A., et al. 2011, A&A, 525, A54

Batalha, N. M., Borucki, W. J., Bryson, S. T., et al. 2011, ApJ, 729, 27

Batalha, N. M., Rowe, J. F., Bryson, S. T., et al. 2013, ApJS, 204, 24

Beatty, T. G., Pepper, J., Siverd, R. J., et al. 2012, ApJ, 756, L39

Bonfils, X., Gillon, M., Udry, S., et al. 2012, A&A, 546, A27

Bonomo, A. S., Santerne, A., Alonso, R., et al. 2010, A&A, 520, A65

Bonomo, A. S., Hébrard, G., Santerne, A., et al. 2012, A&A, 538, A96

Bordé, P., Bouchy, F., Deleuil, M., et al. 2010, A&A, 520, A66

Borucki, W. J., Koch, D. G., Brown, T. M., et al. 2010, ApJ, 713, L126

Borucki, W. J., Koch, D. G., Basri, G., et al. 2011, ApJ, 736, 19

Bouchy, F., Udry, S., Mayor, M., et al. 2005, A&A, 444, L15

Bouchy, F., Hebb, L., Skillen, I., et al. 2010, A&A, 519, A98

Bouchy, F., Bonomo, A. S., Santerne, A., et al. 2011, A&A, 533, A83

Bryan, M. L., Alsubai, K. A., Latham, D. W., et al. 2012, ApJ, 750, 84

Buchhave, L. A., Bakos, G. Á., Hartman, J. D., et al. 2010, ApJ, 720, 1118

—. 2011a, ApJ, 733, 116

Buchhave, L. A., Latham, D. W., Carter, J. A., et al. 2011b, ApJS, 197, 3

Burke, C. J., McCullough, P. R., Valenti, J. A., et al. 2007, ApJ, 671, 2115

—. 2008, ApJ, 686, 1331

Butler, R. P., Vogt, S. S., Marcy, G. W., et al. 2004, ApJ, 617, 580

Cabrera, J., Bruntt, H., Ollivier, M., et al. 2010, A&A, 522, A110

Carter, J. A., Winn, J. N., Gilliland, R., & Holman, M. J. 2009, ApJ, 696, 241

Carter, J. A., Winn, J. N., Holman, M. J., et al. 2011, ApJ, 730, 82

Carter, J. A., Agol, E., Chaplin, W. J., et al. 2012, Science, 337, 556

Charbonneau, D., Brown, T. M., Latham, D. W., & Mayor, M. 2000, ApJ, 529, L45

Charbonneau, D., Winn, J. N., Everett, M. E., et al. 2007, ApJ, 658, 1322

Charbonneau, D., Berta, Z. K., Irwin, J., et al. 2009, Nature, 462, 891

Christian, D. J., Gibson, N. P., Simpson, E. K., et al. 2009, MNRAS, 392, 1585

Cochran, W. D., Fabrycky, D. C., Torres, G., et al. 2011, ApJS, 197, 7

Collier Cameron, A., Bouchy, F., Hébrard, G., et al. 2007, MNRAS, 375, 951

Csizmadia, S., Moutou, C., Deleuil, M., et al. 2011, A&A, 531, A41

Deeg, H. J., Moutou, C., Erikson, A., et al. 2010, Nature, 464, 384

Deleuil, M., Deeg, H. J., Alonso, R., et al. 2008, A&A, 491, 889 Désert, J.-M., Charbonneau, D., Demory, B.-O., et al. 2011, ApJS, 197, 14 Doyle, L. R., Carter, J. A., Fabrycky, D. C., et al. 2011, Science, 333, 1602 Dunham, E. W., Borucki, W. J., Koch, D. G., et al. 2010, ApJ, 713, L136 Endl, M., MacQueen, P. J., Cochran, W. D., et al. 2011, ApJS, 197, 13 Endl, M., Robertson, P., Cochran, W. D., et al. 2012, ApJ, 759, 19 Enoch, B., Cameron, A. C., Anderson, D. R., et al. 2011, MNRAS, 410, 1631 Faedi, F., Barros, S. C. C., Anderson, D. R., et al. 2011, A&A, 531, A40 Fischer, D. A., Vogt, S. S., Marcy, G. W., et al. 2007, ApJ, 669, 1336 Fortney, J. J., Demory, B.-O., Désert, J.-M., et al. 2011, ApJS, 197, 9 Fridlund, M., Hébrard, G., Alonso, R., et al. 2010, A&A, 512, A14 Gandolfi, D., Hébrard, G., Alonso, R., et al. 2010, A&A, 524, A55 Gautier, III, T. N., Charbonneau, D., Rowe, J. F., et al. 2012, ApJ, 749, 15 Gilliland, R. L., Marcy, G. W., Rowe, J. F., et al. 2013, ApJ, 766, 40 Gillon, M., Anderson, D. R., Triaud, A. H. M. J., et al. 2009, A&A, 501, 785 Gillon, M., Hatzes, A., Csizmadia, S., et al. 2010, A&A, 520, A97 Gillon, M., Doyle, A. P., Lendl, M., et al. 2011, A&A, 533, A88

Guenther, E. W., Díaz, R. F., Gazzano, J.-C., et al. 2012, A&A, 537, A136

Hartman, J. D., Bakos, G. Á., Torres, G., et al. 2009, ApJ, 706, 785

Hartman, J. D., Bakos, G. Á., Sato, B., et al. 2011a, ApJ, 726, 52

Hartman, J. D., Bakos, G. Á., Kipping, D. M., et al. 2011b, ApJ, 728, 138

Hartman, J. D., Bakos, G. Á., Torres, G., et al. 2011c, ApJ, 742, 59

Hartman, J. D., Bakos, G. Á., Béky, B., et al. 2012, AJ, 144, 139

Hebb, L., Collier-Cameron, A., Loeillet, B., et al. 2009, ApJ, 693, 1920

Hebb, L., Collier-Cameron, A., Triaud, A. H. M. J., et al. 2010, ApJ, 708, 224

Hébrard, G., Evans, T. M., Alonso, R., et al. 2011, A&A, 533, A130

Hébrard, G., Collier Cameron, A., Brown, D. J. A., et al. 2013, A&A, 549, A134

Hellier, C., Anderson, D. R., Collier Cameron, A., et al. 2009a, Nature, 460, 1098

Hellier, C., Anderson, D. R., Gillon, M., et al. 2009b, ApJ, 690, L89

Hellier, C., Anderson, D. R., Collier Cameron, A., et al. 2010, ApJ, 723, L60

- —. 2011, A&A, 535, L7
- —. 2012, MNRAS, 426, 739

Henry, G. W., Marcy, G. W., Butler, R. P., & Vogt, S. S. 2000, ApJ, 529, L41

Howard, A. W., Marcy, G. W., Bryson, S. T., et al. 2012, ApJS, 201, 15

Jenkins, J. M., Borucki, W. J., Koch, D. G., et al. 2010, ApJ, 724, 1108

Johns-Krull, C. M., McCullough, P. R., Burke, C. J., et al. 2008, ApJ, 677, 657

Johnson, J. A., Winn, J. N., Bakos, G. Á., et al. 2011, ApJ, 735, 24

Johnson, J. A., Gazak, J. Z., Apps, K., et al. 2012, AJ, 143, 111

Joshi, Y. C., Pollacco, D., Collier Cameron, A., et al. 2009, MNRAS, 392, 1532

Kipping, D. M., Bakos, G. Á., Hartman, J., et al. 2010, ApJ, 725, 2017

Kipping, D. M., Hartman, J., Bakos, G. A., et al. 2011, AJ, 142, 95

Koch, D. G., Borucki, W. J., Rowe, J. F., et al. 2010, ApJ, 713, L131

Kovács, G., Bakos, G. Á., Torres, G., et al. 2007, ApJ, 670, L41

Kovács, G., Bakos, G. Á., Hartman, J. D., et al. 2010, ApJ, 724, 866

Latham, D. W., Bakos, G. Á., Torres, G., et al. 2009, ApJ, 704, 1107

Latham, D. W., Borucki, W. J., Koch, D. G., et al. 2010, ApJ, 713, L140

Léger, A., Rouan, D., Schneider, J., et al. 2009, A&A, 506, 287

Lissauer, J. J., Fabrycky, D. C., Ford, E. B., et al. 2011, Nature, 470, 53

Lister, T. A., Anderson, D. R., Gillon, M., et al. 2009, ApJ, 703, 752

Maciejewski, G., Errmann, R., Raetz, S., et al. 2011, A&A, 528, A65

Mandushev, G., O'Donovan, F. T., Charbonneau, D., et al. 2007, ApJ, 667, L195

Mandushev, G., Quinn, S. N., Buchhave, L. A., et al. 2011, ApJ, 741, 114

Maness, H. L., Marcy, G. W., Ford, E. B., et al. 2007, PASP, 119, 90

Maxted, P. F. L., Anderson, D. R., Gillon, M., et al. 2010a, AJ, 140, 2007

Maxted, P. F. L., Anderson, D. R., Collier Cameron, A., et al. 2010b, PASP, 122, 1465

—. 2011, PASP, 123, 547

—. 2013, PASP, 125, 48

McArthur, B. E., Endl, M., Cochran, W. D., et al. 2004, ApJ, 614, L81

McCullough, P. R., Stys, J. E., Valenti, J. A., et al. 2006, ApJ, 648, 1228

Moutou, C., Bruntt, H., Guillot, T., et al. 2008, A&A, 488, L47

Moutou, C., Hébrard, G., Bouchy, F., et al. 2009, A&A, 498, L5

Naef, D., Latham, D. W., Mayor, M., et al. 2001, A&A, 375, L27

Noyes, R. W., Bakos, G. Á., Torres, G., et al. 2008, ApJ, 673, L79

O'Donovan, F. T., Charbonneau, D., Mandushev, G., et al. 2006, ApJ, 651, L61

Ollivier, M., Gillon, M., Santerne, A., et al. 2012, A&A, 541, A149

Pál, A., Bakos, G. Á., Torres, G., et al. 2008, ApJ, 680, 1450

—. 2010, MNRAS, 401, 2665

Penev, K., Bakos, G. Á., Bayliss, D., et al. 2013, AJ, 145, 5

Pollacco, D., Skillen, I., Collier Cameron, A., et al. 2008, MNRAS, 385, 1576

Pont, F., Tamuz, O., Udalski, A., et al. 2008, A&A, 487, 749

Queloz, D., Bouchy, F., Moutou, C., et al. 2009, A&A, 506, 303

Queloz, D., Anderson, D., Collier Cameron, A., et al. 2010, A&A, 517, L1

Rauer, H., Queloz, D., Csizmadia, S., et al. 2009, A&A, 506, 281

Rouan, D., Parviainen, H., Moutou, C., et al. 2012, A&A, 537, A54

Santerne, A., Bonomo, A. S., Hébrard, G., et al. 2011, A&A, 536, A70

Sato, B., Fischer, D. A., Henry, G. W., et al. 2005, ApJ, 633, 465

Sato, B., Hartman, J. D., Bakos, G. Á., et al. 2012, PASJ, 64, 97

Shporer, A., Bakos, G. Á., Bouchy, F., et al. 2009, ApJ, 690, 1393

Simpson, E. K., Pollacco, D., Cameron, A. C., et al. 2011a, MNRAS, 414, 3023

Simpson, E. K., Faedi, F., Barros, S. C. C., et al. 2011b, AJ, 141, 8

Skillen, I., Pollacco, D., Collier Cameron, A., et al. 2009, A&A, 502, 391

Smalley, B., Anderson, D. R., Collier Cameron, A., et al. 2010, A&A, 520, A56

—. 2011, A&A, 526, A130

Smalley, B., Anderson, D. R., Collier-Cameron, A., et al. 2012, A&A, 547, A61

Smith, A. M. S., Anderson, D. R., Collier Cameron, A., et al. 2012, AJ, 143, 81

Snellen, I. A. G., Koppenhoefer, J., van der Burg, R. F. J., et al. 2009, A&A, 497, 545

Street, R. A., Simpson, E., Barros, S. C. C., et al. 2010, ApJ, 720, 337

Tingley, B., Endl, M., Gazzano, J.-C., et al. 2011, A&A, 528, A97

Torres, G., Winn, J. N., & Holman, M. J. 2008, ApJ, 677, 1324

Torres, G., Bakos, G. Á., Kovács, G., et al. 2007, ApJ, 666, L121

Torres, G., Bakos, G. Á., Hartman, J., et al. 2010, ApJ, 715, 458

Triaud, A. H. M. J., Queloz, D., Hellier, C., et al. 2011, A&A, 531, A24

Triaud, A. H. M. J., Anderson, D. R., Collier Cameron, A., et al. 2013, A&A, 551, A80

Tripathi, A., Winn, J. N., Johnson, J. A., et al. 2010, ApJ, 715, 421

Udalski, A., Pont, F., Naef, D., et al. 2008, A&A, 482, 299

Weiss, L. M., Marcy, G. W., Rowe, J. F., et al. 2013, ApJ, 768, 14

West, R. G., Anderson, D. R., Gillon, M., et al. 2009a, AJ, 137, 4834

West, R. G., Collier Cameron, A., Hebb, L., et al. 2009b, A&A, 502, 395

Wilson, D. M., Gillon, M., Hellier, C., et al. 2008, ApJ, 675, L113

Winn, J. N., Johnson, J. A., Albrecht, S., et al. 2009, ApJ, 703, L99

Winn, J. N., Johnson, J. A., Howard, A. W., et al. 2010, ApJ, 718, 575

This manuscript was prepared with the AAS LATEX macros v5.2.