

Trace Metal Parameter Data

Table 2. Conditional stability constants for metals in seawater (Millero, 2001b)

Metal	[M]	[L]	log K _c
Cu(II)	1–10 nM	2–60 nM	8.5
Zn(II)	0.1–2 nM	1.2 nM	12
Cd(II)	2–800 pM	100 pM	12
Pb(II)	17–49 pM	200–500 pM	11
Ni(II)	1.7–4.3 nM	2–4 nM	17–19
Co(II)	10–103 pM	9–83 pM	11–16
Fe(III)	0.2–8 nM	0.4–13 nM	19–23

Table 2 Compilation of selected conditional stability constants and ligand concentrations of natural organic ligands for cobalt, nickel, copper, and zinc

Location	Values measured			Method, competing ligand ^{a,b}	Reference
	Metal concentration	Ligand concentration	$\log K'_{(\text{Metal})-L}$		
Cobalt	pM	pM			
Antarctic polar front	10–120	15–50	–	CSV, nioxime	(Ellwood et al. 2005)
Costa Rica upwelling dome	57–12 ^c 45–93 ^d	50	≥ 16.8	CSV, DMG	(Saito et al. 2005)
East equatorial Pacific	27–315 ^c	–	–	CSV, DMG	(Saito et al. 2004)
Atlantic Ocean, Sargasso Sea	17–73 ^d 20 \pm 10 ^{e,f} 19–133 ^c	–	–	CSV, DMG	(Saito & Moffett 2002)
Sargasso Sea	19–73 ^d	9–83	16.3 \pm 0.9	CSV, DMG	(Saito & Moffett 2001b)
Northeast Atlantic Ocean	25–103 ^c	22–60	15.6–16.1	CSV, nioxime	(Ellwood & van den Berg 2001)
Nickel	nM	nM			
Costa Rica upwelling dome	3.0 \pm 0.3 ^f	–	–	CSV, DMG	(Saito et al. 2005)
Coastal Britain	–	2–4	17.3–18.7	CSV, DMG	(van den Berg & Nimmo 1987)
Copper	nM	nM			
Subarctic northwest Pacific	3–4	3.7–5 (1500–2500 m)	12.7–14.1	CSV, salicylaldoxime, and benzoylacetone	(Moffett & Dupont 2007)
Estuarine waters	9–23 ^g	L ₁ = 10–33 L ₂ = 14–300	L ₁ = 14.8 – 15.8 L ₂ = 13–13.5	CSV, salicylaldoxime	(Laglera & van den Berg 2003)
North Pacific	0.58–1.88 ^d	L ₁ = 1.5–3 L ₂ = 5–10	L ₁ = 11.6 L ₂ = 8.6	DPASV	(Coale & Bruland 1990)
Zinc	nM	nM			
Northeastern Atlantic Ocean	0.3–2.0	0.4–2.5	10.0–10.5	CSV, PDC	(Ellwood & van den Berg 2000)
Central North Pacific	0.1–3.0 ^d	1.2	11.0	DPASV	(Bruland 1989)

Distributive Characteristics of Dissolved Trace Metals

- Oxyanions and big, standalone ions with one valence electron (like Cs⁺ and Rb⁺) tend to exist in small oceanic concentrations.
- Metals with greatest variation in oceanic concentration seem to be greatly related to biological processes (iron, zinc, cadmium ex.)

Types of Trace Metal Distributions:

- 1) Conservative Distributions: Trace metal retains relative concentration for long periods of time.
- 2) Nutrient-dependent distributions: metals that are involved in nutrient cycles and whose concentrations also depend on concentrations of other nutrients; includes zinc, cadmium, and (surprisingly) silver (and co.).
- 3) Scavenged-type distribution: scavenged very often; high concentrations near source (e.g. aluminum).

Some like iron and copper have hybrid distributions (where iron follows a nutrient-dependent and scavenged-type distribution).

Mixed distributions (where different forms of the metal undergo different distributions) also exist with elements like germanium.

--- the above is from Bruland 2003