

Lecture A2. Probability Review

Sim, Min Kyu, Ph.D., mksim@seoultech.ac.kr



서울과학기술대학교 데이터사이언스학과

1 I. Probability

2 II. Random Variables

3 III. Uniform

4 IV. Exponential

5 V. Poisson

6 VI. Some Exercises

I. Probability

Probability

Definition 1

Probability is a function from “Event” to real number between 0 and 1.

$$\mathbb{P} : S \rightarrow [0, 1],$$

where S is the set of all possible events.

Remark 1

- For any event E , $0 \leq \mathbb{P}(E) \leq 1$.
- For $E = \emptyset$, $\mathbb{P}(E) = 0$.
- $\mathbb{P}(S) = 1$, where S is whole space, or a space for all possible events.

Remark 2

- $\mathbb{P}(E_1 \cup E_2) = \mathbb{P}(E_1) + \mathbb{P}(E_2) - \mathbb{P}(E_1 \cap E_2)$.
- If $E_1 \cap E_2 = \emptyset$, then $\mathbb{P}(E_1 \cup E_2) = \mathbb{P}(E_1) + \mathbb{P}(E_2)$.

Conditional Probabilities

조건부 확률

Definition 2

$\mathbb{P}(E|F)$, "probability of E ^{bar}given F ", is the probability that event E occurs given that F has occurred. In a math notation,

$$\mathbb{P}(E|F) = \frac{\mathbb{P}(E \cap F)}{\mathbb{P}(F)}$$

- Ex) (from Ross) Suppose cards numbered one through ten are placed in a hat, mixed up, and then one of the cards is drawn. If we are told that the number on the drawn card is at least five, then what is the conditional probability that it is ten?

1. 10개 카드 중 1개 뽑을 경우의 수 $\Rightarrow \frac{1}{10}$
2. 뽑은 카드 한 개 중 5이상의 확률 $\Rightarrow \frac{6}{10} \left(\frac{2}{5} \right)$
3. 5이상의 카드 중 10인 확률 $\Rightarrow \frac{1}{10}$

$$\left[\frac{\frac{1}{10}}{\frac{6}{10}} \right] = \frac{10}{60} = \frac{1}{6}$$

Independent Probability

Definition 3

If $\mathbb{P}(E) = \mathbb{P}(E|F)$, then we say the events E and F are independent events.

- That is, the occurrence of E happening has nothing to do with the occurrence of F .
- Just as Galilei once said "*And yet it rotates (even though people do not believe so)*".
People believing whether or not the earth rotates is independent event of earth's rotation.

Properties

- ❶ $\mathbb{P}(E_1 \cup E_2) = \mathbb{P}(E_1) + \mathbb{P}(E_2)$, if $E_1 \cap E_2 = \emptyset$.
- ❷ $\mathbb{P}(E) = \mathbb{P}(E \cap F) + \mathbb{P}(E \cap F^c)$ ($\because (E \cap F) \cap (E \cap F^c) = \emptyset$)
- ❸ If $F_i \cap F_j = \emptyset$ for all $i \neq j$ and $\mathbb{P}(F_1 \cup \dots \cup F_n) = 1$ where $1 \leq i, j \leq n$, then $\mathbb{P}(E) = \sum_{i=1}^n \mathbb{P}(E \cap F_i)$

Bayes' rule

비이탈

$F_1 \cap E, F_2 \cap E$



Theorem 1

상호 배반 사건

Suppose F_1, \dots, F_n are **mutually exclusive events** (or, $F_i \cap F_j = \emptyset$, or, $\mathbb{P}(F_i \cap F_j) = 0$, for any $i \neq j$) and $\mathbb{P}(F_1 \cup \dots \cup F_n) = 1$ (In other words, exactly one and only one events among F_1, \dots, F_n will occur.), then the following holds.

$$\mathbb{P}(E) = \mathbb{P}(E \cap F_1) + \dots + \mathbb{P}(E \cap F_n)$$

$$= \sum_{i=1}^n \mathbb{P}(E \cap F_i)$$

$$= \sum_{i=1}^n \underbrace{\mathbb{P}(E|F_i)}_{\frac{\mathbb{P}(E \cap F_i)}{\mathbb{P}(F_i)}} \mathbb{P}(F_i)$$

E: 비가 올 확률

F_1 : 온도

ex) $F_1: 0 \sim 10^\circ\text{C}$ 0.3

$F_2: 10 \sim 20^\circ\text{C}$ 0.5

$$0.3 \times 0.7 + 0.1 = 0.3$$



Exercise 1

Show that $\mathbb{P}(A|B \cap C)\mathbb{P}(B|C) = \mathbb{P}(A \cap B|C)$.

$$\frac{\mathbb{P}(A \cap (B \cap C))}{\cancel{\mathbb{P}(B \cap C)}} \times \frac{\cancel{\mathbb{P}(B \cap C)}}{\mathbb{P}(C)} = \frac{\mathbb{P}(A \cap (B \cap C))}{\mathbb{P}(C)} = \frac{\mathbb{P}((A \cap B) \cap C)}{\mathbb{P}(C)} = \mathbb{P}(A \cap B|C)$$

II. Random Variables

discrete vs continuous

Discrete r.v. 이산형

Definition 4

A random variable X is called a discrete random variable if (*complete sentence*)

- Example
 - flip a coin)
 - throw a dice)

Continuous r.v. 연속형

Definition 5

A random variable X is called a continuous random variable if (*complete sentence*)

- Example
 - weights and heights of a person)
 - temperature of a room)

pmf and pdf

확률 질량 함수
pmf(probability mass function) for a discrete r.v. X

Definition 6

A pmf $p(x)$ is a function that gives the **probability** that a discrete random variable is exactly equal to some value.

$$\mathbb{P}(X = x) = p(x)$$

- Suppose you throw a dice and X be a r.v. of the outcome. What is the pmf of X ?

① answer in a math form: $P(X) = \frac{1}{6} \quad x=1, 2, 3 \dots, 6$

② answer in a tabular form:

결과	1	2	3	4	5	6
확률	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$

pdf(probability density function) for a continuous r.v. X

정의 *정의* *정의*

Definition 7


A pdf $f(x)$ is a function that gives the **relative likelihood** for this continuous r.v. to take on a given value.

$$\mathbb{P}(a \leq X \leq b) = \int_a^b f(x) dx$$

- Suppose a random variable takes a real values between 0 and 1 with equal likelihood, what is the pdf of X ?

① answer in a math form: $P(0 \leq X \leq 1) = \int_0^1 f(x) dx$

② draw a graph:



Properties of pmf and pdf

AE

- Mathematical property? *Something that always happen.*

Remark 3

The functions (pmf and pdf) are nonnegative everywhere. That is,

$$p(x) \geq 0 \text{ and } f(x) \geq 0$$

Remark 4

Its summation or integral over the entire area is equal to one. That is,

$$\sum_{\text{pmf}} p(x) = 1 \text{ and } \int_{\text{pdf}} f(x) dx = 1$$

- One can derive a cdf (cumulative distribution function) from a pmf or from a pdf. (See the upcoming definition of cdf.)

Expectation

$$\text{이산형} \quad \text{Var}(X) = E(X^2) - (E(X))^2$$

x	1	2	3	4	5	6
$p(x)$	$\frac{1}{6}$					$\frac{1}{6}$

- For a discrete random variable X with pmf $p(x) \sum_{n=1}^6 x_n p(x)$

- $EX = \sum xp(x)$
- $E[g(X)] = \sum g(x)p(x)$
- $Ex) EX^2 = \sum x^2 p(x)$
- $Ex) E(X^2 - 2X) = \sum$

- 연속형 For a continuous random variable X with pdf $f(x)$

- $EX = \int_{-\infty}^{\infty} xf(x)dx$
- $E[g(X)] = \int_{-\infty}^{\infty} g(x)f(x)dx$
- $Ex) EX^2 = \int_{-\infty}^{\infty} x^2 f(x)dx$

cdf 누적분포함

Definition 8

For a random variable X , the cdf(cumulative distribution function) $F(x)$ is a function of probability that the random variable X is found at a value less than or equal to x .

$$F(x) := \mathbb{P}(X \leq x) \quad \text{확률, 스 등.}$$

- If discrete,

$$F(x) = \mathbb{P}(X \leq x) = \sum_{y=-\infty}^x \mathbb{P}(X = y) = \sum_{y=-\infty}^x p(y)$$

- If continuous,

$$F(x) = \mathbb{P}(X \leq x) = \int_{-\infty}^x f(y) dy$$

III. Uniform

Definition 9

A continuous random variable X is said to follow **uniform distribution** with parameter a and b , and write $X \sim U(a, b)$ if (reads "X follows a uniform distribution with parameter a and b .")

확률 밀도


$$\text{pdf } \underline{f(x)} = \begin{cases} \frac{1}{b-a} & \text{if } a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$$

$$\text{cdf } \underline{F(x)} = \begin{cases} 0 & \text{if } x \leq a \\ \frac{x-a}{b-a} & \text{if } a \leq x \leq b \\ 1 & \text{if } x > b \end{cases}$$


✓ (The value $\frac{1}{b-a}$ is chosen as a constant so that it will make integrations over $-\infty$ to ∞ to be equal to 1.)

$U(0, 1)$

• pdf



• cdf



Exercise 2

$\nearrow^a \nearrow^b$

$\nwarrow^{\cancel{x}} \nearrow^x$

$$f(x) = \frac{1}{b-a} = \frac{1}{20-10} = \frac{1}{10}$$

$X \sim U(10, 20)$, then what is $F(10)$? and $F(15)$?

$$10 \leq x \leq 20$$

0 otherwise

$$F(10) : x \leq 10 \Rightarrow 0$$

$$10 \leq x \leq 20 \Rightarrow \frac{x-a}{b-a}$$

$$20 < x \Rightarrow 1$$

$$= \frac{10-10}{20-10}$$

$$= \frac{0}{10}$$

$$= 0$$

$$F(15) : x \leq 10 \Rightarrow 0$$

$$10 \leq x \leq 20 \Rightarrow \frac{x-a}{b-a}$$

$$20 < x \Rightarrow 1$$

$$= \frac{15-10}{20-10}$$

$$= \frac{5}{10}$$

$$= 0.5$$




IV. Exponential

포아송분포

$$P(x) = \frac{\lambda^x \cdot e^{-\lambda}}{x!}$$

λ : 단위시간당
 사건의 평균 발생 횟수
 $x!$: 사건 발생 횟수

⇒ 길잡이를 기다릴 때까지 걸리는 시간이 T 이하일 확률



Definition 10

A nonnegative continuous random variable X is said to follow **exponential distribution** with parameter λ and write $X \sim \text{exp}(\lambda)$, if

ex) $X \sim \text{exp}(3)$


3분마다 한 번씩.

$$\text{pdf } f(x) = \begin{cases} \lambda e^{-\lambda x}, & \text{if } x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$


λ 시간 안에 X 가 생길 확률

$$\text{cdf } F(x) = \begin{cases} 1 - e^{-\lambda x}, & \text{if } x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

● pdf



● cdf



- TFAE (The followings are equivalent - If any one of them is true, the rest of them are all true. If any one of them is false, then the rest of them are all false.)

① $X \sim \exp(\lambda)$

② pdf $f(x) = \lambda e^{-\lambda x}$, if $x \geq 0$; $f(x) = 0$, otherwise

③ cdf $F(x) = 1 - e^{-\lambda x}$, if $x \geq 0$; $F(x) = 0$, otherwise

Exercise 3

Prove that pdf \rightarrow cdf

$$\int_{-\infty}^x f(x) dx = F(x)$$

$$\text{if } x \geq 0$$

$$\int_0^x \lambda e^{-\lambda x} dx = [-e^{-\lambda x}]_0^x$$

$$= -e^{-\lambda x} - (-1)$$

$$= 1 - e^{-\lambda x}$$

$$e^{2x} = 1 e^{2x} \\ e^{x^2}$$

More statistics (통계량) for r.v.

- Variance: $Var(X) = \mathbb{E}X^2 - (\mathbb{E}X)^2$
- Standard Deviation: $sd(X) = \sqrt{Var(X)}$
- Coefficient of Variation (cv): $cv(X) = \frac{sd(X)}{\mathbb{E}X}$
- (A cv measures 'relative' variation and often more meaningful.)
- Examples
 - ① cv of $N(10, \sqrt{10}^2)$ vs $N(10000, 10^2)$. $\frac{\sqrt{10}}{10} \searrow \frac{1}{1000}$
 - ② For an exponential r.v., $X \sim \exp(\lambda)$, then $cv(X) = \frac{1}{\lambda}$ (next slide)
 - ③ For a normal r.v. $X \sim N(\mu, \sigma^2)$, $cv(X) = \frac{\sigma}{\mu}$
 - ④ For deterministic variable X , $c_x = 0$
 $\underbrace{\quad}_{\parallel}$
 $CV(X)$

Properties of exponential distribution

- Statistics for a r.v. $X \sim \exp(\lambda)$,

71% ① $EX = 1/\lambda$

45% ② $Var(X) = 1/\lambda^2$ $\sqrt{Var(X)} = SD(X) = \frac{1}{\lambda}$

③ $cv(X) = 1$ (Exponential r.v. has c.v. equal to 1, always.)

0 ~ 2 한생 x

0 ~ 3 = 1 ~ 5 한생이 1인

- Theorems

① Memoryless property $X \sim \exp(\lambda)$ $\rightarrow P(X > t+s | X > t) = P(X > s)$

② Suppose $X_1 \sim \exp(\lambda_1)$, $X_2 \sim \exp(\lambda_2)$, and independent, then

$$P(X_1 < X_2) = \frac{\lambda_1}{\lambda_1 + \lambda_2}$$

③ If $X_1 \sim \exp(\lambda_1)$, $X_2 \sim \exp(\lambda_2)$, and independent, then

$$\min(X_1, X_2) \sim \exp(\lambda_1 + \lambda_2)$$

Exercise 4

Show that $EX = 1/\lambda$

$$E(x) = \int_0^{\infty} x \cdot f(x) dx = \int_0^{\infty} \underbrace{x \lambda e^{-\lambda x}}_{\text{density}} dx \quad (\text{density})$$

$$\text{if } x \geq 0$$

$$= [-x e^{-\lambda x}]_0^{\infty} - \int_0^{\infty} -e^{-\lambda x} dx$$

$$= [-x \cdot e^{-\lambda x}]_0^{\infty} + [-\frac{1}{\lambda} e^{-\lambda x}]_0^{\infty}$$

$$= \lim_{x \rightarrow \infty} (-x \cdot e^{-\lambda x}) - 0 + \lim_{x \rightarrow \infty} (-\frac{1}{\lambda} e^{-\lambda x}) - (-\frac{1}{\lambda} e^{-\lambda \cdot 0})$$

\nearrow
 $\underbrace{\quad}_{= +\frac{1}{\lambda}}$

$$= \lim_{x \rightarrow \infty} (-x \cdot e^{-\lambda x}) + \frac{1}{\lambda}$$

$$= -\lim_{x \rightarrow \infty} \left(\frac{x}{e^{\lambda x}} \right) + \frac{1}{\lambda}$$

$$\text{L'Hôpital's rule} \rightarrow -\frac{1}{\lambda} \left(\lim_{x \rightarrow \infty} \frac{\lambda x}{e^{\lambda x}} \right) + \frac{1}{\lambda} = \frac{1}{\lambda}$$

Exercise 5

Show that $\text{Var}(X) = 1/\lambda^2$. (Hint: need to do EX^2 first)

$$\text{Var}(X) = E(X^2) - \underbrace{\{E(X)\}^2}_{\frac{1}{\lambda^2}}$$

$$\text{Var}(X) = E(X^2) - \frac{1}{\lambda^2}$$

$$E(X) = \int_0^{\infty} x f(x) dx$$

$$E(X^2) = \int_0^{\infty} x^2 \cdot \underline{\lambda e^{-\lambda x}} dx = [x^2(-e^{-\lambda x})]_0^{\infty} - \int_0^{\infty} 2x \cdot (-e^{-\lambda x}) dx \quad (\text{부분적분})$$

$$= [x^2 \cdot (-e^{-\lambda x})]_0^{\infty} + \frac{2}{\lambda} \cdot \boxed{\int_0^{\infty} x \cdot e^{-\lambda x} dx}$$

$$\hookrightarrow E(X) = \frac{1}{\lambda}$$

$$= [x^2 \cdot (-e^{-\lambda x})]_0^{\infty} + \frac{2}{\lambda^2}$$

$$= \lim_{x \rightarrow \infty} x^2 \cdot (-e^{-\lambda x}) + \frac{2}{\lambda^2}$$

$$= \left(\lim_{x \rightarrow \infty} \frac{x^2}{e^{\lambda x}} \right) + \frac{2}{\lambda^2} \rightarrow 0$$

Common usage of exponential distribution

- Since exponential r.v. is continuous and nonnegative, it is frequently used to describe time.
- In such cases, the parameter λ can be understood as "rate" as an inverse value of the time.
- For example, $X \sim \text{exp}(\lambda)$ and $\mathbb{E}X = 1/\lambda = 5 \text{ yrs}$ imply $\lambda = 1/5$ per year.
- For example, consider the series of events that occur in every time interval that follows exponential distribution with mean 3 minutes. (In post office, you expect one customer arrival in 3 minutes on average and the time follows exponential distribution). We say that it follows exponential arrival times with rate 1/3 per minute.

$2 \sim 5 \text{ 시간}$ $2 \sim 5 \text{ 분}$

$0 \sim 3 \text{ 시간}$ $0 \sim 3 \text{ 분}$

Memoryless property 무기억성

$$0 \sim 2$$

$$2 \sim 5 = 0 \sim 3$$

Definition 11

A r.v. X is *memoryless*, if $\mathbb{P}(X > s + t | X > t) = \mathbb{P}(X > s)$, for $s, t \geq 0$.

Theorem 2

Exponential random variable is memoryless.

- For example, we shall assume that bus arrival time follows $\exp(1/5)$. In other words, it follows an exponential distribution and its expected arrival time is 5 minutes. You are waiting for bus, and have been waiting for 3 minutes, what is the probability that bus will not come in 5 minutes from now.

$$0 \sim 3 \text{ 안 올 } , \quad 0 \sim 5 \text{ 을 확률 같음}$$

$$3 \sim 8$$

Exercise 6

Prove the previous theorem.

$$P(X > s+t | X > t) = \frac{P(X > s+t \text{ and } X > t)}{P(X > t)} = \frac{P(X > s+t)}{P(X > t)}$$

$$\begin{aligned} \textcircled{1} \quad P(X > s+t) &= 1 - P(X \leq s+t) \\ &= 1 - F(s+t) \\ &= 1 - (1 - e^{-\lambda(s+t)}) \\ &= e^{-\lambda(s+t)} \end{aligned}$$

$$\begin{aligned} \textcircled{2} \quad P(X > t) &= 1 - P(X \leq t) \\ &= 1 - F(t) \\ &= 1 - (1 - e^{-\lambda t}) \\ &= e^{-\lambda t} \end{aligned}$$

$$\frac{e^{-\lambda(s+t)}}{e^{-\lambda t}} = e^{-\lambda s} = P(X > s)$$

한 번 일의 수를 지어 걸리는 시간

Theorem 3

Suppose $X_1 \sim \exp(\lambda_1)$, $X_2 \sim \exp(\lambda_2)$ and *they are independent*, then

$$\mathbb{P}(X_1 < X_2) = \frac{\lambda_1}{\lambda_1 + \lambda_2}$$

- Smith and Jones came to post office together and they are served by two clerks, A and B, respectively. Server A has service time following $\exp(1/3)$ and server B has service time following $\exp(1/5)$. What is the chance that Smith will be done first?

$$\frac{\frac{1}{3}}{\frac{1}{3} + \frac{1}{5}} = \frac{\frac{1}{3}}{\frac{8}{15}} = \frac{15}{24}$$

- Suppose that Smith came to post office earlier than Jones by 2 minutes, but Smith was still being served at the moment that Jones started to being served. Would this assumption change your previous answer? Why or why not?

Memoryless

변하지 않습니다. Smith는 사건이 발생하지 않았기 때문에
자신의 과거의 정보에 영향을 주지 않습니다.

V. Poisson

Definition 12

시간안에 일어난 횟수.

A discrete random variable X is said to follow Poisson distribution with parameter λ , and write $X \sim \text{poi}(\lambda)$, if its pmf is

$$\mathbb{P}(X = k) = \frac{\lambda^k e^{-\lambda}}{k!} \text{ for } k = 0, 1, 2, \dots$$

- What is cdf of $\text{poi}(\lambda)$? $\sum_{k=0}^x \frac{\lambda^k e^{-\lambda}}{k!}$

Remark 5

$$\mathbb{E}X = \text{Var}(X) = \lambda$$

기대값 = 분산 = λ

Exercise 7

For $X \sim \text{poi}(\lambda)$, prove that $\mathbb{E}X = \lambda$.

- cf) $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots = \sum_{n=0}^{\infty} \frac{x^n}{n!}$
- pf) We have $\mathbb{P}(X = k) = \frac{\lambda^k e^{-\lambda}}{k!}$ for $k = 0, 1, 2, \dots$, and

$$\begin{aligned}
 \mathbb{E}X &= \sum_{x=-\infty}^{\infty} xp(x) \text{ (this is common for all discrete r.v.)} \\
 &= \sum_{k=0}^{\infty} k \frac{\lambda^k e^{-\lambda}}{k!} \text{ (x=0을 넣으면 전체 항이 0)} \\
 &= \sum_{k=1}^{\infty} k \frac{\lambda^k e^{-\lambda}}{k!} \\
 &= \lambda e^{-\lambda} \sum_{k=1}^{\infty} \frac{\lambda^{k-1}}{(k-1)!} \\
 &= \lambda e^{-\lambda} \sum_{n=0}^{\infty} \frac{\lambda^n}{n!} \text{ (k-1을 n으로 치환)} \\
 &= \lambda e^{-\lambda} e^{\lambda} \\
 &= \lambda
 \end{aligned}$$

VI. Some Exercises

Maximum and minimum

Definition 13

$$x \wedge y = \min(x, y) = \begin{cases} x & , \text{ if } x \leq y \\ y & , \text{ otherwise} \end{cases}$$

$$x \vee y = \max(x, y) = \begin{cases} x & , \text{ if } x \geq y \\ y & , \text{ otherwise} \end{cases}$$

$$x^+ = \max(x, 0) \text{ (positive part)}$$

- Ex) $x \wedge 25 = \min(x, 25)$
- Ex) $(25 - x)^+ = \max(25 - x, 0)$

$$\mathbb{E}[U] = \frac{1}{b-a}$$

Exercise 8

For $X \sim U(20, 40)$, evaluate $\mathbb{E}[X \wedge 25]$ and $\mathbb{E}[(25 - X)^+]$. $x^+ = \max(x, 0)$ (positive part)

$$X \wedge 25 \begin{cases} X & (X \leq 25) \\ 25 & (X > 25) \end{cases}$$

$$\begin{aligned} \mathbb{E}[X \wedge 25] &= \int_{-\infty}^{\infty} (X \wedge 25) f_X(x) dx \\ &= \int_{20}^{25} (X \wedge 25) \frac{1}{20} dx + \int_{25}^{40} (X \wedge 25) \frac{1}{20} dx \\ &= \int_{20}^{25} x \left(\frac{1}{20}\right) dx + \int_{25}^{40} 25 \cdot \frac{1}{20} dx \\ &= \left[\frac{1}{40} x^2 \right]_{20}^{25} + \left[\frac{5}{4} x \right]_{25}^{40} \\ &= \frac{1}{40} (25^2 - 20^2) + \frac{5}{4} (40 - 25) \\ &= \frac{225}{40} + \frac{75}{4} \\ &= \frac{300}{40} \\ &= \frac{15}{2} \end{aligned}$$

$$(25 - X)^+ = \max(25 - X, 0)$$

$$\max(25 - X, 0) \begin{cases} 25 - X & (25 \geq X) \\ 0 & (25 < X) \end{cases}$$

$$\begin{aligned} \mathbb{E}[(25 - X)^+] &= \int_{-\infty}^{\infty} (25 - X)^+ f_X(x) dx \\ &= \int_{20}^{25} (25 - X)^+ \frac{1}{20} dx \\ &= \int_{20}^{25} (25 - X)^+ \frac{1}{20} dx + \int_{25}^{40} (25 - X)^+ \frac{1}{20} dx \\ &= \int_{20}^{25} 0 \cdot \frac{1}{20} dx + \int_{25}^{40} (25 - X) \cdot \frac{1}{20} dx \\ &= 0 + \frac{1}{20} \left[25X - \frac{1}{2} X^2 \right]_{25}^{40} \\ &= \frac{1}{20} \left[\left(25 \cdot 40 - \frac{1}{2} 40^2 \right) - \left(25 \cdot 25 - \frac{1}{2} 25^2 \right) \right] \\ &= \frac{1}{20} \left[\left(200 - \frac{800}{2} \right) - \left(625 - \frac{625}{2} \right) \right] \\ &= \frac{1}{20} \left[\left(200 - 400 \right) - \left(625 - 312.5 \right) \right] \\ &= \frac{1}{20} \left[-200 - 312.5 \right] = -\frac{45}{8} \end{aligned}$$

이산

Exercise 9

↗ $\lambda = 8$ For $X \sim \text{Poi}(8)$,

① $\mathbb{P}(X = 0) =$

② $\mathbb{P}(2 \leq X \leq 4) =$

③ $\mathbb{P}(X > 2) =$

$$\mathbb{P}(X = k) = \frac{\lambda^k e^{-\lambda}}{k!} \text{ for } k = 0, 1, 2, \dots$$

$$1. \mathbb{P}(X=0) = \frac{8^0 e^{-8}}{0!} = \frac{1 \times e^{-8}}{1} = e^{-8}$$

$$3. \mathbb{P}(X > 2) = 1 - \underbrace{\mathbb{P}(X=0)}_{e^{-8}} - \underbrace{\mathbb{P}(X=1)}_{8e^{-8}} - \underbrace{\mathbb{P}(X=2)}_{32e^{-8}}$$

$$= 1 - 25e^{-8}$$

$$2. \mathbb{P}(2 \leq X \leq 4) = \mathbb{P}(X=2) + \mathbb{P}(X=3) + \mathbb{P}(X=4)$$

$$\mathbb{P}(X=2) = \frac{8^2 e^{-8}}{2!} = \frac{32 \times e^{-8}}{2} = 16e^{-8}$$

$$\mathbb{P}(X=3) = \frac{8^3 e^{-8}}{3!} = \frac{512 \times e^{-8}}{6} = \frac{256}{3} e^{-8}$$

$$\mathbb{P}(X=4) = \frac{8^4 e^{-8}}{4!} = \frac{4096 \times e^{-8}}{24} = \frac{512}{3} e^{-8}$$

$$\therefore \frac{16}{3} e^{-8} + \frac{256}{3} e^{-8} + \frac{512}{3} e^{-8}$$

$$= \frac{864}{3} e^{-8}$$

$$= 288 e^{-8}$$

Exercise 10 *nonnegative random variable*

$$\text{자살률} = \frac{1}{\lambda}$$

For $X \sim \exp(7)$, evaluate $\mathbb{E}[\max(X, 7)]$.

$$\begin{aligned}
 \mathbb{E}[\max(X, 7)] &= \int_{-\infty}^{\infty} (x \vee 7) f_X(x) dx & \max(X, 7) = \begin{cases} x & (x \geq 7) \\ 7 & (x < 7) \end{cases} \\
 &= \int_0^7 (x \vee 7) \cdot 7e^{-7x} dx + \int_7^{\infty} (x \vee 7) \cdot 7e^{-7x} dx \\
 &= \int_0^7 7 \cdot 7e^{-7x} dx + \int_7^{\infty} x \cdot 7e^{-7x} dx \\
 &= 7 \int_0^7 e^{-7x} dx + 7 \int_7^{\infty} x e^{-7x} dx \\
 &= 7 \left[-\frac{1}{7} e^{-7x} \right]_0^7 + 7 \left[-\frac{1}{7} x e^{-7x} - \frac{1}{49} e^{-7x} \right]_7^{\infty} \\
 &= -e^{-49} - 0 + 7e^{-49} \\
 &= 6e^{-49}
 \end{aligned}$$

$$\text{pdf } f(x) = \begin{cases} \lambda e^{-\lambda x}, & \text{if } x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$\text{cdf } F(x) = \begin{cases} 1 - e^{-\lambda x}, & \text{if } x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$x \wedge y = \min(x, y) = \begin{cases} x, & \text{if } x \leq y \\ y, & \text{otherwise} \end{cases}$$

$$x \vee y = \max(x, y) = \begin{cases} x, & \text{if } x \geq y \\ y, & \text{otherwise} \end{cases}$$

$$x^+ = \max(x, 0) \text{ (positive part)}$$

Exercise 11

For $X \sim \text{exp}(8)$, find x^* such that $F(x^*) = 0.6$.

$$\ln = \log_e.$$

$$F(x) = 1 - e^{-\lambda x}$$

$$\text{cdf } F(x) = \begin{cases} 1 - e^{-\lambda x}, & \text{if } x \geq 0 \\ 0, & \text{otherwise} \end{cases}$$

$$F(x^*) = 1 - e^{-8x^*} = 0.6$$

$$= e^{-8x^*} = 0.4$$

$$= \log_e e^{-8x^*} = \log_e 0.4$$

$$-8x^* = \log_e 0.4$$

$$x^* = -\frac{1}{8} \log_e 0.4$$

Exercise 12



For $X \sim U(10, 20)$, find x^* such that $\underline{F}(x^*) = 0.7$.

$$F(x) = \frac{x-a}{b-a}$$

$$F(x^*) = \frac{x^*-a}{b-a}$$

$$= \frac{x^*-10}{20-10} = 0.7$$

$$= x^*-10 = 7$$

$$= x^* = 17$$

"Man can learn nothing unless he proceeds from the known to the unknown. - Claude Bernard"