

**LA TOXICITY**

All local anaesthetics given must not exceed the standard toxic limits:

LIGNOCAINE (plain)	3- 4 mg/kg
LIGNOCAINE ET ADRENALINE 1:200 000	7 mg/kg
BUPIVACAINE	2.0- 2.5 mg/kg (children) 1.5- 2.0 mg/kg(neonates)
LEVOBUPIVACAINE	2.0- 2.5 mg/kg
ROPIVACAINE	2.5- 3.0 mg/kg

Treatment guidelines for serious systemic toxicity of local anesthetics with 20% lipid emulsion by Association of Anaesthetists of Great Britain and Ireland (AAGBI)- see next page.


\*In KKH, Lipofundin can be found in the black LAST boxes at MOT recovery and outside DSOT 2.

Otherwise call Pharmacy 3086/2464 or IPAS 2209/8474.


Pharmacist: on-call 91690643.

It should be given as soon as it is available. Normal resuscitative measures should be instituted in the meantime.

# PAEDIATRIC ANAESTHESIA



**An approximate dose regimen for a 70-kg patient would be as follows:**



This AAGBI Safety Guideline was produced by a Working Party that comprised:  
Grant Cave, Will Harrop-Griffiths (Chair), Martyn Harvey, Tim Meek, John Picard, Tim Short and Guy Weinberg.

**This Safety Guideline is endorsed by the Australian and New Zealand College of Anaesthetists (ANZCA).**