

Extra Dimensions & Gravitons

FYS4560

Sean B.S. Miller

University of Oslo

June 6, 2016

Introduction

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

There are 3 different types of spatial extra dimensions (ED) models.

- *Large*
- *Warped*
- *Universal*

Arkani-Hamed, Dimopoulos, and Dvali (1998) proposed the ADD-model¹ (large ED)

Randall and Sundrum (1999) proposed RS1-model² (warped ED)

¹arXiv: hep-ph/9803315

²arXiv: hep-ph/9905221

Kaluza-Klein theory

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

From linearised gravity,

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad (1)$$

we can find an interaction term $h_{\mu\nu} T^{\mu\nu}$ and a kinetic term for $h_{\mu\nu}$ in the Lagrangian density. However, a mass term for $h_{\mu\nu}$ can be suggested as

$$a h_{\mu\nu} h^{\mu\nu} + b (\eta_{\mu\nu} h^{\mu\nu})^2 \quad (2)$$

Markus Fierz and Wolfgang Pauli (1939) showed[4] that $a = -b$ in order to avoid unphysical results. The result of this was the Fierz-Pauli Lagrangian for massive gravity.

Kaluza-Klein theory

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

Kaluza assumed the metric could be written:

$$\hat{h}_{ab} = V_d^{-1/2} \begin{pmatrix} h_{\mu\nu} + \eta_{\mu\nu}\phi & A_{\mu i} \\ A_{\nu j} & 2\phi_{ij} \end{pmatrix} \quad (3)$$

Kaluza assumed that these fields could be written as expansions, i.e.

$$h_{\mu\nu}(x, y) = \sum_{\substack{n=\{n_1, n_2, \dots, n_5\}; \\ n_i \in \mathbb{Z} \forall i}} h_{n, \mu\nu}(x) Y_n(y) \quad (4)$$

and similarly for $A_{\mu i}$ and ϕ . The modes have to satisfy the Fierz-Pauli equations of motion. These, when combined, will show that $h_{\mu\nu}$ satisfies:

$$(\square + m_n^2)(h_{n, \mu\nu} - \frac{1}{2}\eta_{\mu\nu}h_{n, \sigma}^{\sigma}) = 0 \quad (5)$$

where $m_n^2 = \frac{4\pi^2 n^2}{R^2}$.

Kaluza-Klein theory

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

Through some non-trivial steps, one can find a Lagrangian with mass eigenstates $\tilde{h}_{n,\mu\nu}$, $\tilde{A}_{n,\mu i}$, and $\tilde{\phi}_{n,ij}$, defined from the previous fields. From the Hilbert-Einstein action:

$$S_d = \int d^4x \sqrt{-\hat{g}} \mathcal{L}(\hat{g}, S, V, F) \quad (6)$$

where $\hat{g}_{\mu\nu} = \eta_{\mu\nu} + \kappa(h_{\mu\nu} + \eta_{\mu\nu}\phi_{ii})$, we can find Feynman rules:

$$G_n^{\mu\nu\rho\sigma} = i \frac{\eta^{\mu\rho}\eta^{\nu\sigma} + \eta^{\mu\sigma}\eta^{\nu\rho} - \frac{2}{3}\eta^{\mu\nu}\eta^{\rho\sigma}}{k_G^2 - m_n^2 + i\epsilon} \quad (7)$$

$$-\frac{i\kappa}{8} [\gamma_\mu(k_1 + k_2)_\nu + \gamma_\nu(k_1 + k_2)_\mu - 2\eta_{\mu\nu}(\not{k}_1 + \not{k}_2 - 2m_f)] \quad (8)$$

ADD-model

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

The ADD model is mainly based on 3 features:

- There exists d new spatial compact dimensions, with compactification volume V_d .
- The Planck scale is very low, at the order of one TeV,
- The SM degrees of freedom are localized on a 3D-brane, stretching along 3 non-compact spatial dimensions (i.e. the SM particles move in normal spacetime, not in the new dimension(s)).

The main idea is that $\bar{M} \sim 1\text{TeV}$. By demanding $S_4 = S_{4+d}$, one finds the reduction formula:

$$M_{Pl}^2 = V_d \bar{M}^{d+2} \sim R^d \bar{M}^{d+2} \quad (9)$$

ADD-model

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

With $M_{Pl} \approx 1.22 \times 10^{16}$ TeV, one finds:

$$d = 1 \quad \rightarrow \quad R \sim 10^{11} \text{ m}$$

$$d = 2 \quad \rightarrow \quad R \sim 0.1 \text{ mm}$$

$$d = 3 \quad \rightarrow \quad R \sim 10^{-7} \text{ mm}$$

...

$$d = 6 \quad \rightarrow \quad R \sim 10^{-11} \text{ mm}$$


But Newton's law of gravity must still hold for $r \gg R$.

RS1-Model

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

RS1-model is quite different. They assumed space had S^1/\mathbb{Z}^2 "orbifold" structure. At two points there existed a 3-brane:



From Poincare invariance, it was found:

$$ds^2 = e^{-2A(y)} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2 \quad (10)$$

Due to \mathbb{Z}^2 -symmetry, have $A(y) = A(-y) = A(|y|)$. Solving the Einstein equations let Randall and Sundrum find $A(y) = k|y|$,

$$k = \sqrt{\frac{-\Lambda}{24M^3}}.$$

RS1-Model

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

The metric forces a new reduction formula:

$$M_{Pl}^2 = \frac{M_5^3}{k} [1 - e^{-2kR\pi}] \quad (11)$$

For M_5 to solve the hierarchy problem, one has to require $kR \sim 12 \rightarrow e^{k\pi R} \sim 10^{15}$.

With a new metric form, one has to repeat the KK mode expansion:

$$h_{\mu\nu}(x, \theta) = \sum_{n=0}^{\infty} h_{\mu\nu}^{(n)}(x) \frac{\chi_n(\theta)}{R} \quad (12)$$

where $\chi_0(y) = 2\sqrt{kR}e^{-2kR|\theta|}$, and

$$\chi_n(y) = N_n \left[C_1 Y_2 \left(\frac{m_n}{k} e^{kR|\theta|} \right) + C_2 J_2 \left(\frac{m_n}{k} e^{kR|\theta|} \right) \right], \quad n \neq 0 \quad (13)$$

RS1-model

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

On TeV brane, KK mode masses are

$$m_n = \beta_n k e^{-kR\pi}, \quad J_1(\beta_n) = 0 \quad \forall n \in \mathbb{N} \quad (14)$$

Unlike the ADD-model, the masses are clearly separated. This means the sum over the KK tower is not necessary.

Comparing RS1 Hilbert-Einstein action with the general KK action, one will see that:

$$\kappa = \sqrt{2} \frac{\beta_1}{m_1} \frac{k}{M_{Pl}} \quad (15)$$


meaning RS1 is completely determined by to constants.

Proton-proton collisions

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

One channel for measuring the models is through the
 $pp \rightarrow G \rightarrow l^+ l^-$ ($q\bar{q} \rightarrow G \rightarrow l^+ l^-$):



Proton-proton collisions

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

After a little math...


Proton-proton collisions

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

$$\frac{d\sigma_G}{d\cos(\theta)} = \frac{1}{2}\pi Q_f^2 s^6 |C_4|^2 [1 - 3\cos^2(\theta) + 4\cos^4(\theta)] \quad (16)$$

where $C_4 = \frac{\kappa^2}{8\pi} D(s)$ and $D(s) \equiv \sum_n \frac{1}{k_G^2 - m_n^2 + i\epsilon}$.



Gravitons at the LHC

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

Conclusions

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

References

Extra Dimensions
& Gravitons
FYS4560

Sean B.S.
Miller

-  E.W. Dvergsnes, " *Extra Dimensions at Particle Colliders*", 2004, UiB
-  T. Han, J.D. Lykken, and R.-J. Zhang, *Phys. Rev. D* **59** (1999) 105006, doi:10.1103/PhysRevD.59.105006 hep-ph/9811350v4
-  B.C. Allanach, K. Odagiri, M.A. Parker, and B.R. Webber, *JHEP* 0009:019,2000, doi:10.1088/1126-6708/2000/09/019, hep-ph/0006114v2
-  M. Fierz, P. Wolfgang, *Proc. Roy. Soc. Lond. A*173: 211232, doi: 10.1098/rspa.1939.0140
-  E.E. Boos, Y.A. Kubyshin, M.N. Smolyakov, and I.P. Volobuev, *NPI MSU* 2001-21/661, hep-th/0105304
-  H. Davoudiasl, J.L Hewett, and T. G. Rizzo, *Phys. Rev. Lett.* **84** 3370 (1999), doi:10.1103/PhysRevLett.84.2080, hep-ph/9911262v2
-  Y.A. Kubyshin, (2001), hep-ph/0111027v2