

1	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
0	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Registration number

Signature

- Note:**
- Cross your Registration number(with leading zero). It will be evaluated automatically.
 - Sign in the corresponding signature field.

Discrete Mathematics for Engineers

Exam:	EDE1202 / Endterm	Date:	Tuesday 23 rd February, 2021
Examiner:	Prof. Dr. Ing. Ulf Schlichtmann	Time:	16:30 – 18:00

Problem 1 Propositional Logic (8 credits)

Given is the ring normal form $RNF[A]$ of the propositional form $A(x, y)$.

$$RNF[A(x, y, z)] \iff x \longleftrightarrow y \longleftrightarrow t$$

0		
1		
2		

a)* Give the ring normal form $RNF[dual(A)]$ of the dual form of A .

[illegible]

0		
1		
2		

b)* Give the canonical disjunctive normal form $CDNF[A]$ of A .

[illegible]

0		
1		

c)* Give the value pattern $\underline{\hat{W}}[A]$ of A .

[illegible]

0		
1		
2		
3		

d)* Give all non-trivial conclusions of A .


[illegible]

Problem 2 Propositional Logic (5 credits)

Prove the following implication by means of the deductive proof scheme:

$$(a \longleftrightarrow c) \implies (b \longrightarrow a) \longleftrightarrow (b \longrightarrow c)$$

Name all laws you use.



Row		From Row(s)	Law(s)
1	$a \longleftrightarrow c \implies (b \longrightarrow a) \longleftrightarrow (b \longrightarrow c)$		
2			
3			
4			
5			
6			
7			
8			
9			
10			
11			
12			
13			
14			
15			
16			
17			
18			

Problem 3 Propositional Logic (5 credits)

a)* Given is the following propositional form $B(a, b, c, d)$:

$$B(a, b, c, d) \iff (a \vee \neg b) \wedge (c \vee d) \wedge (\neg a \vee \neg d) \wedge (b \vee \neg d) \wedge \neg c.$$

Prove by means of the layer algorithm that $B(a, b, c, d)$ is a contradiction.

$(a \vee \neg b) \wedge (c \vee d) \wedge (\neg a \vee \neg d) \wedge (b \vee \neg d) \wedge \neg c$	Layer
	1
	2
	3
	4
	5
	6

b)* Additionally, the propositional form $C(a, b, c)$ is given:


$$C(a, b, c) \iff c \vee (a \wedge b)$$

Is the implication $B(a, b, c, d) \implies C(a, b, c)$ always valid? Justify your answer.

[illegible]

Problem 4 Propositional Logic (6 credits)

Given is the *ROBDD*[C] of the propositional form $C(x, y, z)$:



- a)* Give the premise normal form $PNF[C]$ of the propositional form $C(x, y, z)$.

[illegible]

0
1

- b)* Give the number of t- and f-assignments in the propositional form $C(x, y, z)$.

[illegible]

	0
	1
	2

- c)* Draw the $ROBDT[C]$ of the propositional from $C(x, y, z)$ with the **variable ordering** $\mathbf{x,y,z}$.

[illegible]

	0
	1
	2

- d)* Draw the $ROBDT[C]$ of the propositional from $C(x, y, z)$ with the **variable ordering** $\mathbf{x, z, y}$.

[illegible]

A diagram showing a 2x2 grid of squares. To the right of the grid, the number 0 is aligned with the top row and the number 1 is aligned with the bottom row.

Problem 5 Propositional Logic (6 credits)

Given is the propositional form $D(a, b, c)$:

$$D(a, b, c) \iff [a \leftrightarrow (b \longrightarrow c)] \vee \neg(a \longrightarrow \neg b)$$

Give the canonical conjunctive normal form $CCNF[D]$ of D .

Name all laws you use.

0		
1		
2		
3		
4		
5		
6		

[illegible]

		0
		1
		2
		3
		4
		5
		6
		7
		8

0
1
2
3
4
5
6
7
8

3
4
5
6
7
8

73

73

73

This image shows a full page of blank graph paper. The grid consists of small, equal-sized squares formed by thin, light gray lines. There are no margins, text, or other markings on the page.

	0
	1
	2
	3
	4
	5

	0
	1
	2
	3
	4
	5



	0
	1
	2
	3
	4
	5

	0
	1
	2
	3
	4
	5



This image shows a full page of blank graph paper. The grid consists of small, uniform squares formed by thin, light gray lines. There are no margins, text, or other markings on the page.

Given are the following sets and predicates:


- S : Set of all students.
- L : Set of all lectures.
- Pxy : Student x is present in lecture y .
- Cy : Lecture y is being cancelled.

0  a)* $A_1 \iff$ For every lecture applies: If there is at least one student present in the lecture, it will not be
1  cancelled.

[illegible]

0  b) $A_2 \iff$ If statement A_1 is being observed and there is at least one student present in every lecture, then
1  all lectures will take place.

[illegible]

0  c) $A_3 \iff$ If there is a lecture that has been cancelled, then no student was present in that lecture.

[illegible]

	0
	1
	2
	3
	4
	5

	0
	1
	2
	3
	4
	5

	0
	1
	2
	3
	4
	5

	0
	1
	2
	3
	4
	5


- | | |
|--|---|
| | 0 |
| | 1 |
| | 2 |
| | 3 |
| | 4 |
| | 5 |

	0
	1
	2
	3
	4
	5

This image shows a full page of blank graph paper. The grid consists of small, equal-sized squares formed by thin gray lines. There are 20 columns and 20 rows of squares, creating a total of 400 square units. The grid covers the entire area of the page, leaving no margins or additional markings.

Problem 10 Predicate Logic and Sets (6 credits)

0
1
2
3
4
5
6



Prove the following implication by means of the deductive proof scheme:

$$(A \triangle B \subseteq B) \wedge (\overline{C} \cup A = G) \implies C \subseteq B$$



G is the universal set. Name all laws you use.

Row		From Row(s)	Law(s)
1	$A \triangle B \subseteq B$		
2	$\overline{C} \cup A = G \implies C \subseteq B$		
3			
4			
5			
6			
7			
8			
9			
10			
11			
12			
13			
14			
15			
16			
17			
18			



Problem 11 Relations (8 credits)

a)* Prove that the following subset relations are logically equivalent:
Name all laws you use.

$$W^{-1}\overline{X^{-1} \cap Y^{-1}} \subseteq \overline{Z} \iff Z^{-1}W^{-1} \subseteq X \cap Y$$



b)* Given is the Hasse diagram $G_H = (A, H)$ of the partial order relation $R \subseteq A^2$ with $A = \{1, 2, 3, 4, 5, 6, 7\}$:




Calculate the following bounds and extrema of the set $B = \{2, 3, 4, 5\}$:
 $ub(B)$, $max(B)$, $lst(B)$, $glb(B)$



Problem 12 Relations (5 credits)

0		
1		
2		
3		
4		
5		

Given is the following graph $G = (A, R)$ of the relation $R \subseteq A^2$ with $A = \{1, 2, 3, 4, 5\}$:



Calculate the transitive closure R^+ by means of the Warshall Algorithm. Use the variable order given in the table below. For each iteration cycle, give $\Gamma^-(i)$, $\Gamma^+(i)$, and $G = (A, R_i)$.

i	$\Gamma^-(i)$	$\Gamma^+(i)$	$G = (A, R_i)$
1			
2			
3			
4			
5			

Given is the relation R by the following successor table, with $R \subseteq A^2$:

x	$\Gamma^+\{x\}$
1	$\{2,3\}$
2	$\{3,4,5,6\}$
3	$\{4,6\}$
4	$\{1,5\}$
5	$\{1,3,6\}$
6	$\{1,4,6\}$

0		
1		
2		
3		

[illegible]

Mark correct answers with a cross

To re-mark an option, use a human-readable marking

- ☐ $R = R^+$
- ☐ The nodes 3 and 1 are indirectly accessible.
- ☐ R is antisymmetric.
- ☐ R is semiconnex.
- ☐ $|pred(\{3\})| = 4$
- ☐ R is a homogeneous relation.
- ☐ R is asymmetric.
- ☐ R is connex.

$$\begin{array}{|c|c|} \hline & \\ \hline & \\ \hline \end{array} \begin{array}{l} 0 \\ 1 \end{array}$$

	0
	1
	2

- | | |
|--|---|
| | 0 |
| | 1 |
| | 2 |

	0
	1

[illegible][illegible][illegible]

[illegible]

0
1

[illegible]

	0
	1
	2

[illegible]

- $|\Pi_i| = 2$
- $\exists_{K \in \Pi_i} |K| = 1$

[illegible]

Additional space for solutions—clearly mark the (sub)problem your answers are related to and strike out invalid solutions.

