PharmaGuide

SECOND EDITION · 2026

SUPPLEMENT SAFETY INTELLIGENCE BRIEF

Know what you're *actually* taking.

The clinician-reviewed field guide to labels, doses, hidden ingredients and interactions — built on primary research, written for real life.

60.2%	776	347%
of U.S. adults take a supplement	supplements FDA found hiding drugs	of labeled dose, in one tested product

CLINICIAN-REVIEWED BY

Laurie Pham, PharmD

12 SECTIONS · 21 CITED SOURCES

pharmaguide.io

INSIDE THIS BRIEF

Twelve sections. *One habit:* check before you take.

01	Label accuracy	What's on the label vs. what's in the bottle	04
02	Read a label	The five lines that matter	05
03	The proprietary blend trap	Why hidden amounts limit dose-risk checks	06
04	Forms that absorb	Same milligrams, different results	07
05	Third-party seals	What they prove — and what they don't	08
06	Upper limits & stacking	The dose that adds up across bottles	09
07	Interactions people miss	Supplement–medication pairs to know	10
08	Hidden drugs	What FDA keeps finding inside supplements	11
09	Your liver	The organ that processes everything	12
10	Timing & spacing	A sample day that avoids common conflicts	13
11	Kids & the household	Gummies, candy and safe storage	14
12	The checklist	Eight checks before you buy	15
✦	Questions for your pharmacist	Bring these to your next visit	16
✦	My stack worksheet	Fill in, bring to appointments	17
✦	Sources	Every claim, traceable	18

HOW TO READ SEVERITY IN THIS BRIEF

CONTRAINDICATED	AVOID	CAUTION	MONITOR	INFORMATIONAL
Do not combine	Strong reason to separate or skip	Risk depends on context	Manage with timing or awareness	No action needed


Ratings describe general risk from published references. Your doses, conditions and medications can move an item up or down a tier.

THE STATE OF THE CABINET

A label tells you what's in one bottle. Not what happens with your *other* bottles.

Many people take several supplements, sometimes alongside prescriptions. The risk usually isn't one ingredient — it's the combination, the dose that stacks, or the timing.

WHO TAKES SUPPLEMENTS



CDC/NCHS Data Brief No. 561, 2026 · U.S. adults, 2021–2023 [06]

WHEN IT GOES WRONG

~23,000

ER visits a year linked to supplement adverse events

~2,150

hospitalizations a year

Geller et al., NEJM 2015 (PMID 26465986) [05]

HIDDEN INGREDIENTS

776

supplements FDA found containing unapproved drugs, 2007–2016, from 146 companies

Tucker et al., JAMA Netw Open 2018 [03]

LIVER INJURY SHARE



of liver-injury cases at U.S. study centers tied to herbs & supplements, 2004–2013

Navarro et al., Hepatology 2014 [04]

THE TAKEAWAY

Many of these events may be avoidable by knowing exactly what you take, how much, and how it interacts. That is what this brief teaches.


01

LABEL ACCURACY

What's on the label isn't always *what's in the bottle.*

In a study published in 2023, researchers tested 25 melatonin gummy products and measured what was actually inside. It's the clearest recent picture of how far a supplement label can drift from reality.

25 PRODUCTS TESTED



- 3 within ±10% of label
- 21 off by more than 10%
- 1 had no melatonin, only CBD

347%

The highest amount found, relative to the label. The range across products ran from 74% to 347% of the stated dose — 1.3 to 13.1 mg per serving.



Why the dose matters. The same researchers note that as little as 0.1–0.3 mg can lift blood melatonin into the normal nighttime range in young adults. Many gummies delivered far more.

The limits. The study tested gummies only, one sample per product. It doesn't tell us about tablets or batch-to-batch variation — but it shows why "what the label says" is a claim, not a guarantee.

WHAT TO DO

Prefer products with an independent verification seal (Section 05). For sleep aids especially, start at the lowest available dose and ask your pharmacist.

Cohen PA, Avula B, Wang YH, Katragunta K, Khan I. Quantity of melatonin and CBD in melatonin gummies sold in the US. JAMA. 2023;329(16):1401-1402. doi:10.1001/jama.2023.2296

02

READ A LABEL

The Supplement Facts panel is your *starting point.*

Supplement Facts

Serving Size 2 Capsules
Servings Per Container 30

	Amount Per Serving	%DV
1 Magnesium (as magnesium oxide)	400 mg	95%
4 Vitamin B6 (as pyridoxine HCl)	50 mg	2,941%
3 Calm Blend	450 mg	†
L-theanine, ashwagandha root, valerian		

Other ingredients: rice flour, gelatin, magnesium stearate.

5 *"Clinically proven to cure stress."*

Illustrative label · not a real product

1 The form, not just the milligrams

"Magnesium (as magnesium oxide)" — the form sits in parentheses and changes how much is absorbed (Section 04).

2 Serving size vs. what you'll take

"2 capsules" means each capsule carries half the listed dose. Always divide.

3 Blends hide the doses

A "Calm Blend" lists ingredients but not amounts, so its dose risks can't be fully assessed against your meds (Section 03).

4 %DV far above 100%

2,941% isn't a bonus. For some nutrients, it pushes you toward the upper limit (Section 06).

5 Claims that sound medical

"Cures" or "treats" a disease isn't legal language for a supplement. Treat it as a red flag about the brand.

03

THE PROPRIETARY BLEND TRAP

If the doses are *hidden,* dose risks can't be fully assessed.

A proprietary blend lists several ingredients under one total weight, without saying how much of each. It's legal — and common in sleep, energy and pre-workout products.

Sleep & Calm Blend · 1,200 mg	HIDDEN DOSES
Magnesium	?
L-theanine	?
Valerian root	?
Melatonin	?
Herbal extracts	?

Fully disclosed formula	DISCLOSED
Magnesium glycinate	200 mg
L-theanine	200 mg
Valerian root extract	300 mg
Melatonin	0.5 mg

01

Dose can't be confirmed

You can't tell if an active is at a meaningful amount — or mostly filler.

02

Dose risks can't be fully assessed

Known ingredients still flag, but a dose you can't see can't be weighed against your meds.

03

Limits can't be tracked

Unknown amounts can't be added up against upper limits (Section 06).

WHAT TO DO

Prefer fully disclosed formulas, where every active ingredient lists its own amount. Transparency is the fastest signal that a brand expects scrutiny.

Fair note: a blend isn't automatically unsafe. Hidden doses limit dose-risk assessment; if you take prescription medications, discuss the blend with your pharmacist.

04

FORMS THAT ABSORB

Same milligrams. *Not* the same result.

The form decides how much is absorbed and how well it's tolerated. It hides in parentheses on the label.

NUTRIENT	OFTEN LESS ABSORBED / HARSHER	OFTEN BETTER ABSORBED / TOLERATED	WHY
Magnesium	Oxide	Citrate · Chloride · Lactate	NIH notes these forms absorb more completely than oxide in small studies. Glycinate is often chosen for being gentle on the stomach. Oxide may still suit some uses (e.g. constipation).
Calcium	Carbonate (no food)	Citrate	Carbonate needs stomach acid and food; citrate absorbs with or without — helpful if you take acid reducers.
Iron	Ferrous sulfate (GI upset)	Ferrous bisglycinate	Sulfate works but is hard on the gut for many. In trials — mostly in pregnant women — bisglycinate caused fewer stomach side effects than other iron salts.
Folate	—	Folic acid · L-methylfolate	Folic acid has established evidence for preventing neural-tube defects. L-methylfolate is an active form, but common MTHFR variants are not a reason to avoid folic acid.
Vitamin B12	—	Cyanocobalamin · Methylcobalamin	Both forms work for most people; methylcobalamin is an active form some clinicians and consumers prefer.
CoQ10	—	Ubiquinone · Ubiquinol	Small studies in older adults are mixed: one found ubiquinol raised CoQ10 more, another found no significant difference. The formulation matters too.
Zinc	Oxide	Citrate · Gluconate	In a controlled study, oxide was absorbed less than citrate or gluconate — and barely at all by some people.

Reality check: for many people a well-made standard form is fine. Form matters most when you have a specific reason — ask your pharmacist.

Sources: NIH ODS fact sheets [09] · Firoz 2001 [08] · CDC [14] · Wegmüller 2014 [15] · Fischer 2023 [16] · Zhang 2018 [17] · Pravst 2020 [18]

05

THIRD-PARTY SEALS

The FDA does *not* approve supplements before they're sold.

Unlike drugs, supplements rarely need pre-market proof of safety or label accuracy. An independent seal is the closest thing to outside verification.

- **USP Verified**
  Identity, potency, purity and dissolution; made under good manufacturing practices.
- **NSF Contents Certified**
  What's on the label is in the bottle; screened for contaminants.
- **NSF Certified for Sport**
  Everything above, plus screening for substances banned in sport.
- **Informed Sport / Informed Choice**
  Banned-substance tests; Sport checks every batch.
- **BSCG Certified Drug Free**
  Every finished lot tested for banned substances; label claims and contaminants tested yearly.
- **ConsumerLab**
  Independent product testing and reviews (subscription).

A SEAL DOES TELL YOU

✓ (USP, NSF) Contents match the label
✓ (USP, NSF) Screened for contaminants
✓ (Sport seals) Tested for banned substances

A SEAL DOES NOT TELL YOU

✕ That it works for you
✕ That the dose is right for you
✕ That it's safe with your medications

WHAT TO DO

Treat a seal as a floor, not a guarantee. Verify it on the certifier's own website — seals can be copied onto packaging.

Program descriptions from each certifier's website [13]. Seal names link to the certifier.

06

UPPER LIMITS & STACKING

The dose that *stacks* is the one people miss.

UL = Tolerable Upper Intake Level — the most of a nutrient you can take each day before the risk of side effects starts to rise. It's a ceiling, not an intake target.



EXAMPLE · VITAMIN D ACROSS THREE PRODUCTS ILLUSTRATIVE



Each product is within the limit alone. Together they exceed the adult upper limit by 50%.

ADULT UPPER LIMITS (19+)

NUTRIENT	UPPER LIMIT (UL) / DAY	WHAT GOES WRONG ABOVE IT
Vitamin D	4,000 IU (100 mcg)	Raises blood calcium; almost always from over-supplementing.
Calcium	2,000-2,500 mg	Food + supplements; limit set for kidney-stone risk. Can also cause constipation.
Magnesium (supplemental)	350 mg	Counts magnesium from supplements and medicines; magnesium naturally present in food and beverages is not counted. Above this, loose stools are common.
Zinc	40 mg	Chronic high intake lowers copper.
Iron	45 mg	Above this, stomach side effects become more likely. High-dose iron should be clinician-guided; very large overdoses can be toxic.
Vitamin A (preformed)	10,000 IU (3,000 mcg)	Liver toxicity; birth-defect risk in pregnancy.
Vitamin B6	100 mg	U.S. UL is 100 mg/day. In 2023, EFSA set 12 mg/day for adults, after finding neuropathy may occur at intakes as low as about 50 mg/day [10].
Niacin	35 mg (added)	Supplements + fortified food. Nicotinic acid can flush; high doses need medical supervision.

NIH Office of Dietary Supplements fact sheets [09] · ULs vary by age, sex and pregnancy; most include food and supplements. ULs are not targets — most people need far less.

07

INTERACTIONS PEOPLE MISS

The one nobody mentioned is often the one that *matters.*

Well-documented pairs. Not a complete list — a prompt to review yours with a pharmacist.

SUPPLEMENT	WITH	TIER	MECHANISM	WHAT TO DO
St. John's Wort	SSRIs / antidepressants	CONTRAINDICATED	Risk of serotonin syndrome; also speeds breakdown of many drugs, weakening them.	Don't combine.
Red yeast rice	Statins	AVOID	Contains a natural statin — doubling up raises muscle & liver risk.	Treat as a statin; don't add to one.
Vitamin K	Warfarin	CAUTION	Directly opposes the blood thinner; intake swings destabilize INR.	Keep intake consistent; tell your clinician.
Fish oil · Vitamin E	Blood thinners / antiplatelets	CAUTION	Possible added antiplatelet / bleeding effect, mainly at higher doses.	Review with your blood-thinner prescriber; flag before surgery.
Calcium · Magnesium · Iron	Levothyroxine	CAUTION	Bind the drug in the gut, reducing absorption — about 20–25% with calcium in one study.	Separate by at least 4 hours.
Calcium · Magnesium · Iron	Tetracycline & fluoroquinolone antibiotics	CAUTION	Minerals chelate the antibiotic, reducing its effect.	Ask your pharmacist for the exact before-and-after spacing for your antibiotic, and follow its label.
Potassium	ACE inhibitors / ARBs	AVOID	Can push potassium too high (hyperkalemia).	Don't add potassium without supervision.



Don't stop a prescription on your own. Bring the combination to your pharmacist — the fix is usually timing or monitoring.

Sources: Zamfirescu 2011 [07] · NCCIH [12] · FDA labeling · NIH resources · severity is general.

08

HIDDEN DRUGS

Some supplements aren't supplements. They're *unlabeled drugs.*

Researchers analyzed a decade of FDA warnings about supplements found to contain unapproved pharmaceutical ingredients.

776

adulterated supplements identified by FDA, 2007–2016 · 146 companies

WHAT THEY WERE SOLD FOR



Muscle building 11.9% · other 1.8%

MARKETED FOR	MOST COMMON HIDDEN DRUG	SHARE OF THAT CATEGORY
Sexual enhancement	Sildenafil (the active drug in Viagra)	47.0%
Weight loss	Sibutramine (withdrawn from the U.S. market)	84.9%
Muscle building	Synthetic steroids or steroid-like ingredients	89.1%

20.2%

contained more than one unapproved drug

28

products named in repeat FDA warnings, months apart — warnings don't always end it

RED FLAGS ON THE PACKAGE OR LISTING

- ✕ Promises fast or dramatic results
- ✕ Words like guaranteed, miracle, or scientific breakthrough
- ✕ "Natural alternative" to a prescription drug
- ✕ Sold mainly online or shipped from overseas

WHAT TO DO

Be most skeptical in these three categories. Before buying, search the product on FDA's tainted-supplements and recall listings, and turn on FDA recall alerts.

Tucker J et al. Unapproved pharmaceutical ingredients included in dietary supplements associated with US FDA warnings. JAMA Netw Open. 2018;1(6):e183337. doi:10.1001/jamanetworkopen.2018.3337 · Red flags: FDA health-fraud guidance [20].


09

YOUR LIVER

The organ that processes *everything* you take.


The U.S. Drug-Induced Liver Injury Network tracks liver damage from medications and supplements at specialist referral centers.

SHARE OF LIVER-INJURY CASES FROM HERBS & SUPPLEMENTS



839 patients
8 U.S. centers
2004–2013

DEATH OR LIVER TRANSPLANT



Non-bodybuilding products caused the more severe cases — mostly in middle-aged women.

COMMONLY IMPLICATED

Bodybuilding products with steroid-like ingredients · green tea extract [19] · multi-ingredient products

WARNING SIGNS – SEE A CLINICIAN

Yellowing skin or eyes · dark urine · persistent nausea · upper-right abdominal pain · unusual fatigue

LOWER YOUR RISK

Tell every clinician

List herbal products and powders, not just pills — they're the ones most often left off.

Fewer ingredients

Multi-ingredient "fat burner" and bodybuilding formulas are harder to trace when something goes wrong.

If warning signs appear

Stop the suspected supplement and contact a healthcare professional promptly. Bring the bottle or label.

Read this carefully: these are referral-center cases, not national rates — the authors caution it doesn't prove liver injury is rising population-wide. It does show supplements are a real, growing share of the cases specialists see.

Navarro VJ et al. Liver injury from herbals and dietary supplements in the U.S. Drug-Induced Liver Injury Network. Hepatology. 2014;60(4):1399-1408. doi:10.1002/hep.27317 · Warning signs: NIH LiverTox [19].

10

TIMING & SPACING

When you take it can matter as much as *what* you take.

A sample day for someone on thyroid medication who also takes common minerals. It's an example of the spacing logic — not a prescription.

WAKE

Thyroid medication (levothyroxine)

Empty stomach, water only. Wait 30–60 min before food.

BREAKFAST

Fat-soluble vitamins (A, D, E, K)

Take with a meal that has some fat. Keep iron and calcium away from thyroid meds (4+ hrs), including a multivitamin with minerals.

MIDDAY

Iron

Away from calcium and coffee/tea; vitamin C helps absorption.

DINNER

Calcium

Carbonate with food. Separate from iron; 4+ hrs after thyroid meds.

EVENING

Magnesium

One example slot; split large doses.


Metformin users: long-term use is linked to B12 depletion — worth periodic monitoring with your clinician.

High-dose zinc: can deplete copper over time — iron doses of 25 mg or more taken together can reduce zinc absorption [09].

11

KIDS & THE HOUSEHOLD

Gummies look like *candy*. Kids treat them that way.



+530%

rise in yearly pediatric melatonin ingestions reported to poison centers. Most involved young children, and most were unintentional.

HOUSEHOLD RULES THAT WORK

Store it like medicine

Up high, out of sight, child-resistant cap closed — every time.

Never call it candy

Kids copy adults. Take yours where they can't watch.

Ask before giving it to a child

Check with your pediatrician first, including sleep aids.

Count the gummies

Knowing how many were in the bottle helps if one goes missing.

Child swallowed too much? Call Poison Control right away. **1-800-222-1222**

Lelak K et al. Pediatric melatonin ingestions – United States, 2012-2021. MMWR Morb Mortal Wkly Rep. 2022;71:725-729. doi:10.15585/mmwr.mm7122a1 · Poison Help: HRSA [21].

12

THE CHECKLIST

Eight checks before you buy.

By hand, this takes a few minutes per bottle. Screenshot this page for the store — or let PharmaGuide help.

- [ ] **Every ingredient shows its amount** 01
  No "blend" that hides how much of each thing is inside.
- [ ] **It has a testing seal** 02
  USP or NSF on the label. Look it up on the seal's website to make sure it's real.
- [ ] **It uses a form your body absorbs well** 03
  Example: magnesium citrate instead of magnesium oxide. See page 7.
- [ ] **You're not doubling up** 04
  Add up the same vitamin across all your bottles. Stay under the daily limit on page 9.
- [ ] **You've checked for potential interactions with your medicines** 05
  See page 10, and ask your pharmacist.
- [ ] **It's not a risky type of product** 06
  Be extra careful with sex-boost, weight-loss and muscle products.
- [ ] **The brand has no FDA warnings or recalls** 07
  Search the brand name plus "FDA warning" or "recall."
- [ ] **The promises sound realistic** 08
  No "cures," "miracle" or "works overnight."

PharmaGuide helps you review checks 1–5.

Scan your supplement, then add your medications and supplement stack to review potential concerns.

pharmaguide.io →

At home, that part's on you

Keep bottles up high, away from kids. Throw out anything expired.

✦ TOOL · QUESTIONS FOR YOUR PHARMACIST

Bring these questions to *your pharmacist.*



Bring your bottles — or the worksheet on the next page — and ask these seven questions.

Q1	Is this supplement safe with everything else I take — including over-the-counter medicines?
Q2	Should I separate it from any of my medications? By how many hours?
Q3	Is this the right form and dose for me, given my age and conditions?
Q4	Does anything I take deplete a nutrient I should be monitoring?
Q5	Should I stop it before surgery or a procedure? How far ahead?
Q6	Is this product in a category with a history of hidden ingredients?
Q7	What side effects would mean I should stop and call you?

> PRO TIP
>
> Use one pharmacy for all prescriptions so their system can catch drug interactions — then tell them about your supplements, which their system usually can't see.

✦ TOOL · MY STACK WORKSHEET

Everything you take, *on one page.*

Include prescriptions, over-the-counter medicines, vitamins, herbs and powders. Print it, fill it in, bring it to every appointment.

PRODUCT / MEDICATION	DOSE	WHEN	WHY I TAKE IT	PRESCRIBED? Y/N

CONDITIONS & ALLERGIES

QUESTIONS TO ASK

SOURCES

Every claim, *traceable.*

01 Cohen PA, Avula B, Wang YH, Katragunta K, Khan I. Quantity of melatonin and CBD in melatonin gummies sold in the US. JAMA. 2023;329(16):1401–1402. doi:10.1001/jama.2023.2296

02 Lelak K, Vohra V, Neuman MI, Toce MS, Sethuraman U. Pediatric melatonin ingestions — United States, 2012–2021. MMWR. 2022;71:725–729. doi:10.15585/mmwr.mm7122a1

03 Tucker J, Fischer T, Upjohn L, Mazzera D, Kumar M. Unapproved pharmaceutical ingredients included in dietary supplements associated with US FDA warnings. JAMA Netw Open. 2018;1(6):e183337. doi:10.1001/jamanetworkopen.2018.3337

04 Navarro VJ, Barnhart H, Bonkovsky HL, et al. Liver injury from herbals and dietary supplements in the U.S. Drug-Induced Liver Injury Network. Hepatology. 2014;60(4):1399–1408. doi:10.1002/hep.27317

05 Geller AI, Shehab N, Weidle NJ, et al. Emergency department visits for adverse events related to dietary supplements. N Engl J Med. 2015;373:1531–1540. PMID 26465986

06 National Center for Health Statistics. Dietary supplement use among adults: United States, August 2021–August 2023. NCHS Data Brief No. 561. 2026. cdc.gov/nchs/data/databriefs/db561.pdf

07 Zamfirescu I, Carlson HE. Absorption of levothyroxine when coadministered with various calcium formulations. Thyroid. 2011;21(5):483–486. doi:10.1089/thy.2010.0296

08 Firoz M, Graber M. Bioavailability of US commercial magnesium preparations. Magnes Res. 2001;14(4):257–262. PMID 11794633

09 NIH Office of Dietary Supplements. Health professional fact sheets: vitamin D, calcium, magnesium, zinc, iron, folate, vitamin B12, vitamin A, vitamin B6, niacin. ods.od.nih.gov/factsheets/list-all

10 EFSA. Scientific opinion on the tolerable upper intake level for vitamin B6. EFSA J. 2023;21(5):8006. doi:10.2903/j.efsa.2023.8006

11 U.S. Food and Drug Administration. Tainted products marketed as consumer products; supplement recalls and safety alerts. fda.gov/drugs/medication-health-fraud

12 NIH National Center for Complementary and Integrative Health. St. John's wort and other herb–drug interaction fact sheets. nccih.nih.gov/health/st-johns-wort

13 Certifier program descriptions: USP, NSF International, LGC Informed Sport, BSCG, ConsumerLab. quality-supplements.org · nsf.org · wetestyoutrust.com · bscg.org · consumerlab.com

14 Centers for Disease Control and Prevention. MTHFR gene variant and folic acid facts. cdc.gov/folic-acid/data-research/mthfr

15 Wegmüller R, Tay F, Zeder C, Brnic M, Hurrell RF. Zinc absorption by young adults from supplemental zinc citrate is comparable with that from zinc gluconate and higher than from zinc oxide. J Nutr. 2014;144(2):132–136. PMID 24259556

16 Fischer JAJ, Cherian AM, Bone JN, Karakochuk CD. The effects of oral ferrous bisglycinate supplementation on hemoglobin and ferritin concentrations in adults and children: a systematic review and meta-analysis of randomized controlled trials. Nutr Rev. 2023;81(8):904–920. PMID 36728680

17 Zhang Y, Liu J, Chen XQ, Chen CYO. Ubiquinol is superior to ubiquinone to enhance coenzyme Q10 status in older men. Food Funct. 2018;9(11):5653–5659. doi:10.1039/c8fo00971f

18 Pravst I, Rodríguez Aguilera JC, Cortes Rodriguez AB, et al. Comparative bioavailability of different coenzyme Q10 formulations in healthy elderly individuals. Nutrients. 2020;12(3):784. ncbi.nlm.nih.gov/pmc/articles/PMC7146408

19 National Institutes of Health. LiverTox: clinical and research information on drug-induced liver injury. ncbi.nlm.nih.gov/books/NBK547852

20 U.S. Food and Drug Administration. Health fraud scams: consumer guidance on spotting fraudulent health products. fda.gov/consumers/health-fraud-scams

21 Health Resources and Services Administration. Poison Help line, 1-800-222-1222. poisonhelp.hrsa.gov

This brief is educational and was clinician-reviewed for accuracy and clarity by Laurie Pham, PharmD. It is not a substitute for individual medical advice.

PharmaGuide

THIS BRIEF IS THE MANUAL VERSION

Check your *whole stack* in seconds.

Sourced from NIH's 200,000+ supplement labels, PharmaGuide checks your stack against your medications and conditions — with the reasoning and sources behind every flag. Core checks run privately on your device.

- ✓ Drug–supplement interactions
- ✓ Supplement–supplement interactions
- ✓ Recalled & banned ingredients
- ✓ Duplicate ingredients
- ✓ Upper-limit tracking
- ✓ Condition-specific risks
- ✓ Medication nutrient depletion
- ✓ Third-party testing status

Get early access · pharmaguide.io

Found this useful? Forward it to someone who takes a few.

FDA LABELS · NIH RESOURCES · PUBMED-INDEXED RESEARCH | CLINICIAN-REVIEWED | PRIVATE BY DESIGN

EDUCATIONAL INFORMATION ONLY — NOT MEDICAL ADVICE. These statements have not been evaluated by the Food and Drug Administration. This brief is not a substitute for professional care; consult your physician or pharmacist before changing any medication or supplement. Severity ratings reflect general references and may differ for your doses and health profile. Clinician-reviewed by Laurie Pham, PharmD. © 2026 PharmaGuide Inc. · Boston, MA · pharmaguide.io