Effects of Early Forming Massive Stars

Sean C. Lewis

December 30, 2022

Early Forming Massive Stars

Disruption of gas collapse, star formation, and cluster assembly

- The feedback from massive stars likely dominates the self-regulation of star formation.

Early Forming Massive Stars

Disruption of gas collapse, star formation, and cluster assembly

- The feedback from massive stars likely dominates the self-regulation of star formation.
- Gas evacuation (via stellar feedback) is crucial to the completion of star cluster assembly.[1]

[1] Bressert et al. 2010; Longmore et al. 2014; Grudić et al. 2018; Dobbs et al. 2022

Early Forming Massive Stars

Disruption of gas collapse, star formation, and cluster assembly

- The feedback from massive stars likely dominates the self-regulation of star formation.
- Gas evacuation (via stellar feedback) is crucial to the completion of star cluster assembly.[1]
- *How* gas is removed (rapidly, or slowly) may affect cluster structure.[2]

[1] Bressert et al. 2010; Longmore et al. 2014; Grudić et al. 2018; Dobbs et al. 2022

[2] Smith et al. 2013; Portegies Zwart et al. 2010; Banjeree & Kroupa 2017

Early Forming Massive Stars

Disruption of gas collapse, star formation, and cluster assembly

- The feedback from massive stars likely dominates the self-regulation of star formation.
- Gas evacuation (via stellar feedback) is crucial to the completion of star cluster assembly.[1]
- *How* gas is removed (rapidly, or slowly) may affect cluster structure.[2]

- What about *when* massive stars form? Using our computational model we test the effects of early forming massive stars.

[1] Bressert et al. 2010; Longmore et al. 2014; Grudić et al. 2018; Dobbs et al. 2022
[2] Smith et al. 2013; Portegies Zwart et al. 2010; Banjeree & Kroupa 2017

Torch

Stars from gas

- Couples N-body, stellar evolution, and feedback in `AMUSE` with self-gravitating magnetized gas in MHD code `FLASH`.
- Resolved dynamics of stars and gas; study star cluster formation within collapsing GMCs.
- Form stars from sink particles which each have a randomized star mass list sampled from the Kroupa IMF.[3]

[3] Kroupa 2001

A Controlled Experiment





Lewis et al. 2023

Effects on Gas Energy





Effects on Gas Accretion and Star Formation





Effects on Gas Accretion and Star Formation



Early forming massive stars reduces sink accretion and star formation rates.

Effects on Gas Accretion and Star Formation



Early forming massive stars reduces sink accretion and star formation rates.

Run	$\langle \epsilon_{ff} \rangle$
Fid	0.23
50M	0.08
70M	0.03
100M	0.04

Effects on Star Clustering, Cluster Assembly



- DBSCAN to identify cluster with at least 50% bound members and 100 $M_\odot$ at $2\tau_{\rm ff}$.
- Clusters in runs with early massive stars are less massive and more fragmented compared to the fiducial run.

Run	Mass in Clusters $10^3\ M_\odot$	Frac Mass M_c/M_{tot}	r_h MMC pc	E_{bind} MMC 10^{46} erg
Fid	3.6	0.99	0.25	-140
50M	1.4	0.97	0.17	-12
70M	0.86	0.85	0.21	-4.2
100M	0.62	0.46	0.18	-3.8

Effects of Early Forming Massive Stars

- Significantly disrupt the natal gas structure, resulting in premature unbinding of GMC.
- The star formation rate per free-fall time is suppressed by up to a factor of seven, reducing the total mass of stars formed.
- Stifle the hierarchical assembly process of massive star clusters, instead promoting the formation of spatially separate and more loosely bound subclusters.

The Problem with Initial Conditions

- Self consistent galactic scale simulations with resolution down to sub-tenth parsec scales and include Nbody individual stellar dynamics and individual stellar feedback all at once? A little tough.

- Creating our own isolated clouds from scratch? "Creative liberties..."

Clouds from Galactic Simulations



GMC identification[4]

[4] Li, H. et al. 2020

From AREPO to FLASH

Try CIC Mapping?



Cloud from raw AREPO data represented using SPH kernels



Cloud-in-cell mapping onto AMR FLASH grid


300
200
100
0
−100
−200
−300
y (pc)
−300
−200
−100
0
100
200
300
x (pc)


300
200
100
0
−100
−200
−300
y (pc)
−300
−200
−100
0
100
200
300
x (pc)


300
200
100
0
−100
−200
−300
y (pc)
−300
−200
−100
0
100
200
300
x (pc)


300
200
100
0
−100
−200
−300
y (pc)
−300
−200
−100
0
100
200
300
x (pc)

Voronoi Mesh to AMR Grid





VorAMR: Logic path



VorAMR: The Big Wins

- Significantly expands Torch's horizon and moves Torch to "completion".
- Opens wide avenue of collaboration; code bases do not have to be exclusive!
- More accurate visualizations (no more estimating Voronoi meshes as SPH kernels in `yt`).

Thank You!

-

Questions?

Appendix

$$\epsilon_{\mathrm{ff}} = \dot{M}_* \frac{t_{\mathrm{ff}}}{M_{\mathrm{g}}} \tag{1}$$