

Summer Research: Hypervelocity Globular Cluster



July 11, 2018


M87



Identification of HVGC-1

- Caldwell et al. 2014
- Extremely blue-shifted object in M87

Figure 1. from A Globular Cluster toward M87 with a Radial Velocity < -1000 km s $^{-1}$: The First Hypervelocity Cluster
Caldwell et al. 2014 ApJL 787 L11 doi:10.1088/2041-8205/787/1/L11
<http://dx.doi.org/10.1088/2041-8205/787/1/L11>
© 2014. The American Astronomical Society. All rights reserved.




Possible mechanisms for acceleration

- HVGC-1 received kick from potential well of merging dark matter halos.
 - ▶ Samsing 2015
- Kick from interaction with a supermassive binary black hole.
 - ▶ Survival of an extended Globular Cluster has not been simulated/investigated.


Blackhole circular orbits.

- Can set any BH mass ratio and separation.
- Caldwell calls for HVGC-1 to pass 1pc from 10:1 ratio BBH.




10:1 Mass ratio: HVGC-1 Pass

- Perturbation: $\gamma = \frac{M_{BH}}{M_{GC}} \left(\frac{R}{D}\right)^3$
- Minimize γ while still producing a hyper-velocity object.
- Prograde-planar interaction.



Next Steps - 10:1 Mass ratio: HVGC-1 Pass

- Input params: GC starting position (closest approach), BH separation, BH mass ratio
- Output: Velocity of GC, max perturbation experienced.
- Tidal radius of 0.1pc
- Cluster $r_h = 6\text{pc}$



Questions and Preparing for Meetings

- Make Slack channel for astrophysics group?
 - ▶ Useful for quick questions, document sharing.
- Continue making slides for weekly meetings.
 - ▶ Helpful for making progress and developing presenting skills.
 - ▶ Allows entire group to be involved, up-to-date, and able to provide informed advice.

Backup Slides

3:1 Mass ratio

- 2-3 pc pass from larger BH.
- Tidal radius of 0.3-0.4 pc

