
kmerseek 982a055, 70 settings grouped by alphabet size (2-3, 4-8, 12-18, 20 letters); below the dashed line is the pass side
Ced9 vs human proteome (notebook 241)
Ced9 vs BCL2 alone, predicted
Ced9 vs BCL-2 family (14), predicted
no region on BCL2
region on BCL2, no E-value (λ = 0)
--- E = 0.05 (decision line)
Ced9 → BCL2: best E-value
100_000
10_000
1_000
100
10
1
0.1
0.01
P66 vs human proteome (notebook 241)
P66 vs CD47 alone, predicted
no region on CD47
region on CD47, no E-value (λ = 0)
--- E = 0.05 (decision line)
P66 → CD47: best E-value
hp_kyte_doolittle2 k17
hp_kyte_doolittle2 k21
hp_kyte_doolittle2 k26
hp_kyte_doolittle2 k30
hp_kyte_doolittle2 k34
hp_kyte_doolittle2 k38
hp_lehninger2 k16
hp_lehninger2 k17
hp_lehninger2 k19
hp_lehninger2 k20
hp_lehninger2 k24
hp_lehninger2 k28
hp_lehninger2 k32
hp_lehninger_c_nonpolar2 k16
hp_lehninger_c_nonpolar2 k20
hp_lehninger_c_nonpolar2 k24
hp_lehninger_c_nonpolar2 k28
hp_lehninger_c_nonpolar2 k32
hp_pbotc_1st_ed2 k16
hp_pbotc_1st_ed2 k20
hp_pbotc_1st_ed2 k24
hp_pbotc_1st_ed2 k28
hp_thomas_dill2 k17
hp_thomas_dill2 k21
hp_thomas_dill2 k29
hp_thomas_dill_no_c2 k17
hp_thomas_dill_no_c2 k21
hp_thomas_dill_no_c2 k25
hp_thomas_dill_no_c2 k29
hp_thomas_dill_no_c2 k34
hp_lehninger_hpc3 k14
hp_lehninger_hpc3 k18
hp_lehninger_hpc3 k25
gbmr4 k10
gbmr4 k13
gbmr4 k16
gbmr4 k18
gbmr4 k21
polarity4 k9
polarity4 k11
polarity4 k13
polarity4 k16
wwmj5 k7
wwmj5 k9
wwmj5 k11
wwmj5 k13
dayhoff6 k7
dayhoff6 k9
dayhoff6 k11
dayhoff6 k12
gbmr7 k8
funcgroups8 k12
mmseqs12 k5
mmseqs12 k6
mmseqs12 k7
mmseqs12 k8
sdm12 k5
sdm12 k6
sdm12 k8
sdm12 k9
wass14 k6
wass14 k7
wass14 k8
hsdm17 k5
hsdm17 k6
uniprot18 k5
uniprot18 k6
protein20 k5
protein20 k6
protein20 k7