

13. human TNFAIP3 ZN\_FING "A20-type 3" (34 aa) and mouse Q60769

TOOLS: kmerseek hp\_thomas\_dill\_no\_c2 k19 (chosen for ZN\_FING in Stage 0) and dayhoff6 k8; phmmer, MMseqs2, MMseqs2 iterative, Foldseek, ProstT5, Reseek; Kyte-Doolittle scan

- kmerseek: call lands on the feature (>= 80% of call inside, >= 30% covered)

call mostly inside the feature, lower IoU than kmerseek

call overlaps the feature but < 50% of the call is inside it



call reaches kmerseek's IoU or higher

no call of this feature type overlaps the feature
- the tool's best call is on another target protein

Kyte-Doolittle scan: no target protein, it searches nothing

the Swiss-Prot feature being scored (human) / transferred from (target)

other Swiss-Prot features, labelled by type



Hypothesis: kmerseek places the human ZN\_FING feature "A20-type 3" on its mouse counterpart and the structure tools do not.  
Conclusion: kmerseek IoU 0.83, 29 of 33 residues identical; Foldseek spills; ProstT5 spills; Reseek spills; phmmer spills; MMseqs2 spills; MMseqs2 iterative spills; Kyte-Doolittle scan did not land; pLDDT 69, disorder fraction 0.24.