
BHF in the dark-set run, every kmerseek setting; target Swiss-Prot minus Ascidiacea. Rule 3: shared k-mers outside low complexity on BHF and on the target
kmerseek setting, scaled 1: best region
kmerseek setting, scaled 10: best region
phmmer and jackhmmer best hit (E = 0.16)
E = 1
best region E < 1, before the rules
best region kept by rule 3
a
Best region E-value on BHF (lower is better)
hp_kyte_doolittle2
hp_lehninger2
hp_lehninger_c_nonpolar2
hp_pbotc_1st_ed2
hp_thomas_dill2
hp_thomas_dill_no_c2
hp_lehninger_hpc3
gbmr4
polarity4
wwmj5
dayhoff6
funcgroups8
mmseqs12
sdm12
wass14
hsdm17
uniprot18
protein20
Alphabet, by number of letters (2-3, 4-8, 12-18, 20)
b
gbmr4.k16.s1
gbmr4.k18.s1
gbmr4.k20.s1
hp_lehninger_hpc3.k17.s1
gbmr4.k22.s1
hp_thomas_dill2.k20.s1
hp_thomas_dill_no_c2.k26.s10
hp_lehninger_c_nonpolar2.k21.s10
hp_lehninger_hpc3.k25.s10
hp_thomas_dill_no_c2.k24.s10
none kept
Region E-value