
calibrated: 0.5 to 2
E* = 0.1
E* = 1
E* = 10
composition shuffles
dipeptide shuffles
kmerseek 0a22970, mmseqs12 k=4, extension penalty 2, give-up margin 8; queries: shuffles of held-out windows; target: whole collection
15 aa, not low-complexity, low hydrophobic (n=28,980)
15 aa, not low-complexity, middle hydrophobic (n=21,040)
15 aa, not low-complexity, high hydrophobic (n=54,820)
15 aa, SEG low-complexity, low hydrophobic (n=5,138)
15 aa, SEG low-complexity, middle hydrophobic (n=1,180)
15 aa, SEG low-complexity, high hydrophobic (n=1,960)
30 aa, not low-complexity, low hydrophobic (n=22,000)
30 aa, not low-complexity, middle hydrophobic (n=24,200)
30 aa, not low-complexity, high hydrophobic (n=45,120)
30 aa, SEG low-complexity, low hydrophobic (n=11,380)
30 aa, SEG low-complexity, middle hydrophobic (n=5,360)
30 aa, SEG low-complexity, high hydrophobic (n=5,060)
53 aa, not low-complexity, low hydrophobic (n=21,140)
53 aa, not low-complexity, middle hydrophobic (n=20,600)
53 aa, not low-complexity, high hydrophobic (n=35,160)
53 aa, SEG low-complexity, low hydrophobic (n=19,700)
53 aa, SEG low-complexity, middle hydrophobic (n=9,400)
53 aa, SEG low-complexity, high hydrophobic (n=7,120)
0.05
0.1
0.2
0.5
1
2
5
Chance regions at E ≤ E*: observed / predicted (shuffles x E*)