

exact: k class matches in a row (dot = last position of a seed)
spaced: matches at the 1s of a pattern
chained: n exact k-mers within W positions
window: at least m matches among L positions
BH1 block, CED-9 154-190 / BCL2 130-166
(n chance seeds) = seed positions of that scheme on the other 19,731 human proteins, CED-9 query
a
CED-9 against BCL2 on the BH1 diagonal; hp_thomas_dill2 (H = ACFILMVWY, P = the rest)
CED-9 residues
BCL2 residues
CED-9 H/P class
same class (33 of 61)
BCL2 H/P class
exact k=12 (1,192,037 chance seeds)
exact k=18 (23,357 chance seeds)
exact k=20 (6,377 chance seeds)
spaced 1101101101101101101101 (94,660 chance seeds)
spaced 111010010100110111 (2,457,623 chance seeds)
chained 2 x k=8 in 40 (1,071,133 chance seeds)
chained 3 x k=8 in 60 (73,344 chance seeds)
window 18 of 20 (1,129,984 chance seeds)
window 27 of 30 (39,671 chance seeds)
window 36 of 40 (1,363 chance seeds)
140 145 150 155 160 165 170 175 180 185 190 195 200
CED-9 position (BCL2 position = CED-9 - 24)
b
CED-9 against BCL2 on the BH1 diagonal; hp_lehninger2 (H = AFGILMPVWY, P = the rest)
CED-9 residues
BCL2 residues
CED-9 H/P class
same class (34 of 61)
BCL2 H/P class
exact k=12 (847,057 chance seeds)
exact k=18 (15,332 chance seeds)
exact k=20 (3,853 chance seeds)
spaced 1101101101101101101101 (52,227 chance seeds)
spaced 111010010100110111 (1,504,807 chance seeds)
chained 2 x k=8 in 40 (601,116 chance seeds)
chained 3 x k=8 in 60 (31,150 chance seeds)
window 18 of 20 (640,516 chance seeds)
window 27 of 30 (13,576 chance seeds)
window 36 of 40 (306 chance seeds)
140 145 150 155 160 165 170 175 180 185 190 195 200
CED-9 position (BCL2 position = CED-9 - 24)