

6. human COL12A1 DOMAIN "Collagen-like 2" (51 aa) and chicken P13944

TOOLS: kmerseek hp_thomas_dill_no_c2 k19 (chosen for DOMAIN in Stage 0) and dayhoff6 k11; phmmer, MMseqs2, MMseqs2 iterative, Foldseek, ProstT5, Reseek; Kyte-Doolittle scan

- kmerseek: call lands on the feature ($\geq 80\%$ of call inside, $\geq 30\%$ covered)

call mostly inside the feature, lower IoU than kmerseek

call overlaps the feature but $< 50\%$ of the call is inside it



call reaches kmerseek's IoU or higher

no call of this feature type overlaps the feature
- the tool's best call is on another target protein

Kyte-Doolittle scan: no target protein, it searches nothing

the Swiss-Prot feature being scored (human) / transferred from (target)

other Swiss-Prot features, labelled by type



Hypothesis: kmerseek places the human DOMAIN feature "Collagen-like 2" on its chicken counterpart and the structure tools do not.
Conclusion: kmerseek IoU 0.60, 27 of 30 residues identical; Foldseek no call; ProstT5 spills; Reseek no call; phmmer equal or higher IoU; MMseqs2 spills; MMseqs2 iterative spills; Kyte-Doolittle scan did not land; pLDDT None, disorder fraction 1.0.