

25. human EHMT2 REPEAT "ANK 7" (30 aa) and arabidopsis Q38898

TOOLS: kmerseek hp_kyte_doolittle2 k19 (chosen for REPEAT in Stage 0) and hp_pbotc_1st_ed2 k18; phmmer, MMseqs2, MMseqs2 iterative, Foldseek, ProstT5, Reseek; Kyte-Doolittle scan

- kmerseek: call lands on the feature (>= 80% of call inside, >= 30% covered)

call mostly inside the feature, lower IoU than kmerseek

call overlaps the feature but < 50% of the call is inside it



call reaches kmerseek's IoU or higher

no call of this feature type overlaps the feature
- the tool's best call is on another target protein

Kyte-Doolittle scan: no target protein, it searches nothing

the Swiss-Prot feature being scored (human) / transferred from (target)

other Swiss-Prot features, labelled by type



Hypothesis: kmerseek places the human REPEAT feature "ANK 7" on its arabidopsis counterpart and the structure tools do not.
Conclusion: kmerseek IoU 0.81, 15 of 29 residues identical; Foldseek spills; ProstT5 spills; Reseek spills; phmmer equal or higher IoU; MMseqs2 spills; MMseqs2 iterative spills; Kyte-Doolittle scan did not land; pLDDT 95, disorder fraction 0.0.