

145. human ZKSCAN3 ZN\_FING "C2H2-type 6" (23 aa) and mouse P10077

TOOLS: kmerseek hp\_thomas\_dill\_no\_c2 k19 (chosen for ZN\_FING in Stage 0) and gbmr4 k12; phmmer, MMseqs2, MMseqs2 iterative, Foldseek, ProstT5, Reseek; Kyte-Doolittle scan

- kmerseek: call lands on the feature ( $\geq 80\%$  of call inside,  $\geq 30\%$  covered)

call mostly inside the feature, lower IoU than kmerseek

call overlaps the feature but  $< 50\%$  of the call is inside it



call reaches kmerseek's IoU or higher

no call of this feature type overlaps the feature
- the tool's best call is on another target protein

Kyte-Doolittle scan: no target protein, it searches nothing

the Swiss-Prot feature being scored (human) / transferred from (target)

other Swiss-Prot features, labelled by type



Hypothesis: kmerseek places the human ZN\_FING feature "C2H2-type 6" on its mouse counterpart and the structure tools do not. Conclusion: kmerseek IoU 0.81, 15 of 25 residues identical; Foldseek spills; ProstT5 spills; Reseek inside, lower IoU; phmmer inside, lower IoU; MMseqs2 spills; MMseqs2 iterative equal or higher IoU; Kyte-Doolittle scan did not land; pLDDT 71, disorder fraction 0.0.