

Read this: 20% identity or less, grouped on one half, measured on the other, 95% interval
all identities, whole pairs, 95% interval
Biased, not a result: 20% identity or less, grouped and measured on the same positions
two classes
three or more classes (number in brackets)
1 = same evidence as the 20 amino acids
Homologs:
right of the line is better
(more evidence that related proteins are related)
Not homologs (SCOPe):
right of the line is worse
(evidence that unrelated proteins are related)
Pfam seed
SCOPe:
same family
SCOPe: same superfamily,
different family
SCOPe: same fold,
different superfamily
SCOPe: different fold,
same class
protein20 (20)
uniprot18 (18)
hsdm17 (17)
wass14 (14)
mmseqs12 (12)
sdm12 (12)
funcgroups8 (8)
gbmr7 (7)
dayhoff6 (6)
wwmj5 (5)
gbmr4 (4)
polarity4 (4)
hp_lehninger_hpc3 (3)
hp_kyte_doolittle2 (2)
hp_lehninger2 (2)
hp_lehninger_c_nonpolar2 (2)
hp_pbotc_1st_ed2 (2)
hp_thomas_dill2 (2)
hp_thomas_dill_no_c2 (2)
0.5
1
2
I(alphabet) / I(20 aa)
higher is better
I(alphabet) / I(20 aa)
higher is worse