36.5% of dark Pacific oyster proteins have an AlphaFold DB model with mean pLDDT 50 or less, against 3.9% of lit ones (proteins with no model count in both totals).

Proteins: UniProt 2026_03 one-protein-per-gene sets; only the dark ones are drawn. Dark: no InterPro match and no Swiss-Prot hit above 30% identity (MMseqs2 18.8cc5c).
Colour: mean pLDDT of the AlphaFold DB model AF-<accession>-F1 (v6), in AlphaFold DB's colours; bands as in the four-protein figures: above 90, above 70 up to 90, above 50 up to 70, 50 or less.

One tile = one dark protein; 150 tiles per row, the same tile size as the two proteome maps on this page; longest protein first, read row by row.

pLDDT > 90 pLDDT 70-90 pLDDT 50-70 pLDDT 50 or less no AlphaFold DB model ◀ row holding every 1,000th tile

Pacific oyster (Crassostrea gigas): 5,877 dark proteins, 6,017 to 39 aa. pLDDT > 90 41, pLDDT 70-90 1,087, pLDDT 50-70 2,553, pLDDT 50 or less 2,148, no model 48 (the 47 longest, 1,282 aa and up, all have no model)



Human (Homo sapiens): 181 dark proteins, 1,178 to 2 aa. pLDDT > 90 3, pLDDT 70-90 48, pLDDT 50-70 22, pLDDT 50 or less 26, no model 82