## FERMAT'S LAST THEOREM FOR REGULAR PRIMES

### Contents

1.	Introduction	1
2.	Discriminants of number fields	1
3.	Cyclotomic fields	4
4.	Fermats Last Theorem for regular primes	6
References		7

# 1. Introduction

We prove Fermat's Last Theorem for regular primes and give some of the necessary background. It uses [Sam70, Mar18, Was82].

# 2. Discriminants of number fields

**lemma 2.1.** Let K be a number field,  $\alpha \in K$  and let  $\sigma_i$  be the embeddings of K into  $\mathbb{C}$ . Then

$$\operatorname{Tr}_{K/\mathbb{Q}}(\alpha) = \sum_{i} \sigma_{i}(\alpha) \qquad N_{K/\mathbb{Q}}(\alpha) = \prod_{i} \sigma_{i}(\alpha)$$

**lemma 2.2.** Let K be a number field with basis  $B = \{b_1, \ldots, b_n\}$  and let  $\sigma_1, \ldots, \sigma_n$  be the embeddings of K into  $\mathbb{C}$ . Now let M be the matrix

$$\begin{pmatrix} \sigma_1(b_1) & \cdots & \sigma_1(b_n) \\ \vdots & & \vdots \\ \sigma_n(b_1) & \cdots & \sigma_n(b_n) \end{pmatrix}.$$

Then

$$\Delta(B) = \det(M)^2.$$

*Proof.* By Proposition 2.1 we know that  $\operatorname{Tr}_{K/\mathbb{Q}}(b_ib_j) = \sum_k \sigma_k(b_i)\sigma_k(b_j)$  which is the same as the (i,j) entry of  $M^tM$ . Therefore

$$\det(T_B) = \det(M^t M) = \det(M)^2.$$

**lemma 2.3.** Let K be a number field and  $B = \{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$  for some  $\alpha \in K$ . Then

$$\Delta(B) = \prod_{i < j} (\sigma_i(\alpha) - \sigma_j(\alpha))^2$$

where  $\sigma_i$  are the embeddings of K into  $\mathbb{C}$ . Here  $\Delta(B)$  denotes the discriminant.

*Proof.* First we recall a classical linear algebra result relating to the Vandermonde matrix, which states that

$$\det \begin{pmatrix} 1 & x_1 & x_1^2 & \cdots & x_1^{n-1} \\ \vdots & & \vdots & \\ 1 & x_n & x_n^2 & \cdots & x_n^{n-1} \end{pmatrix} = \prod_{i < j} (x_i - x_j).$$

Combining this with Proposition 2.2 gives the result.

**lemma 2.4.** Let f be a monic irreducible polynomial over a number field K and let  $\alpha$  be one of its roots in  $\mathbb{C}$ . Then

$$f'(\alpha) = \prod_{\beta \neq \alpha} (\alpha - \beta),$$

where the product is over the roots of f different from  $\alpha$ .

*Proof.* We first write  $f(x) = (x - \alpha)g(x)$  which we can do (over  $\mathbb{C}$ ) as  $\alpha$  is a root of f, where now  $g(x) = \prod_{\beta \neq \alpha} (x - \beta)$ . Differentiating we get

$$f'(x) = g(x) + (x - \alpha)g'(x).$$

If we now evaluate at  $\alpha$  we get the result.

**lemma 2.5.** Let  $K = \mathbb{Q}(\alpha)$  be a number field with  $n = [K : \mathbb{Q}]$  and let  $B = \{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$ . Then

$$\Delta(B) = (-1)^{\frac{n(n-1)}{2}} N_{K/\mathbb{Q}}(m'_{\alpha}(\alpha))$$

where  $m'_{\alpha}$  is the derivative of  $m_{\alpha}(x)$  (which we recall denotes the minimal polynomial of  $\alpha$ ).

*Proof.* By Proposition 2.3 we have  $\Delta(B) = \prod_{i < j} (\alpha_i - \alpha_j)^2$  where  $\alpha_k := \sigma_k(\alpha)$ . Next, we note that the number of terms in this product is  $1 + 2 + \cdots + (n-1) = \frac{n(n-1)}{2}$ . So if we write each term as  $(\alpha_i - \alpha_j)^2 = -(\alpha_i - \alpha_j)(\alpha_j - \alpha_i)$  we get

$$\Delta(B) = (-1)^{\frac{n(n-1)}{2}} \prod_{i=1}^{n} \prod_{i \neq j} (\alpha_i - \alpha_j).$$

Now, by lemmama 2.4 and Proposition 2.1 we see that

$$N_{K/\mathbb{Q}}(m'_{\alpha}(\alpha)) = \prod_{i=1}^{n} m'_{\alpha}(\alpha_i) = \prod_{i=1}^{n} \prod_{i \neq j} (\alpha_i - \alpha_j),$$

which gives the result.

**lemma 2.6.** Let K be a number field and B, B' bases for  $K/\mathbb{Q}$ . If P denotes the change of basis matrix, then

$$\Delta(B) = \det(P)^2 \Delta(B').$$

**lemma 2.7.** If K is a number field and  $\alpha \in \mathcal{O}_K$  then  $\operatorname{Tr}_{K/\mathbb{Q}}(\alpha)$  and  $N_{K/\mathbb{Q}}(\alpha)$  are both in  $\mathbb{Z}$ .

**lemma 2.8.** Let K be a number field and  $B = \{b_1, \ldots, b_n\}$  be elements in  $\mathcal{O}_K$ , then  $\Delta(B) \in \mathbb{Z}$ .

**lemma 2.9.** Let K be a number field and  $B = \{b_1, \ldots, b_n\}$  be a basis for  $K/\mathbb{Q}$  consisting of algebraic integers. If B is not an integral basis then there

exists an algebraic integer of the form

$$\alpha = \frac{x_1b_1 + \dots + x_nb_n}{p}$$

where p is a prime and  $x_i \in \{0, ..., p-1\}$  with not all  $x_i$  zero. Moreover, if  $x_i \neq 0$  and we let B' be the basis obtained by replacing  $b_i$  with  $\alpha$ , then

$$\Delta(B') = \frac{x_i^2}{p^2} \Delta(B).$$

In particular  $p^2 \mid \Delta(B)$ .

*Proof.* If B is not an integral basis then we can find some element  $\phi \in \mathcal{O}_K$ such that

$$\phi = y_1 b_1 + \dots y_n b_n$$

with not all the  $y_i$  in  $\mathbb{Z}$ . So, let N be the least common multiple of the denominators of the  $y_i$  (meaning  $Ny_i \in \mathbb{Z}$  for all i). Now, let p be a prime factor of N. If we now consider  $(N/p)\phi$  then all of the coefficients of  $b_i$ are in  $\frac{1}{p}\mathbb{Z}$  (so they have denominator 1 or p.) and at least one of them has denominator p (since not all the  $y_i$  where in  $\mathbb{Z}$ ). So by relabelling, wlog we can assume

$$\phi = y_1 b_1 + \dots y_n b_n$$

with  $y_i \in \frac{1}{p}\mathbb{Z}$ Now look at

$$\psi := \lfloor y_1 \rfloor b_1 + \dots + \lfloor y_n \rfloor b_n$$

(here |x| denotes the integer part of x). The both  $\psi$  and  $\phi$  are algebraic integers (as the  $b_i$  are algebraic integers). Therefore, so is  $\theta = \phi - \psi$ . By construction,  $\theta$  has coefficients of the for  $\frac{x_i}{p} := y_i - \lfloor y_i \rfloor$  where  $x_i \in$  $\{0,\ldots,p-1\}$  and not all the  $x_i$  are zero (since, again, not all the  $y_i$  were in  $\mathbb{Z}$ ). This gives the first part of the lemma.

Now, assume  $x_i \neq 0$ , then let us replace  $b_i \in B$  with  $\theta$  to get a new basis B' which again consists of algebraic integers. Next, we note that the change of basis matrix from B to B' is

$$\begin{pmatrix} 1 & 0 & \cdots & \frac{x_1}{p} & \cdots & 0 \\ 0 & 1 & \cdots & \frac{x_2}{p} & \cdots & 0 \\ \vdots & \vdots & & \vdots & & \vdots \\ 0 & 0 & \cdots & \frac{x_n}{p} & \cdots & 1 \end{pmatrix}$$

(here the column of  $x_i/p$ 's is in the *i*-th column).

This matrix has determinant  $\frac{x_i}{p}$ . Therefore, by Proposition 2.6 we see that  $\Delta(B') = \frac{x_i^2}{p^2} \Delta(B)$ . But both  $\Delta(B), \Delta(B')$  are in  $\mathbb{Z}$  by Proposition 2.8, therefore  $p^2 \mid \Delta(B)$  giving the result.

**lemma 2.10.** Let  $K = \mathbb{Q}(\alpha)$  and  $\alpha$  be an algebraic integer such that  $m_{\alpha}$ satisfies Eisensteins Criterion for a prime p. Then none of the elements

$$\phi = \frac{1}{p}(x_0 + x_1\alpha + \dots + x_{n-1}\alpha^{n-1})$$

is an algebraic integer, where  $n = \deg(m_{\alpha})$  and  $x_i \in \{0, \dots, p-1\}$ .

*Proof.* We will only prove the case when  $m_{\alpha}$  is Eisenstein, since the proof of the more general case is identical.

Suppose for contradiction that  $\phi \in \mathcal{O}_K$  and let  $x_d$  be the first non-zero coefficient, so

$$\phi = \frac{1}{p}(x_d \alpha^d + x_{d+1} \alpha^{d+1} + \dots + x_{n-1} \alpha^{n-1}) \in \mathcal{O}_K.$$

Now, rewrite this as  $\phi = \frac{1}{p}(x_d\alpha^d + \alpha^{d+1}\beta)$  for some  $\beta \in \mathcal{O}_K$ . Next, multiply through by  $\alpha^{n-1-d}$ , then we have

$$\frac{x_d\alpha^{n-1}}{p} + \frac{\alpha^n\beta}{p} \in \mathcal{O}_K.$$

Now, since  $m_{\alpha}$  satisfies Eisenstein at p, we see that  $\alpha^n = pf(\alpha)$  for some  $f \in \mathbb{Z}[x]$  and therefore the above gives us that

$$\frac{x_d \alpha^{n-1}}{p} + \beta f(\alpha) \in \mathcal{O}_K.$$

and thus

$$\frac{x_d \alpha^{n-1}}{p} \in \mathcal{O}_K.$$

Lets now calculate the norm of this:

$$N_{K/\mathbb{Q}}\left(\frac{x_d\alpha^{n-1}}{p}\right) = \frac{x_d^n N_{K/\mathbb{Q}}(\alpha)^{n-1}}{p^n}.$$

By Eisenstein the constant coefficient of  $m_{\alpha}$  is divisible by p but not  $p^2$ , so since the constant coefficient of  $m_{\alpha}$  is  $N_{K/\mathbb{Q}}(\alpha)$  we see that  $N_{K/\mathbb{Q}}(\alpha) = pa$  where  $p \nmid a$ . Therefore we have

$$N_{K/\mathbb{Q}}\left(\frac{x_d\alpha^{n-1}}{p}\right) = \frac{x_d^n p^{n-1}a^{n-1}}{p^n} = \frac{x_d^n a^{n-1}}{p}.$$

But this cant be in  $\mathbb{Z}$  since p doesn't divide  $x_d$  or a, and this gives us a contradiction since Proposition 2.7 says that the norm of an algebraic integer must be an integer. So  $\phi$  couldn't have been an algebraic integer.

### 3. Cyclotomic fields

**lemma 3.1.** For n any integer,  $\Phi_n$  (the n-th cyclotomic polynomial) is an irreducible polynomial of degree  $\varphi(n)$  (where  $\varphi$  is Euler's Totient function).

**Theorem 3.2.** Let  $\zeta_p$  be a p-th root of unity for p an odd prime, let  $\lambda_p = 1 - \zeta_p$  and  $K = \mathbb{Q}(\zeta_p)$ . Then  $\mathcal{O}_K = \mathbb{Z}[\zeta_p] = \mathbb{Z}[\lambda_p]$  moreover

$$\Delta(\{1,\zeta_p,\ldots,\zeta_p^{p-2}\}) = \Delta(\{1,\lambda_p,\ldots,\lambda_p^{p-2}\}) = (-1)^{\frac{(p-1)}{2}}p^{p-2}.$$

*Proof.* First note  $[K : \mathbb{Q}] = p - 1$ .

Since  $\zeta_p = 1 - \lambda_p$  we at once get  $\mathbb{Z}[\zeta_p] = \mathbb{Z}[\lambda_p]$  (just do double inclusion). Next, let  $\alpha_i = \sigma_i(\zeta_p)$  denote the conjugates of  $\zeta_p$ , which is the same as the image of  $\zeta_p$  under one of the embeddings  $\sigma_i : \mathbb{Q}(\zeta_p) \to \mathbb{C}$ . Now by

Proposition 2.3 we have

$$\Delta(\{1, \zeta_p, \dots, \zeta_p^{p-2}\}) = \prod_{i < j} (\alpha_i - \alpha_j)^2 = \prod_{i < j} ((1 - \alpha_i) - (1 - \alpha_j))^2$$
$$= \Delta(\{1, \lambda_p, \dots, \lambda_p^{p-2}\})$$

Now, by Proposition 2.5, we have

$$\Delta(\{1,\zeta_p,\cdots,\zeta_p^{p-2}\}) = (-1)^{\frac{(p-1)(p-2)}{2}} N_{K/\mathbb{Q}}(\Phi_p'(\zeta_p))$$

Since p is odd  $(-1)^{\frac{(p-1)(p-2)}{2}} = (-1)^{\frac{(p-1)}{2}}$ . Next, we see that

$$\Phi_p'(x) = \frac{px^{p-1}(x-1) - (x^p - 1)}{(x-1)^2}$$

therefore

$$\Phi_p'(\zeta_p) = -\frac{p\zeta_p^{p-1}}{\lambda_p}.$$

Lastly, note that  $N_{K/\mathbb{Q}}(\zeta_p) = 1$ , since this is the constant term in its minimal polynomial. Similarly, we see  $N_{K/\mathbb{Q}}(\lambda_p) = p$ . Putting this all together, we get

$$N_{K/\mathbb{Q}}(\Phi_p'(\zeta_p)) = \frac{N_{K/\mathbb{Q}}(p)N_{K\mathbb{Q}}(\zeta_p)^{p-1}}{N_{K/\mathbb{Q}}(-\lambda_p)} = (-1)^{p-1}p^{p-2} = p^{p-2}$$

So the last thing we need to prove is that  $\mathcal{O}_K = \mathbb{Z}[\zeta_p]$ . From the calculation we just did, the only prime dividing the discriminant is p, therefore Lemma 2.9 tells us the only prime we need to check is p. But from Lemma 2.10 we know that dividing by p wont give us any new integral elements, so this must be an integral basis which give the result.

**Theorem 3.3.** Let n be a positive integer and  $\zeta_n$  a root of unity. If  $K = \mathbb{Q}(\zeta_n)$  then

$$\mathcal{O}_K = \mathbb{Z}[\zeta_n].$$

**lemma 3.4.** Let  $\alpha$  be an algebraic integer all of whose conjugates have absolute value one. Then  $\alpha$  is a root of unity.

**lemma 3.5.** Any unit u in  $\mathbb{Z}[\zeta_p]$  can be written in the form  $\beta \zeta_p^k$  with k an integer and  $\beta \in \mathbb{R}$ .

**lemma 3.6.** Let p be a prime and  $n = p^k$ . Then

$$p = u(1 - \zeta_n)^{\varphi(n)}$$

where  $u \in \mathbb{Z}[\zeta_n]^{\times}$ .

**lemma 3.7.** Let R be a Dedekind domain, p a prime and  $\mathfrak{a}, \mathfrak{b}, \mathfrak{c}$  ideals such that

$$\mathfrak{ab} = \mathfrak{c}^p$$

and suppose  $\mathfrak{a},\mathfrak{b}$  are coprime. Then there exist ideals  $\mathfrak{e},\mathfrak{d}$  such that

$$\mathfrak{a} = \mathfrak{e}^p \qquad \mathfrak{b} = \mathfrak{d}^p \qquad \mathfrak{ed} = \mathfrak{c}$$

#### 6

## 4. Fermats Last Theorem for regular primes

**Theorem 4.1.** Let  $p \geq 5$  be an prime number,  $\zeta_p$  a p-th root of unity and  $K = \mathbb{Q}(\zeta_p)$ . Assume that we have  $x, y, z \in \mathbb{Z}$  with gcd(xyz, p) = 1 and such that

$$x^p + y^p = z^p.$$

(1) We cannot have

$$x \equiv y \equiv -z \pmod{p}$$

and, therefore, without loss of generality, we may assume

$$x \not\equiv y \mod p$$
.

(2) For  $i \neq j$  we can write

$$(\zeta_p^i - \zeta_p^j) = u(1 - \zeta_p)$$

with u a unit in  $\mathbb{Z}[\zeta_p]$ . From this it follows that the ideals

$$(x+y), (x+\zeta_p y), (x+\zeta_p^2 y), \dots, (x+\zeta_p^{p-1} y)$$

are pairwise coprime.

- (3) Let  $\alpha \in \mathbb{Z}[\zeta_p]$ . Then  $\alpha^p$  is congruent to an integer modulo p.
- (4) Let  $\alpha \in \mathbb{Z}[\zeta_p]$ . Suppose that  $\alpha = a_0 + a_1\zeta_p + \cdots + a_{p-1}\zeta_p^{p-1}$  with at least one  $a_i = 0$ . Then there is an integer n such that  $\alpha/n \in \mathbb{Z}[\zeta_p]$ , then n divides each  $a_i$ .
- (5) Let  $i \in \{1, ..., p-1\}$ . Suppose that  $x + y\zeta_p^i = u\alpha^p$  with  $u \in \mathbb{Z}[\zeta_p]^\times$  and  $\alpha \in \mathbb{Z}[\zeta_p]$ . By considering  $x + y\zeta_p^i$  and its complex conjugate then there is an integer k such that

$$x+y\zeta_p^i-\zeta_p^{2k}x-\zeta_p^{2k-i}y\equiv 0\pmod p.$$

- Proof. (1) Reducing modulo p, using Fermat's little theorem, you get that if  $x \equiv y \equiv -z \pmod{p}$ , then  $3z \equiv 0 \pmod{p}$ . But since p > 3 this means p|z but this contradicts  $\gcd(xyz,p) = 1$ . Now, if  $x \equiv y \pmod{p}$  then  $x \not\equiv -z \pmod{p}$  we can relabel y,z so that wlog  $x \not\equiv y$  (this uses that p is odd).
  - (2) Lemma 3.6 gives that u is a unit. So all that needs to be proved is that the ideals are coprime. Assume not, then for some  $i \neq j$  we have some prime ideal  $\mathfrak p$  dividing by  $(x+y\zeta_p^i)$  and  $(x+y\zeta_p^j)$ . It must then also divide their sum and their difference, so we must have  $\mathfrak p|(1-\zeta_p)$  or  $\mathfrak p|y$ . Similarly,  $\mathfrak p$  divides  $\zeta_p^j(x+y\zeta_p^i)-\zeta_p^i(x+y\zeta_p^j)$  so  $\mathfrak p$  divides x or  $(1-\zeta_p)$ . We can't have  $\mathfrak p$  dividing x,y since they are coprime, therefore  $\mathfrak p|(1-\zeta_p)$ . But from lectures we know that since  $(1-\zeta_p)$  has norm p it must be a prime ideal, so  $\mathfrak p=(1-\zeta_p)$ . Now, note that  $x+y\equiv x+y\zeta_+p^i\equiv 0 \mod \mathfrak p$ . But since  $x,y\in\mathbb Z$  this means we would have  $x+y\equiv 0 \pmod p$ , which implies  $z^p\equiv 0 \pmod p$  which contradicts our assumptions.
  - (3) This is easy. Just use  $(x+y)^p \equiv x^p + y^p \pmod{p}$  and that  $\zeta_p$  is a p-th root of unity.
  - (4) Looking at  $\alpha = a_0 + a_1\zeta_p + \cdots + a_{p-1}\zeta_p^{p-1}$ , if one of the  $a_i$ 's is zero and  $\alpha/n \in \mathbb{Z}[\zeta_p]$ , then  $\alpha/n = \sum_i a_i/n\zeta_p^i$ . Now, as  $\alpha/n \in \mathbb{Z}[\zeta_p]$ , pick the basis of  $\mathbb{Z}[\zeta_p]$  which does not contain  $\zeta_p$  (which is possible

as any subset of  $\{1, \zeta_p, \dots, \zeta_p^{p-1}\}$  with p-1 elements forms a basis of  $\mathbb{Z}[\zeta_p]$ .]). Then  $\alpha = \sum_i b_i \zeta_p^i$  where  $b_i \in \mathbb{Z}$ . Therefore comparing coefficients, we get the result.

(5) Using lemma 3.5 we have  $(x + y\zeta_p^i) = \beta \zeta_p^k \alpha^p$  which is equivalent to  $\beta \zeta_p^k a \pmod{p}$  with a and integer (using part 3). Now, if we consider the complex conjugate we have  $\overline{(x + y\zeta_p^i)} \equiv \beta \zeta_p^{-k} a \pmod{p}$ . Looking at  $(x + y\zeta_p^i) - \zeta_p^{2k} \overline{(x + y\zeta_p^i)}$  then gives the result.

**Definition 4.2.** A prime number p is called regular if it does not divide the class number of  $\mathbb{Q}(\zeta_p)$ .

**Theorem 4.3.** Let p be a regular prime. Then

$$x^p + y^p = z^p$$

has no solutions with  $x, y, z \in \mathbb{Z}$  and gcd(xyz, p) = 1.

*Proof.* First thing is to note that if  $x^p + y^p = z^p$  then

$$z^p = (x+y)(x+\zeta_p y)\cdots(x+y\zeta_p^{p-1})$$

as ideals. Then since we know the ideals are coprime, then by lemma 3.7 we have that each  $(x+y\zeta_p^i)=\mathfrak{a}^p$ , for  $\mathfrak{a}$  some ideal. Note that,  $[\mathfrak{a}^p]=1$  in the class group. Now, since p does not divide the size of the class group we have that  $[\mathfrak{a}]=1$  in the class group, so its principal. So we have  $x+y\zeta_p^i=u_i\alpha_i^p$  with  $u_i$  a unit. So by 4.1 part 5 we have some k such that  $x+y\zeta_p-\zeta_p^{2k}x-\zeta_p^{2k-1}\equiv 0$  (mod p). If  $1,\zeta_p,\zeta_p^{2k},\zeta_p^{2k-1}$  are distinct, then 4.1 part 4 (which uses that p>3) says that p divides x,y, contrary to our assumption. So they cannot be distinct, but checking each case leads to a contradiction, therefore there can not be any such solutions.

**Theorem 4.4.** Let p be a regular prime. Then

$$x^p + y^p = z^p$$

has no solutions with  $x, y, z \in \mathbb{Z}$  and  $xyz \neq 0$ .

## References

[Mar18] Daniel A. Marcus. *Number fields*. Universitext. Springer, Cham, 2018. Second edition of [MR0457396], With a foreword by Barry Mazur.

[Sam70] Pierre Samuel. Algebraic theory of numbers. Translated from the French by Allan J. Silberger. Houghton Mifflin Co., Boston, Mass., 1970.

[Was82] Lawrence C. Washington. Introduction to cyclotomic fields, volume 83 of Graduate Texts in Mathematics. Springer-Verlag, New York, 1982.