


# Hurricane Wake Restratification Mechanisms

Haney S.<sup>1,2</sup>, B. Fox-Kemper<sup>1,2</sup>, S. Bachman<sup>1,2</sup>, B. Cooper<sup>1</sup>, S. Kupper<sup>1</sup>, K. McCaffrey<sup>1,2</sup>, L. Van Roekel<sup>2,3</sup>, S. Stevenson<sup>1,2,4</sup>, A. Webb, R. Ferrari<sup>5</sup>

1. University of Colorado, Boulder, CO
2. Cooperative Institute for Research in Earth Sciences (CIRES), Boulder, CO
3. Northland College, Northland, WI
4. University of Hawaii Honolulu, HI
5. Massachusetts Institute of Technology, Cambridge, MA

# How does a wake warm back up?

- **Goal:** make simple scalings for 1D, 2D, and 3D processes that restratify cold hurricane wakes.
- Scalings depend on readily available observations from satellites, profiling floats, and reanalysis data.
- Scalings for both the thin surface layer and for the sub-surface bolus are derived.



A cross section of the Typhoon Fanapi wake temperature. Image courtesy of Dr. Steve Jayne and the ITOP Group.

# Surface Scalings

$$\tau_{Ekman} = K_1 \frac{f H_{out} L_f}{u_*^2}$$

$K_1 = 2$

Ekman buoyancy fluxes  
(Thomas & Ferrari, 2008)

$$\tau_{SF} = K_2 H_{out}$$

$$K_2 = \frac{\rho C_p}{\lambda} \sim 1 \left( \frac{\text{days}}{m} \right)$$


surface, and penetrating solar  
buoyancy fluxes  
(Price et al., 2008, Kraus & Turner, 1967)

$$\tau_{eddy} = K_3 \frac{L_f^2 |f|}{H_{out} \Delta T}$$

$$K_3 = \frac{0.2 C_e}{g \alpha} \sim 6 \left( \frac{s^2 K}{m} \right)$$

eddy buoyancy fluxes  
(Fox-Kemper & Ferrari, 2008)

# Sub-Surface Bolus Scalings



$$\tau_{sb} = K_4 \frac{(H_{in} - H_{out})\Delta T}{\int_{-H_{in}}^{-H_{out}} (I_1 e^{k_1 z} + I_2 e^{k_2 z}) dz}$$

$$K_4 = \frac{\rho C_p}{S_0}$$

solar in the sub-surface  
bolus

$$\tau_{eb} = K_5 \frac{L_f^2 |f|}{\Delta T (H_{in} - H')}$$

$$K_5 = \frac{1}{7.11 C_e g \alpha}$$

eddies in the sub-surface  
bolus

# Ekman buoyancy fluxes are fastest

## Surface Timescales

Cyclone	$\tau_{\text{Ekman}}$ (days)	$\tau_{\text{SF}}$ (days)	$\tau_{\text{eddy}}$ (days)
Frances	6	30	525
Igor	2	26	287
Katrina	2	15	163


## Sub-Surface Bolus Timescales

Cyclone	$\tau_{\text{eb}}$ (days)	$\tau_{\text{sb}}$ (days)
Frances	122	435,000
Igor	39	325,000
Katrina	23	1,650

# Who Wins Under What Conditions?

surface fluxes beat wind


$$\frac{\tau_{Ekman}}{\tau_{SF}} = \frac{K_1}{K_2} \frac{f L_f}{u_*^2} > 1$$



eddies beat surface fluxes


$\otimes u_*$

$$\frac{\tau_{SF}}{\tau_{eddy}} = \frac{K_2}{K_3} \frac{H^2 \Delta T}{L_f^2 f} > 1$$



eddies beat wind

$$\frac{\tau_{Ekman}}{\tau_{eddy}} = \frac{K_1}{K_3} \frac{H^2 \Delta T}{L_f u_*^2} > 1$$




# Conclusions

- Restratification by Ekman buoyancy fluxes is the fastest mechanism in the thin surface layer for the wakes considered
- Restratification by eddy buoyancy fluxes is the fastest in the subsurface bolus.
- In the subsurface bolus restratification is generally slower, so temperature anomalies will persist
- Who wins may easily change if  $L_f$ ,  $H$ ,  $\Delta T$ ,  $u_*$ ,  $f$  change
- Eddies are particularly sensitive to  $H$  and  $L_f$ .



# References

- Fox-Kemper, B., R Ferrari (2008), Parameterization of mixed layer eddies. part II: prognosis and impact, J. Phys. Oceanography, 38, 1166-1179.
- Kraus, E. B., and J. S. Turner (1967), A one-dimensional model of the seasonal thermocline. II: The general theory and its consequences. Tellus, 19, 98–105.
- Price, J. F., J. Morzel, and P. P. Niiler (2008), Warming of SST in the cool wake of a moving hurricane, J. Geophys. Res., 113, C07010, doi:10.1029/2007JC004393.
- Thomas, L., R. Ferrari (2008), Friction, frontogenesis, and the stratification of the surface mixed layer, J. Phys. Oceanography, 38, 2501-2518.



U.S. Department of Commerce | National Oceanic & Atmospheric Administration | NOAA Research



**Earth System Research Laboratory**  
Global Monitoring Division

NNX09AF38G  
NNX09AO20H

0934737  
0825614