

YSC4204: STATISTICAL COMPUTING PROBLEM SET 2

EAST: SEAN SAITO, JOHN REID, HRISHI OLICKEL, CHONG WOON HAN

- (1) The C code to implement `meanC()` is as follows:

```
#include <stdio.h>
#include <R.h>

/*
 * Function that calculates the mean of vector x
 * Inputs:
 *   x    : vector of doubles
 *   n    : length of vector
 *   res  : placeholder for result
 */
void meanC(double *x, int *n, double *res) {

    int i;
    double temp = 0.0;
    for (i = 0; i < *n; i++) {
        temp += x[i];
    }

    *res = temp / *n;
    return;
}
```

We call this function in R using the following R Code:

```
# Automatically sets working directory to source file location
this.dir <- dirname(parent.frame(2)$ofile)
setwd(this.dir)

# Problem 1 - Writing a mean function in C to be used in R
meanR <- function(x) {
    # Placeholder for result
    res = 0.0
    n = length(x)

    dyn.load("mean.so")
    m = .C("meanC", x=as.double(x), n=n, res=res)

    return(m$res)
}
```

- (2) (a) With reference to Figure 2, let the red line (the "radius") be y_1 and let the blue line (the chord) be y_2 .

We need to find the point (p_1, q_1) . To do so, we will first find the lines y_1 and y_2 , then "step" from (p_0, q_0) to (p_1, q_1) by adding the distance from (p_0, q_0) to the midpoint.

$$\begin{aligned} y_1 &= \frac{q}{2p}x. && \text{(Since } y_1 \text{ bisects the perpendicular from } (p_0, q_0) \text{ to the } x\text{-axis)} \\ y_2 &= \frac{-2p}{q}x + c. && \text{Substitute in } (p, q) \\ q &= \frac{-2p}{q} \cdot p + c. \\ q^2 &= -2p^2 + qc. \\ c &= \frac{q^2 + 2p^2}{q}. \\ \Rightarrow y_2 &= \frac{-2p}{q}x + \frac{q^2 + 2p^2}{q}. \end{aligned}$$

We need to find the intersection of y_1 and y_2 , so we set $y_1 = y_2$.

$$\begin{aligned}\frac{qx}{2p} &= \frac{-2px}{q} + \frac{q^2 + 2p^2}{q}, \\ q^2x + 4p^2x &= 2pq^2 + 4p^3, \\ x(q^2 + 4p^2) &= 2pq^2 + 4p^3, \\ x &= \frac{2pq^2 + 4p^3}{q^2 + 4p^2}.\end{aligned}$$

To find y , we substitute x into the equation for y_1 ,

$$\begin{aligned}y &= \frac{q}{2px}, \\ &= \frac{q}{2p} \cdot \left(\frac{2pq^2 + 4p^3}{q^2 + 4p^2} \right), \\ &= \frac{2pq^3 + 4p^3q}{2pq^2 + 8p^3}, \\ &= \frac{q^3 + 2p^2q}{q^2 + 4p^2}.\end{aligned}$$

Now we need to find (p_1, q_1) . We know that the distance from (p_0, q_0) to the midpoint is equal to the distance from the midpoint to (p_1, q_1) . Breaking it into components:

$$\begin{aligned}p_1 &= p_0 + 2(p_2 - p + 0), \quad \text{where } p_2 \text{ is the } x\text{-coordinate of the mid point} \\ &= 2p_2 - p_0.\end{aligned}$$

Similarly, $q_1 = 2q_2 - q_0$.

$$\begin{aligned}p_1 &= 2p_2 - p, \\ &= 2\left(\frac{2pq^2 + 4p^3}{q^2 + 4p^2}\right) - p, \\ &= \frac{4pq^2 + 8p^3 - pq^2 - 4p^2}{q^2 + 4p^2}, \\ &= \frac{3pq^2 + 4p^3}{q^2 + 4p^2}, \\ &= \frac{p(4p^2 + q^2) + 2pq^2}{4p^2 + q^2}, \\ &= p + 2p\left(\frac{q^2}{4p^2 + q^2}\right), \\ &= p\left(1 + 2\left(\frac{\frac{q^2}{p^2}}{4 + \frac{q^2}{p^2}}\right)\right), \\ &= p\left(1 + 2\left(\frac{r}{4 + r}\right)\right), \\ &= p(1 + 2s).\end{aligned}$$

$$\begin{aligned}
q_1 &= 2q_2 - q_0, \\
&= 2\left(\frac{q^3 + 2p^2q}{q^2 + 4p^2}\right) - q, \\
&= \frac{2q^3 + 4p^2q - q^3 - 4p^2q}{q^2 + 4p^2}, \\
&= \frac{q^3}{q^2 + 4p^2}, \\
&= q \cdot \frac{q^2}{q^2 + 4p^2}, \\
&= q \cdot \left(\frac{q^2}{p^2} \cdot \frac{p^2}{4p^2 + q^2}\right), \\
&= q \cdot \left(\frac{r}{\frac{4p^2 + q^2}{p^2}}\right), \\
&= q \cdot \left(\frac{r}{4 + r}\right), \\
&= q \cdot s.
\end{aligned}$$

We've just shown that (p_1, q_1) are the same as the values of p and q after the $k = 1$ iteration of the loop. Since both points lie on the circle, $\sqrt{p_1^2 + q_1^2} = \sqrt{a^2, b^2}$. Hence we can consider this to be the loop invariant. For each iteration, the line from $(0, 0)$ to (p_k, q_k) remains on the circle while moving towards the x axis. This means

$$\lim_{k \rightarrow \infty} q_k = 0 \Rightarrow \lim_{k \rightarrow \infty} p_k = \sqrt{a^2, b^2}$$

. This becomes true when q_k becomes insignificant. In practice however, 3 iterations are necessary to approach decent accuracy, hence the limit of $k = 3$.

- (b) There are two cases where **pythag2** fails and **pythag** works:

```

x = 3e200
y = 4e200
pythag2(x,y)
> [1] Inf
pythag(x,y)
> [1] 5e200
x = 3e-200
y = 4e-200
> pythag2(x,y)
[1] 0
> pythag(x,y)
[1] 5e-200

```

The reason **pythag2** fails while **pythag** runs is due to possible overflow and underflow generated in the intermediate result of $x^2 + y^2$. For extremely large and small numbers, this value will limit the possible inputs of x and y to a range smaller than that of allowed by the floating points being used. **pythag** prevents this by iteratively approximating the value, where p approaches the result.

- (3) The standard Laplacian distribution is given by $f(x) = \frac{1}{2}e^{-|x|}$.

To obtain the inverse transform $F_X^{-1}(x)$, we must first take the integral of $f(x)$, and then find the inverse of the integral.

We will first separate $f(x)$ into cases

$$f_X(x) = \begin{cases} \frac{1}{2}e^x & \text{if } x < 0 \\ \frac{1}{2}e^{-x} & \text{if } x \geq 0 \end{cases}$$

Taking the integral, we get

$$F_X(x) = \begin{cases} \frac{1}{2}e^x & \text{if } x < 0 \\ 1 - \frac{1}{2}e^{-x} & \text{if } x \geq 0 \end{cases}$$

For the second case, the indefinite integral gives us $-\frac{1}{2}e^{-x} + c$, and we need $F_X(x)$ to be a cumulative distribution, so we have $c = 1$.

For the inverse transform, we must again do case analysis. For $F_X(x) = \frac{1}{2}e^x$:

$$\begin{aligned} y &= \frac{1}{2}e^x \\ 2y &= e^x \\ x &= \ln(2y). \end{aligned}$$

At $x = 0$, $F_X(x) = 0.5$, so this first case of the integral holds true for $x < 0.5$.

Case 2:

$$\begin{aligned} y &= 1 - \frac{1}{2}e^{-x} \\ y - 1 &= -\frac{1}{2}e^{-x} \\ 2 - 2y &= e^{-x} \\ \ln(2 - 2y) &= -x \\ x &= -\ln(2 - 2y). \end{aligned}$$

This holds true for $x \geq 0.5$.

Therefore, $F_X^{-1}(x)$ is given by the following

$$G_X(x) = \begin{cases} \ln(2y) & \text{if } x < 0.5 \\ \ln(2 - 2y) & \text{if } x \geq 0.5 \end{cases}$$

This is easy to translate into R.

```
# Define the standard laplacian pdf
laplacian_pdf <- function (x){
  return(0.5 * exp(-abs(x)))
}

# Define the laplacian cdf (not used)
laplacian_cdf <- function (x){
  return(ifelse(x < 0, 0.5*exp(x), 1 - 0.5*exp(-x)))
}

# Define the inverse transform of the laplacian cdf
inv_trf <- function (x){
  return(ifelse(x < 0.5, log(2*x), -log(2 - 2*x)))
}

# Generate the random sample
result <- inv_trf(runif(10000))

# Graph the sample
hist(result, probability = TRUE, breaks = 100,
      xlim = c(-5, 5), ylim = c(0,0.5))

# Overlay the line
x <- seq(-5,5,0.1)
y <- laplacian_pdf(x)
lines(x,y,col = "red")
```

The resulting histogram is shown below:

- (4) Suppose $x \sim \text{Rayleigh}(\sigma)$. The Rayleigh probability density function is

$$f(x) = \frac{x}{\sigma^2}e^{-x^2/(2\sigma^2)}, \quad x \geq 0, \sigma > 0.$$

The cumulative distribution function is

$$F_X(x) = \int \frac{x}{\sigma^2}e^{-x^2/(2\sigma^2)}dx,$$

Let $u = -x^2/(2\sigma^2)$,

$$\begin{aligned}\frac{du}{dx} &= \frac{-x}{\sigma^2}, \\ du &= \frac{-x}{\sigma^2} dx.\end{aligned}$$

By substitution,

$$\begin{aligned}F_X(x) &= \int -e^u du, \\ &= -e^u + c, \\ &= -e^{-x^2/(2\sigma^2)} + c.\end{aligned}$$

Observe that when $x = 1$, $F_X(x) = 0$. Therefore $c = 1$. Thus, we have the cumulative distribution for the Rayleigh distribution,

$$F_X(x) = 1 - e^{-x^2/(2\sigma^2)} \quad \text{for } x \in [0, \infty).$$

We derive $F_X^{-1}(x)$ as follows:

$$\begin{aligned}y &= 1 - e^{-x^2/(2\sigma^2)}, \\ e^{-x^2/(2\sigma^2)} &= 1 - y, \\ \frac{-x^2}{2\sigma^2} &= \ln(1 - y), \\ x &= \sqrt{-2\sigma^2 \ln(1 - y)}, \\ F_X^{-1}(x) &= \sqrt{-2\sigma^2 \ln(1 - y)}.\end{aligned}$$

For $u \sim \text{Uniform}(0, 1)$, and given that U and $1 - U$ have the same distribution, we can generate Rayleigh random variables by taking

$$F_X^{-1}(u) = \sqrt{-2\sigma^2 \ln(u)}.$$

We implement this equation in R to generate n random samples from a Rayleigh(σ) distribution as shown below:

```
## Define function to generate Rayleigh random variables ##
rray <- function(n, sigma){
  u <- runif(n)
  output <- sqrt(-2*sigma^2*log(u))
  return(output)
}
```

The code used to generate histograms is as follows:

```
## Set seed to ensure consistency before generating numbers and plotting ##
set.seed(0)

##### sigma = 0.5 #####
x1 <- rray(10000, 0.5)
hist(x1, prob=TRUE,
     main="Generated rayleigh random variables with sigma=0.5",
     xlab="x") # plot the histogram


## Superimpose a density line ##
xlines1 <- seq(0, round(max(x1)), 0.01)
yline1 <- (xlines1/0.5^2)*exp(-xlines1^2/(2*(0.5^2)))
lines(xlines1, yline1)

##### sigma = 2 #####
x2 <- rray(10000, 2)
hist(x2, prob=TRUE,
     main="Generated rayleigh random variables with sigma=2",
     xlab="x") # plot the histogram

## Superimpose a density line ##
xlines2 <- seq(0, round(max(x2)), 0.01)
yline2 <- (xlines2/2^2)*exp(-xlines2^2/(2*(2^2)))
lines(xlines2, yline2)
```

The resulting histograms are show below:

Generated rayleigh random variables with sigma=0.5



Generated rayleigh random variables with sigma=2

