

# Elementary Logic (PrepCamp)

Sebastiano Tronto

uni.lu

September 13-14, 2021

- 1 Statements
- 2 Logical operations
- 3 Implication
- 4 Quantifiers
- 5 Proofs

## Slides and exercises:

<https://github.com/sebastianotronto/preplogic>

## Email:

`sebastiano.tronto@uni.lu`

# Statements

- Unambiguous

- Unambiguous

Example (A bad joke)

**Q:** How many months have 30 days?

- Unambiguous

## Example (A bad joke)

**Q:** How many months have 30 days?

**A:** 11, some of them have even more!

- Unambiguous

## Example (A bad joke)

**Q:** How many months have 30 days?

**A:** 11, some of them have even more!

:- (

- Unambiguous

## Example (A bad joke)

**Q:** How many months have 30 days?

**A:** 11, some of them have even more!

:- (

- Objective



- Unambiguous

## Example (A bad joke)

**Q:** How many months have 30 days?

**A:** 11, some of them have even more!

:-(  
(

- Objective

## Example

**Good:** 3 is greater than 4

**Bad:** 3 is nicer than 4

- Mathematical: “*Three is greater than four*” (or “ $3 > 4$ ”)
- ...or not: “*I am 26 years old*”
- **Key point:** statements can be **true** or **false**

- We can combine statements to make new ones
- Negation (**not**), conjunction (**and**), disjunction (**or**)

# Negation (**not**)

If  $A$  is a statement, the statement “not  $A$ ” (in symbols:  $\neg A$ ) is **true** when  $A$  is **false**, and it is **false** when  $A$  is **true**.

# Negation (**not**)

If  $A$  is a statement, the statement “not  $A$ ” (in symbols:  $\neg A$ ) is **true** when  $A$  is **false**, and it is **false** when  $A$  is **true**.

## Example

$\neg(3 > 4)$  is equivalent to  $3 \leq 4$

“3 is not *greater than* 4” is equivalent to “3 is *less or equal than* 4”

# Conjunction (**and**)

The statement “ $A$  and  $B$ ” (in symbols:  $A \wedge B$ ) is **true** when both  $A$  and  $B$  are **true**, and it is **false** if at *at least* one of them is **false**.

# Conjunction (**and**)

The statement “ $A$  and  $B$ ” (in symbols:  $A \wedge B$ ) is **true** when both  $A$  and  $B$  are **true**, and it is **false** if at *at least* one of them is **false**.

## Example

“ $(3 < 4) \wedge (5 \text{ is an odd number})$ ” is **true**

## Example

“(Today is Monday)  $\wedge$  (we are in France)” is **false**

# Disjunction (**or**)

The statement “ $A$  or  $B$ ” (in symbols:  $A \vee B$ ) is **true** when at least one of  $A$  and  $B$  is **true**, and it is **false** if both of them are **false**.



# Disjunction (**or**)

The statement “ $A$  or  $B$ ” (in symbols:  $A \vee B$ ) is **true** when at least one of  $A$  and  $B$  is **true**, and it is **false** if both of them are **false**.

## Example

“(3 = 4)  $\vee$  (5 is an even number)” is **false**

## Example

“(Today is Monday)  $\vee$  (we are in Luxembourg)” is **true**

- **Important:**  $\vee$  is always *inclusive*:

- **Important:**  $\vee$  is always *inclusive*:

## Example (Another bad joke)

Waiter: "Would you like cheese or dessert?"

Mathematician: "Yes."

- **Important:**  $\vee$  is always *inclusive*:

## Example (Another bad joke)

Waiter: "Would you like cheese or dessert?"

Mathematician: "Yes."

- $\neg$  has precedence over  $\wedge$  and  $\vee$ :

$\neg A \wedge B$  means  $(\neg A) \wedge B$ ,       $\neg A \vee B$  means  $(\neg A) \vee B$

(or just use parenthesis)

If  $A$ ,  $B$  and  $C$  are statements:

$A \wedge B = B \wedge A$	$A \vee B = B \vee A$	<b>commutativity</b>
$A \wedge (B \wedge C) = (A \wedge B) \wedge C$	$A \vee (B \vee C) = (A \vee B) \vee C$	<b>associativity</b>
$A \wedge (B \vee C) = (A \wedge B) \vee (A \wedge C)$		<b>distributivity</b>
$A \vee (B \wedge C) = (A \vee B) \wedge (A \vee C)$		<b>distributivity*</b>
$\neg(\neg A) = A$		<b>double negation</b>
$A \wedge \text{true} = A$	$A \wedge \text{false} = \text{false}$	
$A \vee \text{true} = \text{true}$	$A \vee \text{false} = A$	
$(\neg A) \wedge A = \text{false}$	$(\neg A) \vee A = \text{true}$	
$\neg(A \wedge B) = (\neg A) \vee (\neg B)$	$\neg(A \vee B) = (\neg A) \wedge (\neg B)$	<b>De Morgan's laws</b>

- For simplicity: **true** = 1, **false** = 0

# Boolean algebra

- For simplicity: **true** = 1, **false** = 0
- We have a set  $\{0, 1\}$  with some operations  $(\wedge, \vee, \neg)$

- For simplicity: **true** = 1, **false** = 0
- We have a set  $\{0, 1\}$  with some operations  $(\wedge, \vee, \neg)$
- This is called a **Boolean algebra**



# Truth tables

A compact way of describing an operator, or a composition of operators

# Truth tables

A compact way of describing an operator, or a composition of operators  
Example:

$A$	$B$	$\neg A$	$A \wedge B$	$A \vee B$	$(A \vee B) \wedge (\neg A)$
0	0	1	0	0	0
0	1	1	0	1	1
1	0	0	0	1	0
1	1	0	1	1	0

# Truth tables

We can check that two statements are equivalent with truth tables

$A$	$B$	$\neg(A \wedge B)$	$(\neg A) \vee (\neg B)$
0	0	1	1
0	1	1	1
1	0	1	1
1	1	0	0

# Implication

- “ $A \implies B$ ” means “*If A (is true), then B (is true)*”

- “ $A \implies B$ ” means “If  $A$  (is true), then  $B$  (is true)”

## Example

*“If it rains, I will bring an umbrella”*

$(\text{It rains}) \implies (\text{I will bring an umbrella})$

# Implication

- “ $A \implies B$ ” means “*If A (is true), then B (is true)*”

## Example

*“If it rains, I will bring an umbrella”*

(It rains)  $\implies$  (I will bring an umbrella)

## Example

*“If my grandpa had wheels, he would be a bike”*

(My grandpa has wheels)  $\implies$  (My grandpa is a bike)

# Implication

- It is a logical operation: “ $A \implies B$ ” means “ $B \vee (\neg A)$ ”

# Implication

- It is a logical operation: " $A \implies B$ " means " $B \vee (\neg A)$ "

$A$	$B$	$A \implies B$	
0	0	1	No rain, I don't bring an umbrella
0	1	1	No rain, I bring an umbrella anyway
1	0	0	It rains, I don't bring an umbrella
1	1	1	It rains, I bring an umbrella



# Implication

- It is a logical operation: " $A \implies B$ " means " $B \vee (\neg A)$ "

$A$	$B$	$A \implies B$	
0	0	1	No rain, I don't bring an umbrella
0	1	1	No rain, I bring an umbrella anyway
1	0	0	It rains, I don't bring an umbrella
1	1	1	It rains, I bring an umbrella

## Remark

"**false**  $\implies A$ " is always **true**, whatever  $A$  is (*ex falso quodlibet*)

" $A \implies$  **true**" is always true, whatever  $A$  is

Sometimes we use the following symbols:

- “ $A \Leftarrow B$ ” is the same as “ $B \Rightarrow A$ ”
- “ $A \iff B$ ” is the same as “ $(A \Rightarrow B) \wedge (B \Rightarrow A)$ ”.  
It is read “ $A$  is equivalent to  $B$ ” or “ $A$  if and only if  $B$ ”.

- The statement  $(\neg B) \implies (\neg A)$  is called *contrapositive* of  $A \implies B$

# Contrapositive

- The statement  $(\neg B) \implies (\neg A)$  is called *contrapositive* of  $A \implies B$
- It is equivalent to “ $A \implies B$ ”

- The statement  $(\neg B) \implies (\neg A)$  is called *contrapositive* of  $A \implies B$
- It is equivalent to “ $A \implies B$ ”
- Two proofs:
  - 1 Properties of logical operations
  - 2 Truth tables

# See you tomorrow!

## Slides and exercises:

<https://github.com/sebastianotronto/preplogic>

## Email:

`sebastiano.tronto@uni.lu`

## Welcome Back!

### Slides and exercises:

<https://github.com/sebastianotronto/preplogic>

### Email:

`sebastiano.tronto@uni.lu`

Let  $S$  be a set and let  $A(x)$  be a “variable statement” that depends on  $x \in S$  (for example  $S = \mathbb{N}$  and  $A(x) = “x \text{ is an even number}”$ ).



Let  $S$  be a set and let  $A(x)$  be a “variable statement” that depends on  $x \in S$  (for example  $S = \mathbb{N}$  and  $A(x) = “x \text{ is an even number}”$ ).

- **Universal quantifier** ( $\forall$  or “for all”): “ $\forall x \in S, A(x)$ ” means that if we replace “ $x$ ” with any element of  $S$ ,  $A(x)$  is always **true**.
- **Existential quantifier** ( $\exists$  or “there exists”): “ $\exists x \in S, A(x)$ ” means that  $A(x)$  is **true** for at least one value of  $x$  in  $S$ .

Let  $S$  be a set and let  $A(x)$  be a “variable statement” that depends on  $x \in S$  (for example  $S = \mathbb{N}$  and  $A(x) = “x \text{ is an even number}”$ ).

- **Universal quantifier** ( $\forall$  or “for all”): “ $\forall x \in S, A(x)$ ” means that if we replace “ $x$ ” with any element of  $S$ ,  $A(x)$  is always **true**.
- **Existential quantifier** ( $\exists$  or “there exists”): “ $\exists x \in S, A(x)$ ” means that  $A(x)$  is **true** for at least one value of  $x$  in  $S$ .

**You always need a set  $S$**

# Quantifiers - examples

## Example

$S = \text{"the set of all cars"} , A(x) = \text{"x is red"}$

$\forall x \in S, A(x)$  is **false**.

$\exists x \in S, A(x)$  is **true**.

## Example

$S = \mathbb{N}, A(x) = x > 5$

$\forall x \in S, A(x)$  is **false**.

$\exists x \in S, A(x)$  is **true**.

**Today's most important fact:**

$$\neg(\forall x \in S, A(x)) = ?$$

$$\neg(\exists x \in S, A(x)) = ?$$

**Today's most important fact:**

$$\neg(\forall x \in S, A(x)) = \exists x \in S, \neg A(x)$$

$$\neg(\exists x \in S, A(x)) = \quad ?$$

## Example

$\neg$ “every number is even” = “there is at least one odd number”

**Today's most important fact:**

$$\neg(\forall x \in S, A(x)) = \exists x \in S, \neg A(x)$$

$$\neg(\exists x \in S, A(x)) = \forall x \in S, \neg A(x)$$

## Example

$\neg$ “every number is even” = “there is at least one odd number”

## Example

$\neg$ “ $\exists x \in \mathbb{N}, x + 3 = 9$ ” = “ $\forall x \in \mathbb{N}, x + 3 \neq 9$ ”

LUXEMBOURG

- A proof is a sequence of statements, each one logically deriving from the previous.

- A proof is a sequence of statements, each one logically deriving from the previous.
- Proofs are used to derive new statements from statements that are known to be true.



- A proof is a sequence of statements, each one logically deriving from the previous.
- Proofs are used to derive new statements from statements that are known to be true.
- If  $A$  is known to be true and the implication  $A \implies B$  is logically clear, then also  $B$  must be true.

- A proof is a sequence of statements, each one logically deriving from the previous.
- Proofs are used to derive new statements from statements that are known to be true.
- If  $A$  is known to be true and the implication  $A \implies B$  is logically clear, then also  $B$  must be true.
- Every mathematical theorem must be justified with a proof.

# Example: direct proof

## Theorem

*The sum of two even numbers is even.*

# Example: direct proof

## Theorem

*The sum of two even numbers is even.*

## Proof.

- 1 Recall the definition: a natural number  $x$  is called *even* if there is some natural number  $n$  such that  $x = 2n$ .

# Example: direct proof

## Theorem

*The sum of two even numbers is even.*

## Proof.

- 1 Recall the definition: a natural number  $x$  is called *even* if there is some natural number  $n$  such that  $x = 2n$ .
- 2 If  $x$  and  $y$  are even numbers, then there are natural numbers  $n$  and  $m$  such that  $x = 2n$  and  $y = 2m$ .

# Example: direct proof

## Theorem

*The sum of two even numbers is even.*

## Proof.

- 1 Recall the definition: a natural number  $x$  is called *even* if there is some natural number  $n$  such that  $x = 2n$ .
- 2 If  $x$  and  $y$  are even numbers, then there are natural numbers  $n$  and  $m$  such that  $x = 2n$  and  $y = 2m$ .
- 3 Then  $x + y = 2n + 2m = 2(n + m)$ .

# Example: direct proof

## Theorem

*The sum of two even numbers is even.*

## Proof.

- 1 Recall the definition: a natural number  $x$  is called *even* if there is some natural number  $n$  such that  $x = 2n$ .
- 2 If  $x$  and  $y$  are even numbers, then there are natural numbers  $n$  and  $m$  such that  $x = 2n$  and  $y = 2m$ .
- 3 Then  $x + y = 2n + 2m = 2(n + m)$ .
- 4 Then  $x + y$  is even.



# Proofs by contradiction

Idea: I want to show  $A = \mathbf{true}$ . I show that the implication  
“ $(\neg A) \implies \mathbf{false}$ ” is **true**. Then  $\neg A = \mathbf{false}$ , so  $A = \mathbf{true}$ .



# Proofs by contradiction

Idea: I want to show  $A = \mathbf{true}$ . I show that the implication “ $(\neg A) \implies \mathbf{false}$ ” is **true**. Then  $\neg A = \mathbf{false}$ , so  $A = \mathbf{true}$ .

## Definition

A natural number is called *prime* if it is different from 1 and it is only divisible by 1 and itself.

## Theorem

*There are infinitely many prime numbers.*

## Theorem

*There are infinitely many prime numbers.*

# Proofs by contradiction

## Theorem

*There are infinitely many prime numbers.*

## Proof.

- 1 Assume that there are only finitely many prime numbers.

# Proofs by contradiction

## Theorem

*There are infinitely many prime numbers.*

## Proof.

- 1 Assume that there are only finitely many prime numbers.
- 2  $\exists n \in \mathbb{N}$ , there are  $n$  prime numbers. Call them  $p_1, p_2, \dots, p_n$ .

# Proofs by contradiction

## Theorem

*There are infinitely many prime numbers.*

## Proof.

- 1 Assume that there are only finitely many prime numbers.
- 2  $\exists n \in \mathbb{N}$ , there are  $n$  prime numbers. Call them  $p_1, p_2, \dots, p_n$ .
- 3 Let  $u = p_1 \times p_2 \times \dots \times p_n + 1$ .

# Proofs by contradiction

## Theorem

*There are infinitely many prime numbers.*

## Proof.

- ① Assume that there are only finitely many prime numbers.
- ②  $\exists n \in \mathbb{N}$ , there are  $n$  prime numbers. Call them  $p_1, p_2, \dots, p_n$ .
- ③ Let  $u = p_1 \times p_2 \times \dots \times p_n + 1$ .
- ④  $u$  is not divisible by any of the prime numbers  $p_1, \dots, p_n$ .

# Proofs by contradiction

## Theorem

*There are infinitely many prime numbers.*

## Proof.

- ① Assume that there are only finitely many prime numbers.
- ②  $\exists n \in \mathbb{N}$ , there are  $n$  prime numbers. Call them  $p_1, p_2, \dots, p_n$ .
- ③ Let  $u = p_1 \times p_2 \times \dots \times p_n + 1$ .
- ④  $u$  is not divisible by any of the prime numbers  $p_1, \dots, p_n$ .
- ⑤ Therefore  $u$  is only divisible by 1 and itself. So  $u$  is prime.

# Proofs by contradiction

## Theorem

*There are infinitely many prime numbers.*

## Proof.

- ① Assume that there are only finitely many prime numbers.
- ②  $\exists n \in \mathbb{N}$ , there are  $n$  prime numbers. Call them  $p_1, p_2, \dots, p_n$ .
- ③ Let  $u = p_1 \times p_2 \times \dots \times p_n + 1$ .
- ④  $u$  is not divisible by any of the prime numbers  $p_1, \dots, p_n$ .
- ⑤ Therefore  $u$  is only divisible by 1 and itself. So  $u$  is prime.
- ⑥ So  $p_1, \dots, p_n$  are not the only prime numbers.





# Proofs by induction

If I want to prove  $\forall n \in \mathbb{N}, A(n)$ :

- 1 Prove  $A(0)$  (*base step*)
- 2 Prove  $\forall n \in \mathbb{N}, (A(n) \implies A(n+1))$  (*inductive step*)

# Proofs by induction

If I want to prove  $\forall n \in \mathbb{N}, A(n)$ :

- 1 Prove  $A(0)$  (*base step*)
- 2 Prove  $\forall n \in \mathbb{N}, (A(n) \implies A(n+1))$  (*inductive step*)

Theorem (Sum of natural numbers)

$$\forall n \in \mathbb{N}, \quad 0 + 1 + \cdots + n = \frac{n(n+1)}{2}$$

## Proof.

- 1 Base case:  $0 = 0$ .

## Proof.

- 1 Base case:  $0 = 0$ .
- 2 Let  $n$  be any natural number.

## Proof.

- 1 Base case:  $0 = 0$ .
- 2 Let  $n$  be any natural number.  
If  $A(n) = \mathbf{false}$ , then  $A(n) \implies A(n + 1)$  is **true**.

## Proof.

① Base case:  $0 = 0$ .

② Let  $n$  be any natural number.

If  $A(n) = \mathbf{false}$ , then  $A(n) \implies A(n+1)$  is **true**.

If  $A(n) = \mathbf{true}$ , we have to show that  $A(n+1) = \mathbf{true}$ .

## Proof.

① Base case:  $0 = 0$ .

② Let  $n$  be any natural number.

If  $A(n) = \mathbf{false}$ , then  $A(n) \implies A(n+1)$  is **true**.

If  $A(n) = \mathbf{true}$ , we have to show that  $A(n+1) = \mathbf{true}$ .

$$\begin{aligned} 0 + \cdots + (n+1) &= (0 + \cdots + n) + (n+1) = \\ &= \frac{n(n+1)}{2} + (n+1) = && \text{since } A(n) = \mathbf{true} \\ &= \frac{n^2 + n + 2n + 2}{2} \\ &= \frac{(n+1)(n+2)}{2} \end{aligned}$$

so  $A(n+1) = \mathbf{true}$



# The end

What you have learned in this course:

- Basic logic operations and logic implication
- Quantifiers and **their negation**
- Some basic mathematical proofs

**Slides and exercises:**

<https://github.com/sebastianotronto/preplogic>

**Email:**

`sebastiano.tronto@uni.lu`