

Rank-5 fusion rings

A reproducible census through multiplicity 19

Sébastien Palcoux

BIMSA

Computational companion, 19 September 2026

Abstract

We explain the exact enumeration of all based fusion rings of rank 5 through multiplicity 19, comprising 34,133 isomorphism classes. The account separates the mathematical coverage argument, the completed search, and independent verification of the emitted tensors. All duality types are included, and no categorifiability filter is applied. Code, machine-readable counts, full multiplication tensors and provenance records accompany the text.

AI assistance and verification. GPT-6 Astra Pro assisted with code, exposition and computational checks. Exhaustive-search arguments and exact independent tensor verification are recorded separately; no proof-assistant certification or peer-reviewed acceptance is claimed.

Scope. A census of based rings, not a classification of tensor categories. Only completed whole-rank results are released.

Contents

1	Objects, equivalence and the counting convention	3
1.1	Why duality types can be searched separately	3
2	The certified release at rank 5	3
3	An independent census	4
3.1	Commutativity and coordinates	4
3.2	Self-dual rings: reconstruction from a Krylov matrix	5
3.3	Every singular stratum	5
3.4	Nonself-dual rings and isomorphisms	6
3.5	The modular prefilter does not change the solution set	6
4	What the independent verification establishes	7
4.1	Provenance and reproducibility	7
4.2	Computation budgets and release gates	7
5	Reading and reusing the data	8
6	Reproduction commands	8

1 Objects, equivalence and the counting convention

A *fusion ring* here is a free abelian group with a distinguished finite basis $B = \{b_0, b_1, \dots, b_{r-1}\}$, an associative multiplication with unit b_0 , and an involution $i \mapsto i^*$ fixing 0. Its structure constants satisfy

$$b_i b_j = \sum_k N_{ij}^k b_k, \quad N_{ij}^k \in \mathbb{Z}_{\geq 0},$$

$$N_{0j}^k = N_{j0}^k = \delta_{jk}, \quad N_{ij}^0 = \delta_{i^*, j}, \quad (1)$$

$$N_{ij}^k = N_{i^*k}^j = N_{kj^*}^i. \quad (2)$$

These conventions agree with the usual based-ring definition; see [1, 2]. No assumption of categorifiability, commutativity or integrality of Frobenius–Perron dimensions is imposed unless proved in the relevant rank.

A based isomorphism is a bijection of the distinguished bases preserving the unit and multiplication. In tensor notation it is a permutation q fixing 0 such that

$$N_{ij}^k = N_{q(i)q(j)}^{q(k)} \quad \text{for every } i, j, k.$$

The coefficient of the unit determines the involution, so such an isomorphism automatically intertwines duality. The *multiplicity* is the maximum entry of the *full* ordered multiplication tensor. It is at least one, including for group rings. Put

$$c_r(m) = \#\{\text{rank-}r \text{ fusion rings of multiplicity exactly } m\} / \cong, \quad C_r(M) = \sum_{m=1}^M c_r(m).$$

A list through M contains all rings of multiplicity at most M , not only those of multiplicity equal to M . This distinction is essential when exporting OEIS b-files.

1.1 Why duality types can be searched separately

An involution on the $r - 1$ nonunit elements consists of p disjoint transpositions and $r - 1 - 2p$ fixed points, with $0 \leq p \leq \lfloor (r - 1)/2 \rfloor$. Fix one representative d_p of each type. An isomorphism between two tensors with involution d_p commutes with d_p . Hence it is sufficient to quotient that stratum by the centralizer

$$Z_{S_{r-1}}(d_p), \quad |Z_{S_{r-1}}(d_p)| = 2^p p! (r - 1 - 2p)!.$$

Different p cannot be isomorphic. Therefore a whole-rank census is complete exactly when every required stratum has been searched exhaustively. Finishing only the non-self-dual strata does not certify a whole-rank counting term.

2 The certified release at rank 5

The released bound is $M = 19$ and the cumulative number of classes is $C_5(19) = 34,133$. Every duality type is included. The accompanying full tensor file has one record per based-isomorphism class.

Exact multiplicity m	$c_5(m)$
1	16
2	37
3	82
4	134
5	209
6	336
7	477
8	733
9	863
10	1,082
11	1,383
12	1,948
13	2,211
14	2,554
15	2,936
16	4,157
17	4,255
18	5,080
19	5,640

The complete-run evidence is available through `results/census.json` and the corresponding reports under `verification/`. Earlier ranks retain their original completed-run provenance; repository checks and newly executed runs are labelled separately.

3 An independent census

3.1 Commutativity and coordinates

The following based-ring argument justifies the commutative specialization. Formal-codegree facts used here are recalled in [3].

Lemma 3.1. *Every rank-five fusion ring is commutative.*

Proof. The complex algebra of a Frobenius based ring is semisimple: the coefficient-of-unit functional gives the positive inner product making the basis orthonormal and multiplication a $*$ -representation. A noncommutative algebra of dimension five with a one-dimensional representation must be $\mathbb{C} \oplus M_2(\mathbb{C})$. Its unique one-dimensional character is Galois invariant, hence rational, and its values are algebraic integers, hence integers. This character is the Frobenius–Perron dimension character. Its values are positive. Consequently its formal codegree is an integer $D \geq 5$, the sum of five nonzero integer squares. If f denotes the formal codegree of the two-dimensional representation, the standard identity $\sum_E \dim(E)/f_E = 1$ gives

$$f = \frac{2D}{D-1} = 2 + \frac{2}{D-1}.$$

Formal codegrees are algebraic integers. The displayed rational number is not an integer for $D \geq 5$, a contradiction. $\square$

Fix the unit 0 and one of the three dualities

$$(0, 1, 2, 3, 4), \quad (0, 1, 2, 4, 3), \quad (0, 2, 1, 4, 3).$$

Write $C_{ijk} = N_{ij}^{k*}$. Commutativity and reciprocity say that C is a symmetric tensor, invariant under simultaneous duality. The nonunit triples therefore give respectively 20, 13, 10 variables, ordered lexicographically by their smallest orbit representative. All variables range independently in $[0, M] \cap \mathbb{Z}$ before associativity is imposed. The unit and duality determine the other entries. Commutativity of the five multiplication matrices is equivalent to associativity: the common unit is cyclic for the span of the matrices, and commuting matrices acting identically on the unit agree everywhere. Direct expansion gives 21, 12, 13 distinct quadratic equations, respectively. The supplement regenerates these polynomials from the axioms; no source census enters their construction.

3.2 Self-dual rings: reconstruction from a Krylov matrix

In the self-dual case use the ordered triples

$$\begin{aligned} &111, 112, 113, 114, 122, 123, 124, 133, 134, 144, \\ &222, 223, 224, 233, 234, 244, 333, 334, 344, 444 \end{aligned}$$

as coordinates $v_0, \dots, v_{19}$. By relabelling, arrange that v_1 is the minimum of the twelve mixed repeated-index coefficients, and $v_1 \leq v_2 \leq v_3$. The first nontrivial fusion matrix is

$$N_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & a & u^\top \\ 0 & u & Q \end{pmatrix}, \quad u = (v_1, v_2, v_3)^\top, \quad Q = \begin{pmatrix} v_4 & v_5 & v_6 \\ v_5 & v_7 & v_8 \\ v_6 & v_8 & v_9 \end{pmatrix}, \quad a = v_0.$$

For each of the nine bounded coordinates of (u, Q) set

$$K = [u, Qu, Q^2u], \quad s = u^\top u, \quad t = u^\top Qu, \quad H = Q^2 - aQ + uu^\top - I.$$

For $i = 1, 2, 3$, let T_i be the bottom-right 3×3 block of N_{i+1} ; thus $(T_i)_{jk} = C_{i+1, j+1, k+1}$. The vectors e_i below are the three standard coordinate vectors. Commuting with N_1 gives

$$\begin{aligned} T_i u &= H e_i, \\ T_i Q u &= (QH - sQ + u(Qu)^\top) e_i, \\ T_i Q^2 u &= (Q^2 H - sQ^2 - tQ + (Qu)(Qu)^\top + u(Q^2 u)^\top) e_i. \end{aligned} \tag{3}$$

These follow successively from the lower blocks of the commutator. If $\det K \neq 0$, multiplication by $\text{adj}(K)$ reconstructs all ten remaining coefficients, affinely in a . One integrality condition is a linear congruence $A + aB \equiv 0 \pmod{\det K}$. A gcd calculation gives every bounded solution for a , including the case $B = 0$. For each such solution all remaining divisibilities, coefficient bounds and all 21 associativity equations are checked. Thus there is no search over the last ten coefficients in this stratum.

3.3 Every singular stratum

The singular cases are essential to completeness. Multiplication by a ring element is zero if it annihilates the unit. Hence any polynomial annihilating the unit under N_1 annihilates N_1 itself. Apply this to the cyclic block generated by the unit.

If $\text{rank } K = 2$, write $Q^2 u = \alpha u + \beta Qu$ and $\lambda = \text{tr } Q - \beta$. The invariant rational subspaces of Q show that α, β, λ are integers. The remaining one-dimensional eigenspace of Q has eigenvalue λ . The cyclic block condition is

$$(\lambda^2 - a\lambda - 1)(\lambda^2 - \beta\lambda - \alpha) - \lambda(s\lambda + t - \beta s) = 0. \tag{4}$$

This is linear in a . All zero-coefficient cases are retained.

If $\text{rank } K = 1$, write $Qu = \lambda u$ and let $x^2 - Tx + U$ be the characteristic polynomial on $u^\perp$. The cyclic block polynomial is

$$p(x) = x^3 - (a + \lambda)x^2 + (a\lambda - 1 - s)x + \lambda.$$

For two distinct complementary eigenvalues require $x^2 - Tx + U \mid p$; for a repeated eigenvalue $\eta = T/2$ require $p(\eta) = 0$. These are again linear conditions on a , with no division by a potentially zero coefficient.

For both positive-rank singular cases, the 15 commutator equations linear in the ten remaining coordinates are solved by fraction-free elimination. Enumerate every bounded integral value of the free coordinates, back-substitute, and test all 21 quadratic equations.

If $u = 0$, the unit is annihilated by $N_1^2 - aN_1 - I$. For $a > 0$, $x^2 - ax - 1$ is irreducible over $\mathbb{Q}$, and a five-dimensional rational vector space cannot be a module over the resulting quadratic field. Thus $a = 0$ and $Q^2 = I$. A nonnegative integral symmetric orthogonal matrix is a permutation matrix. The transpositions are handled by the same linear completion. For $Q = I$, the remaining ten variables form the weighted rank-four associativity system: the constant 1 in each diagonal rank-four equation is replaced by 2, because both 1 and the invertible element contribute. The Diophantine elimination of the accompanying rank-four derivation applies with this substitution. All zero branches are implemented explicitly in `weighted.hpp`. For example, its terminal branch $b = c = e = 0$ gives $a = 1$, $d = f = 2$ and $gi + hj = h^2 + i^2 - 6$; it must not be omitted.

3.4 Nonself-dual rings and isomorphisms

For one dual pair, use the 13 orbit coordinates $v_0, \dots, v_{12}$. Put $A = v_6 - v_5$, $B = v_{10} - v_9$, $C = v_{11} - v_{12}$. Three associativity consequences are

$$\begin{aligned} Cv_5 &= Av_2 + Bv_4, & Cv_9 &= Av_4 + Bv_8, \\ v_{11}^2 &= 1 + v_{12}^2 - v_5^2 + v_6^2 - v_9^2 + v_{10}^2. \end{aligned}$$

Enumerate $v_5, v_6, v_9, v_{10}, v_{12}$ and recover v_{11} by an exact square test. Enumerate v_4 and solve for v_2, v_8 , retaining all free choices if A or B vanishes. Linear commutators determine or bound the four remaining variables v_0, v_1, v_3, v_7 ; check all equations. For two dual pairs the corresponding initial reductions are

$$v_0^2 = 1 + v_1^2 - v_2^2 - v_3^2 + 2v_4^2, \quad v_7^2 = v_2^2 - v_3^2 + v_5^2.$$

Two linear equations determine v_6 unless their coefficients vanish; that branch retains every bounded value. Solve for v_8, v_9 by linear completion and check all 13 equations. The orbit definitions in `census_system2.json` and `census_system4.json` fix these coordinate conventions exactly.

Finally minimize over the permutations preserving the fixed duality: 24, 4, 8 permutations in the three cases. Distinct duality types cannot be isomorphic. The independent verifier additionally minimizes every output tensor over all 24 unit-fixing permutations.

3.5 The modular prefilter does not change the solution set

In the nonsingular branch each reconstructed coefficient has the form $(A + Ba)/D$. When D is invertible modulo 257, an allowed integer coefficient must have a residue in $[0, M]$. Intersecting precomputed masks for two such coefficients eliminates impossible a values. Every retained value still undergoes exact divisibility and the original integer checks. If $257 \mid D$, the original gcd/congruence calculation is used. All integer-singular branches are unchanged. This is a necessary-condition speedup, not an assumption that a generic determinant is nonzero.

4 What the independent verification establishes

The enumerator and verifier have separate mathematical tasks. Exhaustiveness uses the complete coordinate parameterization, the safe-pruning proof and the normal completion of every required branch. The verifier reads only the emitted *full tensors*; it neither reconstructs the parameterization nor trusts the generator’s polynomial list.

For every tensor it checks nonnegative integral coefficients, both unit laws, uniqueness and involutivity of the dual, both reciprocity identities, and every integer associativity identity for all four indices. It then canonicalizes under *every* unit-fixing basis permutation. A matching hash is only a lookup aid: actual canonical tensors are compared. This detects both literal repetitions and different labellings of an isomorphic based ring.

Passing these checks proves soundness and uniqueness of the emitted data; by itself it would not prove that no ring was omitted. That is why normal completion logs and the mathematical coverage argument are retained separately. Comparison with the original ancillary file is an additional containment/regression test, not an input to enumeration and not a replacement for an exhaustiveness proof.

4.1 Provenance and reproducibility

The retained Vercleyen–Slingerland v4 ancillary file has SHA-256

b587c3f683157369da7cc090f762bbff0e18dcf11acc3f5031d6bc0c643843f8

Its 28,451 records contain 25,138 distinct classes. The 3,313 redundant records occur at rank five and exact multiplicities 9–12. Their original and corrected counts are respectively

$$(1463, 1794, 2283, 3049) \longrightarrow (863, 1082, 1383, 1948).$$

Every deletion has an explicit basis-permutation witness in the audit folder. This audit does not establish completeness of the source’s partial-search cells.

The primary manifest `results/census.json` records every released count, bound, filename and checksum. Compressed tensors use the original line-per-ring, nested-curly-brace format. An entry is $N[i][j][k] = N_{ij}^k$: the output index has already been dualized where the internal parameter convention requires it. The raw tensor needs no additional index conversion.

4.2 Computation budgets and release gates

The private repository provides two manually dispatched GitHub Actions campaigns. The original frontier has a 1,100-second shared driver deadline and a 20-minute job ceiling. The long frontier selects only ranks 5–8: each rank has one 4,200-second (70-minute) shared deadline and a 75-minute job ceiling, at most 300 runner-minutes in total. Compilation, all attempted multiplicities and duality types, export, compression and independent verification share the driver deadline. Enumeration receives 90% of remaining driver time; the balance is reserved for other phases. Threads follow the actual runner CPU allocation.

Only a complete whole-rank run with successful independent verification enters `best/`. A later incomplete attempt never replaces it. Candidates require explicit review and fresh independent verification before integration. Partial logs remain diagnostic evidence and never supply exhaustive census or OEIS terms. All workflows are manual; neither a push nor this manuscript launches a search.

The reviewed long campaign, GitHub run 35440299307 at source revision 931addc686e8 (full revision in the review record), completed rank five through multiplicity 19 with 34,133 classes,

including 5,640 at exact multiplicity 19. The subsequent rank-five bound 20 and the attempted bounds 7, 4 and 2 at ranks six, seven and eight exhausted their search budgets. They yield no larger complete whole-rank counts. Ranks three and four were not recomputed. The accepted rank-five data underwent a fresh exhaustive tensor and based-isomorphism audit on GitHub before integration. The full earlier exact-multiplicity prefix agrees. See `docs/CAMPAIGN_REVIEW.md` and `verification/imports/rank5_through19/`.

5 Reading and reusing the data

The result files count based rings, not monoidal categories or realizations of a ring. One tensor can admit several inequivalent categorifications, or none. No number here is a classification of fusion categories. The rank-specific count files use exact multiplicity; cumulative totals belong only in the summary column. Proposed OEIS additions must be regenerated from a complete manifest, not copied from an interrupted log.

The retained code was generated with AI assistance and is accompanied by explicit mathematical coverage arguments and independent integer checks. These checks are not a formal proof in a proof assistant. The present text is an explanatory computational companion, not a claim of journal acceptance or external peer review.

6 Reproduction commands

From the repository root, a complete run can be requested as follows. A timeout reports failure to finish, not a smaller theorem.

```
python3 code/run_census.py --rank 5 --bound 19 \
  --seconds 4200 --threads "$(nproc)" --verify --out rerun_rank5
```

To verify the supplied release without recomputing it:

```
python3 scripts/check_release.py --full
```

For a shared-budget frontier search, use `scripts/extend_census.py`. For ranks five through eight it starts at the released bound plus one and restarts the unfinished bound without claiming checkpoint resumption. The low-level command above limits enumeration only; the frontier wrapper additionally bounds every phase. Rank three is instead capped at 1000.

References

- [1] P. Etingof, S. Gelaki, D. Nikshych and V. Ostrik, *Tensor Categories*, Mathematical Surveys and Monographs 205, American Mathematical Society, 2015.
- [2] G. Vercleyen and J. Slingerland, *On Low Rank Fusion Rings*, Journal of Mathematical Physics 64 (2023), 091703; [arXiv:2205.15637v4](https://arxiv.org/abs/2205.15637v4).
- [3] V. Ostrik, *On formal codegrees of fusion categories*, Mathematical Research Letters 16 (2009), 895–901; [arXiv:0810.3242](https://arxiv.org/abs/0810.3242).
- [4] J. Dong and S. Palcoux, rank-four and rank-five census derivations, project manuscripts and computational supplements, September 2026. The relevant derivations are retained in the repository’s `methods/` directory.
- [5] OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, entries A348305, A354471, A354472, A354473, A354475, A354476 and A354477. Proposed updates in this repository are not submitted edits.