RANK-EIGHT GROTHENDIECK RINGS: CATEGORIFICATION THROUGH MULTIPLICITY THREE

Abstract. We investigate complex categorification of the supplied 4500 rank-eight fusion rings of multiplicity at most three. The exact-multiplicity-three continuation treats 3527 new tensors: 3479 general exclusions, 14 unitary models, and 34 unresolved cases. The complete classification through multiplicity two is preserved: 896 exclusions and 77 unitary models. New unified obstructions use quadratic-compositum compression, weak-integral and biquadratic rotation constraints, convex central induction, module-category ranks and Morita transport. Positive constructions include group representations, near-group equivariantizations, exceptional subfactor categories, a twisted group-theoretical category and Morita duals. All accepted models are compared with full source tensors. Exact replay certificates and individual candidate attempt inventories accompany the article.

Research status. AI-generated research · independent verification pending. The results, proofs and accompanying data have not been independently verified by a human expert or formalized in Lean. Treat all claims and data with caution; exact computational checks do not establish unstated categorical hypotheses.

1. Introduction and exact scope

The fusion-ring census [11] supplies all 96 rank-eight based fusion rings of exact multiplicity one and all 877 of exact multiplicity two, and the supplied extension adds 3527 of exact multiplicity three. A fusion ring need not be the Grothendieck ring of a fusion category. We work over $\mathbb{C}$ and do not assume pivotality, sphericality, braiding, commutativity or unitarity when generally excluding a ring. A positive construction is a unitary category with the exact based-isomorphism type of its full multiplication tensor.

The lower-rank criteria and their finite verifiers are developed in [12, 13, 14, 16, 17, 18]. We cite their theorems and do not reprove them. These new manuscripts are cited by title, with no byline. Our rank-eight observation, Proposition 4.1, extracts ordinary formal codegrees from the Casimir spectrum even though the ring may be noncommutative. Known near-group, zero-spectrum and cyclotomic tests provide other general obstructions [25, 28, 21]. Standard quantum group, pointed, Tambara–Yamagami and representation categories supply the positive side [29, 41, 22, 42].

The accepted study of integral Grothendieck rings [19, 20] supplies a necessary divisibility test and explicit group-theoretical rank-eight models and exclusions. Being an integral or Drinfeld ring does not itself supply a fusion category, so only its explicit categorical constructions enter our positive list.

Theorem 1.1 (Complete rank-eight classification through multiplicity two)**.** *The rank-eight based fusion rings through multiplicity two have the following classification.*

Exact multiplicity	*Fusion rings*	*Excluded generally*	*Unitarily realized*
1	96	48	48
2	877	848	29
Total	973	896	77

Every excluded ring is ruled out over $\mathbb{C}$ with no pivotality assumption. Each positive model is matched to the entire tensor under a unit-preserving permutation, and is therefore a Grothendieck ring. Thus

Date: 21 October 2026.
2020 *Mathematics Subject Classification.* 18M20, 18M15, 16T05.
Key words and phrases. Grothendieck ring, fusion category, formal codegree, categorification, exact certificate.

there are exactly 48 *Grothendieck rings of exact multiplicity one and* 29 *of exact multiplicity two, and every one admits a unitary realization.*

The general complex exclusion of `R8_2_24` is proved in Section 14. The ring `R8_1_3`, of global FP dimension 20, is realized below by a twisted group-theoretical category. Its dimension vector agrees with that of the commutative ring $\mathrm{Rep}(\mathrm{Dic}_5)$, although its multiplication is noncommutative. Thus full tensor comparison remains essential even after a suitable dimension profile is found.

2. Trusted input and stable records

A rank-r fusion ring has a distinguished basis $B = (b_0 = 1, b_1, \dots, b_{r-1})$, structure constants $N_{ij}^k \in \mathbb{Z}_{\geq 0}$, and a duality involution $i \mapsto i^*$ satisfying

$$b_i b_j = \sum_k N_{ij}^k b_k, \qquad N_{ij}^0 = \delta_{j,i^*}, \qquad N_{ij}^k = N_{i^*k}^j = N_{kj^*}^i.$$

Its exact multiplicity is $\max_{i,j,k} N_{ij}^k$. A based isomorphism permutes the basis, fixes b_0, and preserves every structure constant; it automatically preserves duality. We do not identify tensors by character table, spectrum or FP dimension vector alone.

The historical multiplicity-two source is the public *Fusion Ring Census* [11] at Git commit `b86eb6dc6eeeafe5fc5f6906d7b074e0322e9726`. It contains `results/rank8/tables.txt.gz` and `results/rank8/counts.csv`, with audit file `verification/rank8/audit.json`. The SHA-256 of the uncompressed tensor file is `97b86d2de8ca0352c5c6e14686cd1d43dbd52180dc53b2ff1ccf7a61788c9600`. The census verifies all duality types, associativity, nonnegative coefficients and uniqueness; its terminated-strata evidence justifies completeness through multiplicity two. We trust that source for counting and do not claim an independent exhaustive enumeration here.

Within each exact multiplicity m, sort full flattened $8 \times 8 \times 8$ tensors lexicographically and call the jth one $R8_m_j$. This is a deterministic local index, not an invariant of its original basis labelling. The files `data/decisions.json`, `data/realized_models.json` and `data/rank8_tables.txt.gz` give, respectively, one decision per input, the exact positive model tensors and the source multiplication tables. A SHA-256 of each JSON tensor prevents accidental source-record substitution. The comparison code minimizes the full tensor under all relabellings compatible with inexpensive intrinsic signatures; this is an isomorphism comparison, not a filter by numerical fingerprint.

3. The multiplicity-three continuation

The supplied extension has 3527 rings of exact multiplicity three and 4500 cumulatively. Every historical tensor, identifier, decision and model is preserved. The new raw tensor file has uncompressed SHA-256 `7baaa0a2018e11a0464c3f19da9030195421f04a73806c22363cf9d5eeb92a84`. The independently checked tensor axioms and preservation of earlier tensors do not constitute an independent exhaustive enumeration. The optimized producer source is now supplied and matches the completion receipt; enumeration completeness remains producer-reported.

The census release is pinned to Git commit `30046938fe2bd5cf4013c19eb0341d662f0d5dad`. The source and audit are `results/rank8/tables.txt.gz` and `verification/rank8/audit.json` in that repository.

Theorem 3.1 (Current exact-multiplicity-three status)**.** *The* 3527 *supplied rings of exact multiplicity three have the following recorded categorification status:*

Status	*Rings*
General complex exclusion	3479
Unitary construction	14
Unresolved general categorification	34

Every positive identification compares the entire ordered multiplication tensor under a unit-preserving bijection. A recorded construction proves existence, without claiming uniqueness up to tensor equivalence.

The source-bound combined ledger is `supplements/rank8/output/m3/ledger.json`. Its exact-M3 first-exclusion allocation follows. Overlapping obstructions are retained in the evidence rather than counted twice.

First accepted criterion	Rings
certified negative subring transfer	14
cyclotomic	7
dimension84 adjoint rigidity	5
dnumber	2575
global convex induction lift	2
global duality induction lift	1
integral drinfeld	34
involution module	19
lower rank subring	378
near group	1
nonsolvable dimension60 extension	1
one spectrum	1
oriented signed	67
quadratic casimir compression	64
rank7 certified subring	4
rank8 biquadratic lift	2
rank8 general biquadratic lift	1
rank8 noncomm dnumber	37
rank8 scalar oriented	3
rank8 spherical reduction	11
rank8 split module polynomial	44
rank two	10
rep a5 module rank	1
solvable dimension216 obstruction	4
solvable dimension80 rigidity	1
solvable dimension90 pointed base obstruction	2
split module dimension280 obstruction	2
subring dimension	6
weak group unitarity lift	1
weighted induction	2
zero spectrum	179

The lower-multiplicity classification below retains its original scope and proof inputs. Its historical allocation is separate from this table.

3.1. **Negative subring transfer and simultaneous quadratic fields.** The new census is first compared with the current, general-scope negative rank-six and rank-seven catalogues. Each transfer

certificate gives a dual-closed proper basis subset, verifies product closure, and supplies a unit-preserving permutation identifying every multiplication coefficient with the cited negative tensor. The tensor and its full published decision are hash bound. Any categorification would contain this full fusion subcategory, so the transfer excludes a general complex categorification. The recovered scan reproduces fourteen such transfers from catalogues of 531 rank-six and 1396 rank-seven negatives of multiplicity at most three.

Proposition 3.2 (Simultaneous quadratic compression)**.** *Let R be a fusion ring and $B = \sum_i N_i N_i^t$ its regular Casimir matrix. For each real quadratic field $L = \mathbb{Q}(\sqrt{d})$ appearing among its irreducible quadratic characteristic-polynomial factors, write*

$$Q_d = f_d \sum_j m_j \frac{t_j^2 - 4n_j}{n_j^2}, \qquad (x^2 - t_j x + n_j)^{m_j} \mid \det(xI - B),$$

where the sum includes exactly the factors generating L and f_d is the fundamental discriminant of L. Let r be the rank over $\mathbb{F}_2$ of all these quadratic squareclasses. A complex fusion categorification requires

$$2^{r-1} Q_d \le 4$$

for every displayed field. If a categorification would admit a Galois-conjugate pseudo-unitary form, the stronger bound is $2^{r-1} Q_d \le 1$.

Proof. The full-$I(\mathbf{1})$ compression theorem gives $[K : L] Q_d \le 1$ for a spherical categorification with dimension field K. On a simple representation E, the Casimir eigenvalue is $(\dim E) f_E$ and its regular multiplicity is $(\dim E)^2$; hence the displayed spectral multiplicities already give the correct contribution, also for a noncommutative ring. Each formal codegree, and therefore each Casimir eigenvalue, lies in K. The compositum of the displayed quadratic fields has degree 2^r, so $[K : L] \ge 2^{r-1}$. For an arbitrary category, pass to its sphericalization. Its original fusion-algebra sector has Casimir $2B$, dividing the known reciprocal-square compression contribution by four. This proves the first bound without assuming pivotality. A Galois-conjugate pseudo-unitary realization is spherical and yields the stronger bound directly. □

The implementation checks the stronger hypothesis by the existing weak-integrality or two-distinct-Frobenius–Perron-dimensions reduction. It computes every polynomial factor, rational ratio, squareclass rank, and gate from the source tensor. In particular a scoped spherical obstruction alone is never promoted to general nonexistence. The proof source and replay are `criteria/COMPRESSION_AND_TRANSFER.md` and `criteria/quadratic_compression.py`.

3.2. **Weakly integral quadratic trace fields.** The relative-trace and tensor-rotation engines are extended to the case where the object dimensions generate a real quadratic field but the total Frobenius–Perron dimension D is rational. Weak integrality supplies pseudo-unitarity. The prime support of the Frobenius–Schur exponent is the prime support of D; the quadratic conductor must be supported there. Write every dimension in the basis $1, \sqrt{d}$, retaining both coordinates even though $D = (D, 0)$. The central first- and second-moment and cyclic rotation equations remain identities in this field, with the same complete conductor-bounded relative-trace tables. The extension uses no division by the irrational coordinate of D.

All restriction-column choices and second-indicator assignments are retained. Numerical optimization only proposes a separator; exact integer replay verifies every inequality and the negative right-side pairing. The complete signed lifting remains required outside the independently checked general-to-positive gates. The additive adapter is `criteria/weak_integral_quadratic.py`; the existing frozen engines are left unchanged.

3.3. **Correlated traces in biquadratic dimension fields.** Let $K = \mathbb{Q}(\sqrt{q}, \sqrt{r})$, with coprime squarefree integers $q, r > 1$. For $s \in \{q, r, qr\}$ put $F_s = s$ if $s \equiv 1 \pmod 4$, and $F_s = 4s$ otherwise, and let χ_s be the corresponding primitive quadratic character. Coordinates below use $(1, \sqrt{q}, \sqrt{r}, \sqrt{qr})$.

Proposition 3.3 (Complete finite-height relative traces)**.** *The normalized relative trace of a primitive nth root to K has rational coordinate $\mu(n)/\varphi(n)$. Its $\sqrt{s}$ coordinate is zero unless $F_s \mid n$ and $\gcd(n/F_s, F_s) = 1$; otherwise it equals*

$$\frac{\epsilon_s \mu(n/F_s)\chi_s(n/F_s)\sqrt{F_s/s}}{\varphi(n)}.$$

There are four linked sign choices, with $\epsilon_{qr} = \epsilon_q\epsilon_r$. For the kth power of a primitive Nth root, set $g = \gcd(N, k)$ and $n = N/g$, use the same fixed sign pair, and multiply the quadratic coordinate by $\chi_s(k/g)$.

Suppose the prime support of twist orders is contained in P. For a height h, put

$$B_h = \prod_{p \in P} p^{\max(1, v_p(F_q), v_p(F_r), v_p(F_{qr})) + \lfloor \log_p h \rfloor}.$$

Every joint relative trace of powers $1, \dots, h$ is represented by an order dividing B_h, with one of the four linked sign choices, or by the joint zero vector.

Proof. The rational projection is the Ramanujan sum $\mu(n)$. The quadratic projections are the primitive quadratic Gauss sums $\tau(\chi_s) = \sqrt{F_s}$ followed by the coprime-part Ramanujan sum. Dividing by $\varphi(n)\sqrt{s}$ gives the displayed formula. The three signs come from one field automorphism and therefore satisfy the stated product relation. Reduction of the root order gives the power rule; independent signs for different powers would discard essential correlations.

If $v_p(N)$ exceeds the exponent in B_h, then for every $k \le h$ the p-valuation of $N/\gcd(N, k)$ exceeds one and every conductor valuation. The rational Möbius term vanishes. Each quadratic term vanishes either by noncoprimality with its conductor or by a zero Möbius value. All retained powers consequently have zero trace. Every other order divides B_h. □

For a pseudo-unitary category the spherical Cauchy theorem [61] supplies P from the norm of the global dimension, after the exact integrality and conductor-support checks. These are bounds on retained traces, not on the unknown categorical exponent. The field $\mathbb{Q}(\sqrt{2}, \sqrt{3})$ at height six gives $B_h = 288$; $(q, r) = (3, 7), (3, 5), (3, 13)$ at height four give respectively $1008, 720, 1872$ for the prime supports recorded in the certificates.

Weighted induction and exact separation. The full induction Gram matrix is computed from the tensor, including noncommutative rings. For each simple Grothendieck-algebra representation E, its distinguished central constituent has dimension D/f_E and multiplicity $\dim E$ in $I(1)$; the corresponding restriction column starts with $\dim E$. Its outer square contributes once to the Gram matrix. Thus a two-dimensional representation is handled with first entry two, rather than a commutative first-entry-one shortcut. All compatible low columns and all residual Gram-bounded nonnegative columns are enumerated. Their dimensions satisfy the necessary integrality and d-number tests.

Nonnegative weights for high columns and the complete linked trace types obey the Gram equations, indicator identities and cyclic-rotation traces on $\mathrm{Hom}(\mathbf{1}, X^m)$ through the chosen height. Every pseudo-unitary categorification supplies a feasible point. An exact rational Farkas witness therefore proves a contradiction; the numerical solver only proposes the witness.

General scope and independent checks. Such a rotation contradiction by itself has pseudo-unitary scope. General exclusion additionally checks every original dimension character and every oriented signed quotient in sphericalization, using exact character, parity and gauge-equivalence certificates. The accepted applications include `R8_3_347`, `R8_3_916` and `R8_3_1348`. The independent checker constructs the cyclotomic quotient $\mathbb{Q}[x]/\Phi_{B_h}$ and its quadratic Gauss sums, checks their squares and

correlated product, and compares every bounded-order trace and retained power with the formula. This verifies the finite identities; the preceding argument proves completeness beyond the box.

3.4. **A convex relaxation of central induction.** The following relaxation avoids enumerating all simultaneous choices of restriction columns for the constituents of $I(\mathbf{1})$. It uses the central-induction and relative-trace identities already established above.

Proposition 3.4 (Convex central-induction obstruction)**.** *Let R be a commutative fusion ring, let D be its Frobenius–Perron dimension, and suppose that its formal codegrees split in a real field K for which complete joint relative traces of the required roots of unity have been obtained. For each non-Frobenius–Perron character E, let $\mathcal{C}_E$ be the complete set of possible restriction columns of the central constituent of dimension $\beta_E = D/f_E$ of $I(\mathbf{1})$. Replace the choice of one column by variables*

$$y_{E,c} \geq 0, \qquad \sum_{c \in \mathcal{C}_E} y_{E,c} = 1.$$

Retain the Gram, first and second central trace, and tensor-rotation equations, using $y_{E,c} cc^t$ for the low Gram contribution and $y_{E,c} c_i \beta_E$ for each low twist moment. Fix the central unit column separately. For high columns, use every nonnegative integral column allowed by the full nonunit Gram block and the necessary central-dimension conditions, paired with every joint relative-trace type. If, for every admissible second-indicator assignment, the resulting rational system $Ax = b$, $x \geq 0$, has an exact integer separator $A^t u \geq 0$, $b^t u < 0$, then R has no pseudo-unitary categorification. If the complete spherical-character and oriented-sphericalization checks also succeed, then R has no complex fusion categorification.

Proof. A hypothetical pseudo-unitary category selects one actual column from each $\mathcal{C}_E$, hence supplies a one-hot solution for the corresponding simplex variables. Since R is commutative, each character labels a distinct simple constituent of $I(\mathbf{1})$ with multiplicity one and twist one. Thus its Gram and trace contributions are exactly those stated. Every actual high column is admitted by the full Gram block, whose complement is a sum of column squares. The complete relative-trace tables include every possible twist and cyclic-rotation spectrum. Consequently the category supplies a nonnegative solution of all the displayed equations. An exact separator contradicts $u^t Ax = u^t b < 0$. The final assertion uses the separately checked complete lifting argument; the convex relaxation itself does not assert pivotality or unitarity of an arbitrary fusion category. □

At height four this criterion excludes `R8_3_1453` generally. The relaxation has 15 777 variables and 127 equations; all sixteen second-indicator assignments have exact separators, and the full oriented lifting succeeds. Three fixed-low specializations reproduce the frozen Gram, moment, and rotation equations exactly. Known realized controls `R8_3_334`, `R8_3_342`, and `R8_3_475` survive. The proof, implementation, and validation receipts are stored in `criteria/GLOBAL_CONVEX_INDUCTION.md`, `criteria/global_induction.py`, and `criteria/global_specialization_R8_3_1453.json`. The same relaxation over $\mathbb{Q}(\sqrt{2}, \sqrt{3})$ excludes `R8_3_920` generally. Its height-four system has 97 768 variables and 197 equations; both admissible second-indicator assignments have exact separators and complete signed lifting succeeds. Three fixed-low specializations, selected from all 6891 low frames, reproduce 182 inherited equations each. See `criteria/global_specialization_R8_3_920.json`. The biquadratic dimension filters are additionally checked on 324 exact fixtures against independent rational multiplication-matrix characteristic polynomials, including the coprime-radicand fields used in the generalized trace adapter. One further necessary condition pairs high labels under duality. In the ribbon center the dual of a simple has the same dimension and twist, and its restriction vector is obtained by the source duality permutation. Thus labels (v, τ) and (v^*, τ) have equal multiplicity. Replacing their two columns by their sum gives a stronger nonnegative relaxation; all low and auxiliary rotation variables may remain unchanged. The complete high set and its involution are checked exactly. This refinement has a separate certificate kind and

isolated replay, preserving the earlier uncontracted certificates. Complete coefficient checks and known positive controls are recorded in `criteria/global_duality_validation.json`; the proof is `criteria/GLOBAL_DUALITY.md`. Resource limits or an unsuccessful exact reconstruction of a numerical separator provide no obstruction. This duality refinement excludes `R8_3_986` over $\mathbb{Q}(\sqrt{2},\sqrt{3})$. Its 197-equation system has 52 718 variables, compared with 97 768 before pairing the high labels. Both admissible second-indicator assignments have exact separators, and all sixteen oriented signed branches are excluded by the complete lifting certificate. The source-bound certificate is stored with the `global_family` campaign and has a separate isolated exact replay.

3.5. **A uniform split-module polynomial test.**

Proposition 3.5 (Fixed-point difference-space obstruction)**.** *Let R be a commutative fusion ring, g a nontrivial invertible basis element of order two fixing at least one basis element, and X a self-dual object. Let F be the basis elements fixed by g, and let p be any integer polynomial with $p(N_X)=0$. A categorification of R requires a symmetric integer matrix S, indexed by F, satisfying*

$$S_{V,W}\in\{-N_{X,V}^{W},-N_{X,V}^{W}+2,\dots,N_{X,V}^{W}\},\qquad p(S)=0.$$

Consequently, exhaustive failure of these equations, even after reduction modulo a prime, excludes a complex fusion categorification.

Proof. A simple object fixed by g supplies a rank-one module over the pointed subcategory generated by g, trivializing its associator class. Hence $A=\mathbf{1}\oplus g$ is a connected separable algebra. In $\mathcal{C}_A$, each fixed simple V splits into two module simples V_+,V_-, while each free g-orbit induces one module simple. The induction image is spanned by the free-orbit simples and the sums V_++V_-. The action of self-dual X is symmetric, so its orthogonal complement, spanned by V_+-V_-, is invariant. Induction and symmetry force the block between two fixed pairs to have the form $\left(\begin{smallmatrix}u & a-u\\ a-u & u\end{smallmatrix}\right)$, where $a=N_{X,V}^{W}$ and $0\le u\le a$ is an integer. Its action on differences has entry $2u-a$. The faithful regular representation makes $p(N_X)=0$ a ring identity, which holds on every module and hence on this invariant difference space. This proves the necessary condition. □

Swapping the two labels of a fixed pair conjugates S by a diagonal sign matrix. On a spanning forest of the odd-entry graph, all entries are nonzero and can therefore be made positive. This gives a complete sign reduction. The implementation verifies the source polynomial identity and every finite case exactly; a separate replay uses unbounded integer arithmetic and a different spanning forest. It gives new exclusions $R_{8,3,j}$ for $j\in\{1019,1057,1067,1197,2392,2836\}$, and independently recovers the applicable dimension-84 and dimension-280 obstructions. See `noncommutative/SPLIT_MODULE_POLYNOMIAL.md` and its source-bound certificates.

3.6. **Module ranks and dimension-sixty Morita transport.**

Proposition 3.6. *The ring `R8_3_307` admits no complex categorification.*

Proof. Its positive dimensions, in source order, are $(1,\sqrt{10},\sqrt{10},2\sqrt{10},5,3,3,4)$. The integral basis $\{0,4,5,6,7\}$ is the neutral part of a faithful C_2-grading and has dimension 60. It has no nontrivial proper fusion subring. Any neutral category $\mathcal{D}$ is therefore simple and integral, so [3, Theorem 9.12] gives $\mathcal{D}\simeq\mathrm{Rep}(A_5)$.

The odd component is an invertible $\mathcal{D}$-bimodule of rank three. By the subgroup/projective-representation classification of $\mathrm{Rep}(A_5)$-module categories [75, Theorem 2 and Remark 12], its objects have projective degrees $a(1,1,2)$: the exact action matrices have a one-dimensional simultaneous dimension eigenspace with primitive vector $(1,1,2)$. The certificate checks both its eigenvector equations and rank two of the combined coefficient matrix. For the defining subgroup $H\le A_5$, $|H|=6a^2\mid 60$, whence $a=1$ and $H\cong S_3$. A one-dimensional projective representation trivializes the multiplier.

The dual category has one family of simples for each H-double coset, parametrized by irreducible representations of its stabilizer. The three double cosets have sizes $6, 18, 36$ and stabilizers $S_3, C_2, 1$. Its rank is consequently $3+2+1 = 6$. Invertibility requires its dual to be $\mathcal{D}$, of rank five, a contradiction. The finite certificate checks every order-six embedding through Sylow-three normalizers and the complete double-coset calculation. □

Proposition 3.7 (Pointed transport from dimension sixty)**.** *Let $\mathcal{C}$ be faithfully graded by a finite group, with a nonsolvable neutral category of dimension* 60*. Then $\mathcal{C}$ is group-theoretical and integral. In particular* `R8_3_2132` *has no complex categorification.*

Proof. By [3, Theorem 9.16], the neutral category is weakly group-theoretical. Choose a nilpotent Morita representative. The orders of its extension groups multiply to 60. If all these groups were solvable, Morita invariance and [3, Proposition 4.5] would imply solvability of the neutral category. Every group whose order is a proper divisor of 60 is solvable, while the only nonsolvable group of order 60 is A_5. The representative must therefore be $\mathrm{Vec}^{\omega}_{A_5}$, with arbitrary admissible associator ω.

Transport the entire grading under this neutral Morita equivalence, using [3, Lemma 3.4]. Each component becomes an invertible bimodule over $\mathrm{Vec}^{\omega}_{A_5}$, whose one-sided dual is pointed. The pointed-dual criterion [79, Theorem 3.4] requires the subgroup defining the module category to be abelian and normal in A_5, hence trivial. Each component therefore has 60 simples and dimension 60, so all its simples are invertible. The transported category is pointed; the original category is group-theoretical and thus integral. No assumption $\omega = 1$ has been made.

For `R8_3_2132` the dimensions are $(1, 1, \sqrt{30}, 5, 5, \sqrt{30}, 2, 2)$. Its integral neutral category $\mathcal{D}$ has dimension 60, equals its adjoint subcategory, and has two invertibles. If it were solvable, its last cyclic-prime construction step could not be an extension, since $\mathcal{D}_{\mathrm{ad}} = \mathcal{D}$. It would be a C_2-equivariantization, by [3, Proposition 4.4]. Multiplication by the nontrivial character has neutral orbits $\{0, 1\}, \{3, 4\}, \{6\}, \{7\}$. A fixed underlying simple yields two equivariant simples interchanged by that character; a free underlying orbit yields one character-fixed simple of twice its dimension. De-equivariantization would have dimensions $1, 1, 1, 1, 1, 5$, the near-group ring $\mathrm{NG}(C_5, 4)$. Its whole tensor matches the already excluded `R6_4_19`; the dimension-thirty Morita certificate permits only simple dimensions one and two. Thus $\mathcal{D}$ is nonsolvable. The transport result makes the whole category integral, contradicting its odd dimension $\sqrt{30}$. □

The source-bound replay checks all dimension and grading equations, adjoint closure, character orbits, the complete lower-rank tensor match and the finite normal-subgroup premise. This is a new application and unification of the dimension-sixty dichotomy and integrality reductions; it does not identify an arbitrary categorification from its ring alone.

Theorem 3.8 (Dimension-80 ring rigidity)**.** *Every complex fusion category of type* $(1, 5; 5, 3)$ *has the based ring of* $\mathrm{Rep}((C_2)^4 \rtimes C_5)$ *for the fixed-point-free action. Its maximum fusion multiplicity is two. Hence* `R8_3_275`*,* `R8_3_276`*,* `R8_3_344`*, and* `R8_3_345` *are not categorifiable.*

Proof. The category is solvable by [3, Theorem 1.6]. Use its cyclic-prime extension/equivariantization sequence [3, Definition 1.2(ii), Proposition 4.4]. A final extension would have neutral dimension 40 or 16, neither of the form $a + 25b$ with $a \in \{1, 5\}$ and $0 \le b \le 3$. Thus $\mathcal{C} = \mathcal{D}^{C_5}$, since its invertibles force the equivariantization prime to be five, and $\mathrm{FPdim}\,\mathcal{D} = 16$.

A fixed base simple of dimension d contributes five simples of dimension d, and a free orbit contributes one of dimension $5d$; cyclic stabilizers have trivial projective multipliers. Base dimensions are rational algebraic integers, hence integral. A fixed dimension-five simple is impossible since $25 > 16$, and all free-orbit simples must be invertible. Thus $\mathcal{D} = \mathrm{Vec}^{\omega}_{H}$, $|H| = 16$, with only the unit fixed by C_5. Every invariant subset containing the identity has size one modulo five. The nontrivial characteristic center of the 2-group H therefore has order sixteen; then its characteristic subgroup $H[2]$ also has order sixteen. Hence $H = (C_2)^4$. The order-five action has irreducible minimal polynomial $\Phi_5(t) = t^4 + t^3 + t^2 + t + 1$ over $\mathbb{F}_2$, so it is unique up to conjugacy.

The five unit characters act trivially on the three free-orbit simples X_O, and $X_O^* = X_O$. Reciprocity gives $[X_O X_P : \chi] = \delta_{O,P}$ for every unit character χ. For a nonzero orbit Q, the coefficient at X_Q is the number of pairs $(x, y) \in O \times P$ with $x + y = z$, for any fixed $z \in Q$. This follows by forgetting equivariance and is independent of the associator and action coherence. The exact three-orbit convolution certificate has maximum two, whereas the four source tensors have maximum three. □

Theorem 3.9 (A dimension-216 solvability obstruction)**.** *No complex fusion category has type* $(1, 2; 2, 1; 3, 2; 8, 3)$*. In particular,* `R8_3_810`*,* `R8_3_812`*,* `R8_3_2492`*,* `R8_3_2494`*,* `R8_3_2620`*, and* `R8_3_2622` *are not categorifiable.*

Proof. Use the same solvability sequence. A final extension would have neutral dimension 108 or 72, neither expressible as $a + 4b + 9c + 64d$ with $a \in \{1, 2\}$, $b \le 1$, $c \le 2$, $d \le 3$. Thus $\mathcal{C} = \mathcal{D}^{C_2}$, with base dimension 108. The cyclic orbit rule forces precisely the base types

$$A = (1, 3; 3, 1; 4, 6), \qquad B = (1, 3; 3, 1; 4, 2; 8, 1).$$

Integrality descends by the orbit dimension formula and rational algebraic integrality. Neither base type permits subcategory dimension 54 or 36: enumerate $a + 9b + 16c + 64d$, $a \in \{1, 3\}$, within the displayed bounds. Hence $\mathcal{D} = \mathcal{E}^{C_3}$, with base dimension 36. Type B is impossible because its two dimension-four simples would have to arise from fixed objects in multiples of three. Type A forces $\mathcal{E}$ of type $(1, 4; 4, 2)$, and the actual C_3 action cycles its three nonidentity invertibles. Their group is therefore V_4.

For a dimension-four simple X, reciprocity gives $16 = |\operatorname{Stab}(X)| + 4m$. The stabilizer order divides four, so it is four. Thus all invertibles and a noninvertible belong to $\mathcal{E}_{\mathrm{ad}}$, whose dimension is 20 or 36. The ratio $\operatorname{FPdim}(\mathcal{E})/\operatorname{FPdim}(\mathcal{E}_{\mathrm{ad}}) = |U(\mathcal{E})|$ is integral; therefore $U(\mathcal{E}) = 1$. The final solvability step is consequently a C_2 equivariantization, giving a central Tannakian lift of one nontrivial invertible. Transport it under the actual C_3 action to obtain bosonic central lifts of all three nontrivial invertibles.

The kernel of $\operatorname{Inv} Z(\mathcal{E}) \to \operatorname{Inv} \mathcal{E}$ is $\operatorname{Aut}_{\otimes}(\mathrm{Id}_{\mathcal{E}}) = \operatorname{Hom}(U(\mathcal{E}), \mathbb{C}^{\times}) = 1$. Hence these lifts form a closed V_4, and its quadratic self-braiding form is identically one. This pointed braided subcategory is Tannakian. De-equivariantization gives $\mathcal{E} = \mathcal{F}^{V_4}$, where $\operatorname{FPdim} \mathcal{F} = 9$ and integrality again descends. The integral prime-square theorem [21, 22] makes $\mathcal{F} = \mathrm{Vec}_H^{\omega}$, with $H = C_9$ or C_3^2.

An orbit of length two has cyclic stabilizer and produces forbidden dimension-two simples. A nonidentity fixed point has stabilizer V_4; its projective degrees are either $1, 1, 1, 1$ or 2. These would produce four extra invertibles, beyond the four from the unit, or a forbidden dimension-two simple. Thus all nonidentity elements of H have free V_4 orbits. Each nontrivial involution therefore acts fixed-point-freely. An odd abelian group has only one fixed-point-free involution, namely inversion: $(f - 1)(f + 1) = 0$ and $\ker(f - 1) = 0$ imply $f = -1$. Three nonidentity elements of V_4 cannot all act by inversion. This contradiction proves the theorem. □

Theorem 3.10 (A split-module obstruction in dimension 280)**.** *The based rings* `R8_3_1051` *and* `R8_3_2415` *are not categorifiable over* $\mathbb{C}$*.*

Proof. The exact source certificate puts both rings in dimensions $[1, 1, 5, 5, 6, 8, 8, 8]$, with the nontrivial invertible g swapping the unit pair and the dimension-five pair and fixing the remaining four objects U_i. For a self-dual dimension-five object X, its coefficient block among the U_i is

$$R = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 2 & 1 \\ 1 & 2 & 0 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix},$$

and the source has the exact based-ring identity $p(X) = 0$, where $p(t) = t(t - 5)(t - 1)(t + 1)(t + 2)$.

The fixed dimension-six simple supplies a rank-one module category for $\mathrm{Vec}_{C_2}^{\omega}$, so ω is a coboundary and the regular algebra $A = 1 + g$ exists. In the category of right A-modules, the two free g-orbits

give two simples, and each fixed U_i splits into $U_{i,+}, U_{i,-}$; the stabilizer multiplier vanishes since $H^2(C_2, \mathbb{C}^\times) = 0$.

The action of self-dual X is symmetric. Induction is a module functor, so its block between two split pairs has equal row and column sums R_{ij}. That block is $\left(\begin{smallmatrix} u & R_{ij}-u \\ R_{ij}-u & u \end{smallmatrix}\right)$, with $0 \le u \le R_{ij}$. Coefficients connecting a free simple to the two members of a split pair agree, since the free simple is induced and induction sends a fixed object to the sum of its two summands. Thus the four pair differences form an invariant subspace.

After independently interchanging the two summands of each pair, the matrix on that subspace must be

$$S = \begin{pmatrix} a & 1 & 1 & 1 \\ 1 & 0 & b & c \\ 1 & b & 0 & d \\ 1 & c & d & e \end{pmatrix}, \qquad a, c, d, e \in \{-1, 1\}, \quad b \in \{-2, 0, 2\}.$$

Every module action must satisfy the same based-ring identity, hence $p(S) = 0$. The certificate evaluates the full polynomial over the integers for all 48 possibilities and records a nonzero entry in each case. It also verifies $p(N_X) = 0$ in each source regular representation, which is faithful. This gives the required contradiction without a unitarity assumption. □

Theorem 3.11 (Dimension-84 adjoint rigidity)**.** *No complex fusion category has dimension type* $(1, 4; 2, 2; 6, 2)$ *and* $\mathcal{C}_{\mathrm{ad}} = \mathcal{C}$*. Consequently the five tensors* `R8_3_76`*,* `R8_3_735`*,* `R8_3_738`*,* `R8_3_1422`*, and* `R8_3_1805` *are not categorifiable.*

Proof. By [77, Theorem 1.2], a category of dimension 84 is solvable or group-theoretical. Every group of order 84 is solvable: its Sylow 7 subgroup is normal and its quotient has order 12. Solvability is Morita invariant, so either alternative makes $\mathcal{C}$ solvable. Its last cyclic-prime construction step [3, Definition 1.2, Proposition 4.4] cannot be an extension because $\mathcal{C}_{\mathrm{ad}} = \mathcal{C}$. It is therefore an equivariantization $\mathcal{C} = \mathcal{D}^{C_2}$: the prime divides the number four of invertibles. The base has dimension 42 and is group-theoretical by [3, Theorem 9.2]. It is integral, since the orbit dimension formula makes its simple dimensions rational algebraic integers. Exactly two base invertibles are fixed by C_2. Let g be the sign object of the equivariantization. If n_2, n_6 count its fixed simples of dimensions two and six, respectively, then $n_2, n_6 \in \{0, 2\}$ and the base type is

$$(1, 2 + 2n_2;\ 2, (2 - n_2)/2;\ 3, 2n_6;\ 6, (2 - n_6)/2).$$

Thus the four possible multisets of base dimensions are $\{1, 1, 2, 6\}$, $\{1^6, 6\}$, $\{1, 1, 2, 3^4\}$, and $\{1^6, 3^4\}$. For completeness, all groups of order 42 are $C_7 \rtimes C_6$ with action image of order $1, 2, 3, 6$, and $C_7 \rtimes S_3$ with action image of order $1, 2$. Exhausting their subgroups and double cosets gives the following complete list of group-theoretical bases of the four types. The first and third types never occur. Write $F_{21} = C_7 \rtimes C_3$ and $F_{42} = C_7 \rtimes C_6$ for the faithful semidirect products. Type $\{1^6, 6\}$ occurs only for $(G, H) = (F_{42}, C_6)$ or (F_{42}, F_{42}). Type $\{1^6, 3^4\}$ occurs only for $G = F_{21} \times C_2$, with $|H| = 3, 6, 21, 42$, or $G = F_{42}$, with $|H| = 3, 21$, up to conjugacy. This finite audit includes arbitrary associators: every double-coset stabilizer has squarefree order and trivial Schur multiplier, by restriction/corestriction to its cyclic Sylow subgroups. Its projective character degrees are therefore ordinary character degrees. The exact subgroup and double-coset enumeration is part of the certificate. In the second type the category is $\mathrm{Rep}(F_{42})$. For $H = C_6$, the restriction $H^3(F_{42}, \mathbb{C}^\times) \to H^3(C_6, \mathbb{C}^\times)$ is an isomorphism: the remaining summand $H^3(C_7, \mathbb{C}^\times)^{C_6}$ vanishes because the faithful multiplier acts by its square. Thus the required trivial restriction kills the associator, and the split exact-factorization construction is $\mathrm{Rep}(F_{42})$; compare [64, Corollary 7.4]. In the fourth type the canonical adjoint subcategory is $\mathrm{Rep}(F_{21})$. For $H = C_3$ or F_{21}, support in the normal index-two subgroup F_{21} gives a faithful C_2 grading whose neutral category has type $(1, 3; 3, 2)$. It is $\mathrm{Rep}(F_{21})$ by [64, Theorem 6.3]. Since F_{21} is centerless, this neutral category equals its own adjoint, forcing equality with the adjoint of the whole category. For $G = F_{21} \times C_2$ and $H = C_6$ or G, coprime cohomology decomposition

instead gives $\mathrm{Rep}(F_{21}) \boxtimes \mathrm{Rep}(C_2)$, with the same adjoint. These reductions do not assume a trivial ambient associator. A representation category of a squarefree-order group has a unique fiber functor: the subgroup supporting a nondegenerate projective representation must have square order [75, Section 3.4, Theorem 2 and Example (i)]. Hence every tensor autoequivalence of $\mathrm{Rep}(F_{21})$ or $\mathrm{Rep}(F_{42})$ comes from a group automorphism. In the presentation $a^7 = b^m = 1$, $bab^{-1} = a^\lambda$, with $m = 3$ or 6 and λ of order m, an automorphism $a \mapsto a^u$, $b \mapsto a^v b^w$ forces $\lambda^w = \lambda$. It therefore fixes the abelianization and every linear character. The second base type consequently has six fixed invertibles; in the fourth type the canonical adjoint supplies at least three fixed invertibles. Both contradict the required number two. For each displayed source tensor, the certificate verifies all dimension equations, full adjoint closure, and every possible sign permutation directly in the given basis. □

Remark 3.12 (A restricted dimension-660 candidate). The residual ring `R8_3_731` is integral of dimension 660 and has no nontrivial proper closed based subring. Its exact primary-three witness excludes unitary categorification. By [3, Proposition 9.11], a hypothetical weakly group-theoretical categorification would be a representation category of a nonabelian finite simple group, hence unitary, a contradiction. Thus any categorification of this ring would be nonunitary and not weakly group-theoretical. Independently, the ordinary representation ring scan finds no matching tensor among the 40 groups of order 660. General complex categorification remains unresolved.

Theorem 3.13 (A dimension-90 solvability obstruction)**.** *A fusion ring of type* $(1, 2; 2, 4; 6, 2)$ *is not categorifiable if its adjoint subring is the whole ring and its nonidentity invertible fixes every noninvertible basis element. In particular,* `R8_3_2904` *and* `R8_3_3082` *admit no complex fusion categorification.*

Proof. Every category of dimension 90 is solvable [77, Corollary 4.16]. Its final cyclic-prime step [3, Proposition 4.4] cannot be an extension, since the universal grading is trivial. Two invertibles force this step to be a C_2 equivariantization, say $\mathcal{C} = \mathcal{D}^{C_2}$. The two lifts of a fixed simple of $\mathcal{D}$ are exchanged by the sign character. Thus each of the six sign-fixed noninvertibles of $\mathcal{C}$ comes from a free orbit. Halving their dimensions gives $\mathcal{D}$ of type $(1, 9; 3, 4)$ and dimension 45.

By [45, Theorem 1.1], an integral category of dimension $45 = 5 \cdot 3^2$ is group-theoretical: neither exceptional alternative applies. In fact such a category is pointed. All groups of order 45 have normal Sylow subgroups, so are C_{45} or $C_3^2 \times C_5$. In the presentation $\mathcal{C}(G, \omega, H, \psi)$, every stabilizer is H because G is abelian. A cyclic H has trivial Schur multiplier. A noncyclic H contains C_3^2 and has index 1 or 5. Restriction is then injective on the 3-primary component of $H^3(G, \mathbb{C}^\times)$, by restriction–corestriction. Since $[\omega]|_H = 0$, the remaining class is 5-primary, whereas $H^2(H, \mathbb{C}^\times)$ is 3-primary. The stabilizer-cocycle construction is multiplicative in ω, and the two ψ contributions cancel in cohomology because conjugation is trivial; see [46, Section 5.1]. Consequently every stabilizer multiplier is trivial. Its irreducible projective representations are therefore one-dimensional, and the double-coset index is one. The category is pointed, contradicting the four dimension-three simples forced above.

The exact receipt checks both source tensors, all dimension-character identities, the complete adjoint closure and sign actions. It is independent of the dimension-order linear-programming certificates. □

Proposition 3.14 (The Izumi–Xu inversion obstruction)**.** *The based ring* `R8_3_436` *has no unitary categorification. This statement does not exclude a general complex categorification.*

Proof. The source indices $\{0, 1, 2, 5\}$ form the near-group ring $\mathrm{NG}(C_3, 3)$, with g, g^2, X represented by source indices $1, 2, 5$, respectively, and

$$g^3 = 1, \qquad gX = Xg = X, \qquad X^2 = 1 + g + g^2 + 3X.$$

The invertible object with source index 3, denoted t, satisfies $t^2 = 1$ and $tgt^* = g^2$. The independent crossed-ring construction has model-to-source basis map $[0, 3, 1, 7, 2, 6, 5, 4]$; the certificate checks all 512 coefficients, as well as the displayed neutral subring and conjugation action.

A unitary categorification would therefore contain a unitary $\mathrm{NG}(C_3, 3)$ subcategory with a tensor autoequivalence interchanging its nontrivial invertibles. Izumi's C*-classification [36, Theorem 9.1 and Example 9.4], together with [36, Example 13.7], implies that every such unitary near-group category has trivial tensor automorphism group. This contradicts conjugation by t.

For the known Izumi–Xu categories, the same failure of the proposed construction also follows from the triviality of their outer tensor automorphism groups [83, Section 4.2, following Remark 4.10, and Remark 4.11]: inner conjugation acts trivially on their commutative based ring, whereas the required inversion does not. The C*-classification does not classify all complex near-group categories; the proposed nonunitary extension of its equations remains conjectural in [36, Remark 9.2]. Thus the present argument leaves the general complex categorification problem for this ring open. □

Lemma 3.15 (Graded ranks and fixed simples)**.** *In a faithful finite-group graded complex fusion category with neutral component $\mathcal{D}$ and $\mathrm{BrPic}(\mathcal{D}) = 1$, every homogeneous component has rank $\mathrm{rank}(\mathcal{D})$. If a pointed subcategory Vec_G^{ω} fixes a simple object, then $[\omega] = 0$. Both statements hold without unitarity.*

Proof. Each homogeneous component is an invertible $\mathcal{D}$-bimodule [7, Theorem 6.1(i)], hence belongs to the regular class; equivalence preserves rank. For the second assertion, the fixed simple spans a rank-one module category. Choosing isomorphisms for the group action gives scalar composition factors, and the module pentagon writes ω as their coboundary. □

Proposition 3.16 (The Haagerup graded-component obstruction)**.** *The ring `R8_3_455` has no unitary categorification. The argument does not exclude general complex categorification.*

Proof. The source basis splits into neutral indices $\{0, 1, 2, 3, 6, 7\}$ and odd indices $\{4, 5\}$, giving a faithful C_2-grading. The neutral ring has Haagerup rules

$$g^3 = 1, \qquad \rho g = g^{-1}\rho, \qquad \rho^2 = 1 + \rho + g\rho + g^2\rho.$$

The simple Y at source index 4 satisfies $gY = Yg = Y$ and $Y^2 = 1 + g + g^2$. The certificate independently reconstructs the full tensor and checks all 512 coefficients and the grading.

Let $\mathcal{D}$ be the neutral category in a hypothetical unitary categorification. The additive category generated by Y is a rank-one module over its pointed subcategory $\mathrm{Vec}_{C_3}^{\omega}$. Choosing isomorphisms $g \otimes Y \cong Y$, the module pentagon expresses ω as the coboundary of their composition scalars. Hence $c^{0,3}(\mathcal{D}) = 0$. Also $H^2(C_3, \mathbb{C}^\times) = 0$, so Izumi's second obstruction $c^{1,2}(\mathcal{D}) \in H^1(C_2, H^2(C_3, \mathbb{C}^\times))$ vanishes. These steps establish the hypotheses of [8, Definition 2.7]; omitting the first step would leave the twisted alternatives mentioned immediately after that definition.

By [8, Theorems 5.9 and 9.1], the unitary category $\mathcal{D}$ is one of the two known categories H_2, H_3. Their Brauer–Picard groups are trivial [88, Theorem 1.5]. Every homogeneous component of a faithful extension is an invertible bimodule over its neutral category [7, Theorem 6.1(i)]. The odd component must consequently be equivalent to the regular $\mathcal{D}$-bimodule, whose rank is six. Its rank is two, a contradiction. The classification invoked here is a C*-classification and supplies no such conclusion for hypothetical nonunitary neutral categories. □

3.7. **When a unitary obstruction is a general obstruction.** The negative primary-three form is ordinarily only an obstruction to unitary categorification. We use it for a general exclusion only after establishing that the hypothetical category is weakly group-theoretical. This reduction uses exact integer dimension below 84, solvability for dimensions with at most two prime factors, or a faithful grading with neutral category in that range; weak group-theoreticity is closed under extensions [3, Proposition 4.1 and the small-dimension theorems]. Every weakly group-theoretical category is

completely unitary [39, Theorem 5.20]. The primary-three contradiction then applies to its unitary realization and excludes general existence.

The source-bound verifier checks the dimension hypotheses, every nonzero structure constant of the grading, the neutral dimension and the grading dimension ratio before accepting this implication. In particular it uses no assertion that arbitrary weakly integral categories are unitarizable. When the reduction fails, the same negative form retains its unitary-only scope and does not change a general unresolved status.

3.8. **Source-bound identifications with known constructions.** The new positive inventory includes tensor products of previously constructed unitary categories and representation categories of finite groups. Each entry is checked by a unit-preserving basis bijection against all $8^3 = 512$ source coefficients, including noncommutative products. The certificate specifies the direction of the basis map. Equality of dimension vectors alone is never used as a model identification.

For the Frobenius-group entry `R8_3_342`, the finite orbit-convolution calculation constructs $\mathrm{Rep}(C_{13} \rtimes C_6)$ with its faithful action. The $\mathrm{PSL}_2(11)$ entry `R8_3_724` is reconstructed from its exact finite-group class algebra. The E_8 module-category entry `R8_3_3278` uses the standard unitary E_8 algebra and its module fusion rules [29]. The remaining existing near-group, Fibonacci and E_8 product entries use the unitary $\mathrm{NG}(C_3, 3)$ construction [78, Proposition A.4.1] or the adjoint of the unitary E_8 construction [29], followed by the indicated Deligne product. Their complete tensor comparisons are in the model catalogue. These comparisons establish the based-ring identification; categorical existence uses the cited construction and its stated scope.

3.9. **Unitary scope of the Izumi–Xu extension.** The fusion rules of `R8_3_448` are those of [78, Appendix A.6.1]; in particular $Vg = g^2V$ is essential. The unitary construction is supplied by the symmetrically self-dual 2221 planar algebra of [82, Section 5.1, Theorem 5.5]. The extension classification in [83, Theorem 4.9, Remarks 4.10–4.11] identifies the resulting graded extension with the displayed fusion rules: the two associator choices have the same tensor. Thus the earlier conservative pseudo-unitary label is upgraded to unitary existence.

3.10. **Two equivariantizations of a near-group category.** Use the normalized unitary $C_3 + 6$ solution of [36, Example 13.10], with $(x, y) = (\sqrt{\sqrt{3}/24}, 0)$. Lemma 13.1 describes actual automorphisms $\gamma_{\omega,\theta,u}$, with (ω, θ, u) and $(-\omega, \theta, -u)$ identical. The paragraph before Lemma 13.2 gives the strict conjugation tensor functor. Take $\theta = \mathrm{id}$ and either $(\omega, u) = (1, \mathrm{diag}(1, -1))$ or $(i, R_{\pi/2})$. Both have order two; the latter squares to $\gamma_{-1,\mathrm{id},-I} = 1$. Their actions on the six-dimensional self-fusion space have three eigenvalues of each sign. Their actions on the maps S_g have signs $+1$ and -1, respectively. Hence the complete rules are

$$(g,e)(h,f) = (g{+}h, e{+}f), \quad (g,e)X_f = X_f(g,e) = X_{e+f}, \qquad X_eX_f = \sum_{g\in C_3} (g, e{+}f{+}\epsilon){+}3X_0{+}3X_1.$$

The cases $\epsilon = 0, 1$ identify $R8_3_334, R8_3_274$ by all512 coefficients. These are coherent unitary actions, not abstract based-ring automorphisms.

3.11. **Cyclic permutation of three Fibonacci factors.** The canonical C_3 action on $\mathrm{Fib}^{\boxtimes 3}$ has two fixed simple triples, each splitting into three equivariant simples, and two free orbits. For a fixed output, the multiplicity is the character average on its fusion space; a free input orbit has zero nonidentity trace. For a free output the coefficient is the ordinary multiplicity at a representative. These formulas reconstruct all512 coefficients of `R8_3_475` and give its unitary realization.

3.12. **A Morita dual of the nonsplit $2D2$ extension.** In the known nonsplit extension `R8_2_37`, take the Q-system $A = 1 + r$ from [55, Lemma 4.1]. The neutral component of ${}_A\mathcal{E}_A$ is TT_3, with

basis $1, p, q, pq, qp, pqp$ and relations $p^2 = 1 + p$, $q^2 = 1 + q$, $pqp = qpq$; see [76, Definition 2.5]. The odd free-bimodule Gram matrix, in the order (a, a^3, ar, a^3r), is

$$\begin{pmatrix} 1 & 1 & 3 & 3 \\ 1 & 1 & 3 & 3 \\ 3 & 3 & 13 & 13 \\ 3 & 3 & 13 & 13 \end{pmatrix}.$$

Its nonnegative integral factorizations have the column $(1, 1, 3, 3)^T$ and either $(0, 0, 2, 2)^T$ or four copies of $(0, 0, 1, 1)^T$. After dividing underlying dimensions by $\mathrm{FPdim}(A)$, the latter would give four odd simples of dimension φ, but multiplication by p requires the unavailable dimension φ^2. The two remaining odd dimensions are therefore $2\varphi, 2\varphi^2$. Left and right multiplication by either p or q is forced to be $\left(\begin{smallmatrix} 0 & 1 \\ 1 & 1 \end{smallmatrix}\right)$. Reciprocity determines odd–odd products and yields precisely `R8_3_715`.

3.13. **The conformal G_2 algebra Morita dual.** Let $\mathcal{C} = \mathcal{C}(G_2, 3)$ and let A be the conformal algebra for $G_{2,3} \subset E_{6,1}$. The category ${}_A\mathcal{C}_A$ is unitary by [81, Theorems 2.21,2.28, Corollary 3.32]. Write $d = (3 + \sqrt{21})/2$. The two chiral Izumi–Xu subcategories have pointed local intersection C_3, by [78, Appendix A.4] and [80, Section 3.3, equation (16), Definition 3.12, Remark 3.13]. Their noninvertibles X, Y satisfy $X^2 = 1 + g + g^2 + 3X$ and $Y^2 = 1 + g + g^2 + 3Y$. Rigidity gives $(XY, XY) = 3$; dimension equality forces three further simples Z_i, each of dimension $d + 1$.

The g-action on this triple is nontrivial: otherwise the symmetric integral multiplicity block B has row sums three and satisfies $B^2 + J = 3I + 3B$. Its trace is six, so its off-diagonal sum would be three, a parity contradiction. Write $Z_i = g^i Z_0$ with Z_0 self-dual. Associativity gives $XY = YX = \sum_i Z_i$, $XZ_i = Z_iX = Y + \sum_j Z_j$, and $YZ_i = Z_iY = X + \sum_j Z_j$. The commutative orientation is impossible: the modular center has only self-dual simples, and [21, Lemma 8.49] would force all simples of the commutative based ring to be self-dual, contrary to $g^* = g^2$. Hence $Z_ig = g^{-1}Z_i$. Reciprocity reduces the remaining products to ten nonnegative triples with sum three; exact associativity leaves only $(1, 1, 1)$, so

$$Z_iZ_j = g^{i-j} + X + Y + \sum_k Z_k.$$

The reconstructed full tensor is `R8_3_458`.

3.14. **A twisted group-theoretical construction.**

Theorem 3.17 (A twisted A_6/A_5 model)**.** *There are μ_2-valued cochains $\omega \in Z^3(A_6, \mu_2)$ and $\psi \in C^2(A_5, \mu_2)$ with $d\psi = \omega|_{A_5}$ such that the unitary group-theoretical category $\mathcal{C}(A_6, \omega, A_5, \psi)$ realizes `R8_3_3342`. Its simple dimensions are $1, 3, 3, 4, 5, 10, 10, 10$.*

Proof. We give an exact finite certificate for the relative cocycle and then derive the fusion rules. Let $P = \langle r, s\rangle \cong D_8$ in A_6, on letters $0, \dots, 5$, with

$$r = (0123)(45), \qquad s = (02)(45).$$

Write $r^is^j = (i, j)$, and set $E = \{(i, j) : i \equiv j \pmod 2\}$, $K = \{(i, j) : i \equiv 0 \pmod 2\}$, and $t = s$. Then E fixes both 4 and 5, while K is the other Klein four subgroup. On K, put $x(i, j) = i/2$, $y(i, j) = j$, with values in $\mathbb{F}_2$. The normalized bar equations

$$dw = 0, \qquad w|_E = 0, \qquad w|_K = x^2y + xy^2 + dv$$

have an explicitly saved solution: there are 352 variables, 2455 equations, and rank 312.

For each of the 360 elements $g \in A_6$, the certificate supplies a normalized 2-cochain witnessing equality, modulo coboundaries, of w and its g-conjugate on $P \cap gPg^{-1}$. The intersection orders $1, 2, 4, 8$ occur $256, 64, 32, 8$ times, respectively. Thus the Cartan–Eilenberg stable-elements theorem [84, Chapter XII, Section 10] supplies a class $c \in H^3(A_6, \mathbb{F}_2)$ restricting to $[w]$. For $H = A_5$ fixing 5, E is a Sylow 2-subgroup of odd index 15; restriction and transfer give injectivity of $H^3(H, \mathbb{F}_2)$ into $H^3(E, \mathbb{F}_2)$. Hence $c|_H = 0$. Choose a normalized trivializing 2-cochain on H, extend it to A_6, and

gauge the global cocycle to vanish identically on H. Exponentiation by $a \mapsto (-1)^a$ gives $\omega|_H = 1$, so we may take $\psi = 1$.

The off-diagonal stabilizer is $L = H \cap tHt^{-1} = A_4$. The multiplier convention of [46, Section 5.1], restricted to the involutions $a, b \in E$, has additive form

$$\alpha(a,b) = w(a,b,t) + w(a,bt,t^{-1}bt) + w(abt,t^{-1}bt,t^{-1}at).$$

For $u = r^2$ and $v = rs$, the exact affine certificate proves $\alpha(u,v) + \alpha(v,u) = 1$: it gives a sum of original equations equal to this functional plus 1. The identity therefore holds for every solution of the displayed affine system, including the restriction of the globally gauged cocycle. This last restriction need only be cohomologous to the saved local representative. Consequently the multiplier has commutator -1 on a commuting pair, so its class in $H^2(A_4, \mathbb{C}^\times)$ is nontrivial. This relative computation is essential: the nonzero Dickson cubic itself maps to zero in $H^3(K, \mathbb{C}^\times)$.

There are two double cosets. The diagonal sector is $\mathrm{Rep}(A_5)$. The other sector has the three spin representations of the binary tetrahedral group $2T$:

$$\rho_i = \rho_0 \otimes \chi^i, \qquad i \in \mathbb{Z}/3, \qquad \det \rho_i = \chi^{2i},$$

where ρ_0 is the defining $SU(2)$ representation and χ generates the character group of A_4. The nontrivial μ_2 extension is unique: $H^2(A_4, \mathbb{F}_2)$ is the one-dimensional C_3-invariant part of $H^2(V_4, \mathbb{F}_2)$. The three spin representations have squared dimensions summing to 12, so they exhaust the projective simples. Their corresponding objects X_i have dimension $[A_5 : A_4] \dim \rho_i = 10$.

Every X_i is self-dual. Reversing the endpoints conjugates A_4 by an odd permutation and therefore inverts its character group. Representation duality also inverts determinants, so the two inversions cancel. All remaining transport scalars lie in μ_2 and hence do not change two-dimensional determinants. The three distinct determinants distinguish the X_i. The five diagonal simples are also self-dual; hence all fusion coefficients are symmetric in their three indices.

At distinct endpoints, composing two off-diagonal kernels sums over four intermediate points, a transitive A_4-set with stabilizer C_3. Thus the off-sector product is governed by induction from this C_3. There is a unique μ_2 trivialization of a multiplier on C_3, since $H^1(C_3, \mu_2) = H^2(C_3, \mu_2) = 0$. The order-three lift in $2T$ has eigenvalues ζ, ζ^2 on ρ_0, so $\rho_0|_{C_3} = \chi + \chi^2$. This is preserved by either automorphism of C_3; a residual μ_2 transport character is necessarily trivial. Equivalently, transport preserves the trivial determinant, whereas a nontrivial C_3 character twist would change it. Therefore

$$N_{X_0X_0}^{X_0} = \langle \chi + \chi^2, (\chi + \chi^2)^2 \rangle_{C_3} = 2.$$

For any pair i, j, the tensor of their two-character restrictions has multiplicities $(2,1,1)$ in some order. Pairing with the three spin restrictions gives off-sector multiplicities $(2,3,3)$ in some order. Symmetry and the anchor $N_{X_0X_0}^{X_0} = 2$ have a unique completion (enumerated among the 3^6 choices for unordered pairs):

$$N_{X_iX_j}^{X_k} = 3 - \delta_{i+j+k,0 \pmod 3}.$$

The restrictions of the five diagonal simples to A_4 are $1, 3, 3, 1+3, \chi+\chi^2+3$. As $3 \otimes \rho_i = \rho_0 + \rho_1 + \rho_2$, their actions on the three off-sector objects are $I, J, J, I + J, 2J - I$, where J is the all-ones matrix. Reciprocity supplies the diagonal coefficients of off/off products. The remaining tensor is the ordinary A_5 character tensor. This determines every coefficient.

The verifier constructs that tensor independently, checks the dimension and associativity equations, and matches all 512 coefficients to the authoritative source under the basis map $[0,1,2,6,7,3,4,5]$ from model to source, whose source-to-model inverse is $[0,1,2,5,6,7,3,4]$. All source hashes are recomputed. Finally, unit-modulus cocycles make the pointed category and its subgroup bimodule construction unitary. □

Remark 3.18. The comparison tensor `R8_3_3344` has $N_{XX}^X = 3$ for each dimension-ten simple. Thus it is not the ring of this particular construction. This comparison is not a general obstruction to other constructions of that tensor.

3.15. **Exact model inventory.**

Census ID	Construction	Scope
R8_3_274	Phase-rotation C_2-equivariantization of $\mathrm{NG}(C_3, 6)$	unitary
R8_3_331	$\mathrm{NG}(C_3, 3) \boxtimes \mathrm{Vec}(C_2)$	unitary
R8_3_334	Reflection C_2-equivariantization of $\mathrm{NG}(C_3, 6)$	unitary
R8_3_342	$\mathrm{Rep}(C_{13} \rtimes C_6)$	unitary
R8_3_448	Unitary Izumi–Xu C_2 extension (2221)	unitary
R8_3_458	Morita dual of the conformal $G_{2,3} \subset E_{6,1}$ algebra	unitary
R8_3_471	$\mathrm{NG}(C_3, 3) \boxtimes \mathrm{Fib}$	unitary
R8_3_475	$(\mathrm{Fib}^{\boxtimes 3})^{C_3}$ by cyclic permutation	unitary
R8_3_715	Morita dual of the nonsplit unitary $2D2$ extension	unitary
R8_3_724	$\mathrm{Rep}(\mathrm{PSL}_2(11))$	unitary
R8_3_1456	$\mathrm{Ad}(E_8) \boxtimes \mathrm{Vec}(C_2)$	unitary
R8_3_3278	Unitary E_8 fusion category	unitary
R8_3_3314	$\mathrm{Ad}(E_8) \boxtimes \mathrm{Fib}$	unitary
R8_3_3342	Twisted $\mathcal{C}(A_6, \omega, A_5, \psi)$ from a stable relative μ_2 class	unitary

Every model in this inventory admits a unitary realization. For R8_3_448, the original pseudo-unitary existence result is supplemented by the unitary 2221 planar-algebra construction above.

3.16. **The remaining general questions and replay scope.** There remain 34 unresolved exact-M3 tensors. The source tensors and per-case attempted criteria are recorded in `data/remaining_cases.py` and `evidence/rank8_m3_20261021/pipeline/residual_attempts.json`. The inventory distinguishes an exact contradiction, a necessary-condition survivor, a time limit, and an inapplicable test. Passing these conditions supplies no coherent associator; an unsuccessful construction search supplies no exclusion. Any claim of categorical novelty requires a separate comparison up to tensor equivalence.

3.16.1. *Individually investigated candidates.* These are unresolved categorification candidates, not asserted categories or established exotic examples. A unitary exclusion does not settle the general complex question. The last column gives ranks of nontrivial proper fusion subrings, as checked on the exact source tensors.

Census ID	Commutative	Invertibles	Unitary status	Proper subring ranks
R8_3_65	yes	4	unresolved	2, 4
R8_3_436	no	6	excluded	2, 3, 4, 6
R8_3_439	no	6	unresolved	2, 3, 6
R8_3_455	no	3	excluded	3, 4, 6
R8_3_567	no	6	unresolved	2, 3, 6
R8_3_721	no	1	unresolved	–
R8_3_731	yes	1	excluded	–
R8_3_901	yes	2	unresolved	2, 4
R8_3_903	yes	2	unresolved	2
R8_3_921	yes	2	unresolved	2
R8_3_925	yes	2	excluded	2
R8_3_926	yes	2	unresolved	2
R8_3_928	no	2	unresolved	2
R8_3_941	no	2	unresolved	2
R8_3_982	yes	2	unresolved	2
R8_3_985	yes	2	unresolved	2

Census ID	Commutative	Invertibles	Unitary status	Proper subring ranks
R8_3_1055	yes	2	unresolved	2
R8_3_1154	no	2	unresolved	2
R8_3_1203	yes	2	unresolved	2
R8_3_1350	yes	2	excluded	2
R8_3_1463	yes	2	unresolved	2, 4
R8_3_1615	yes	2	unresolved	2
R8_3_1617	yes	2	unresolved	2
R8_3_1660	yes	2	unresolved	2
R8_3_1661	yes	2	excluded	2
R8_3_1739	yes	2	unresolved	2
R8_3_1768	yes	2	unresolved	2
R8_3_1795	yes	2	unresolved	2
R8_3_2842	yes	2	unresolved	2
R8_3_2885	yes	2	excluded	2
R8_3_3258	yes	1	excluded	–
R8_3_3344	yes	1	unresolved	5
R8_3_3413	yes	1	excluded	–
R8_3_3472	yes	1	unresolved	–

Asaeda–Haagerup and recent odd-order constructions. The ring R8_3_721 has the exact Asaeda–Haagerup global dimension $136 + 32\sqrt{17}$, so a dimension mismatch does not dispose of this lead. The complete known Morita class nevertheless gives no model: AH_1, AH_2, AH_3 have ranks $6, 9, 9$, respectively [85, Section 4.3, Tables 1–3], while AH_4, AH_5, AH_6 have four invertibles and maximum multiplicity two [9, Lemmas 3.4, 3.7, 3.15 and Theorem 4.21]. The source ring has one invertible and maximum multiplicity three. The audit independently reconstructs the two rank-eight AH based tensors and matches all 512 coefficients to the existing positive controls R8_2_39 and R8_2_484. Of the 277 baseline survivors, fourteen have the AH global dimension, checked using exact positive dimension characters; none is a member of this known family.

The September 2026 constructions [86, Theorem 2.1] and [87, Theorems A, 4.4 and 6.3] were also checked. For odd $N \geq 3$, write $d^2 = Nd + 1$ with $d > 0$. The HI family has global dimension $D_N = N(1 + d^2)$, and the specified near-group family has global dimension ND_N. Exact Casimir characteristic-polynomial division, supplemented by integer spectral bounds, rules out both dimensions for every baseline survivor and every odd N; the bounds $D_N > N^3$ and $ND_N > N^4$ certify the infinite tail. This also rules out their Morita duals, since Morita equivalence preserves FP dimension. The cited general existence statements are spherical and pseudo-unitary; the stated Morita classification has additional unitary, cyclic, untwisted and Q-system hypotheses. These completed model-family checks make no general nonexistence claim for the remaining rings. The aggregate command is

`python supplements/rank8/software/verify_m3_continuation.py`.

It checks the accepted ledger against the source, baseline criteria, continuation certificates and explicit model tensors. Its full receipt binds the exact ledger digest, manifests and proof-source hashes. Under Python optimization, inherited assertion-based checkers run in an isolated interpreter with assertions enabled. Metadata-only mode checks these identities and reconciliation without replaying the mathematics.

4. Historical multiplicity-two criteria and their exact application

4.1. **Short tensor and subring tests.** If X is self-dual and $X^2 = 1 + kX$ with $k \geq 2$, its span with 1 is a fusion subring that cannot arise over $\mathbb{C}$ by [12, 23]. When exactly seven basis elements

are invertible and one additional object X satisfies

$$gX = Xg = X, \qquad X^2 = \sum_{g \in G} g + \kappa X,$$

the rank-eight near-group case with $\kappa = 1$ or 2 is commutative and satisfies $0 < \kappa < |G| - 1$. Schopieray's general Proposition 3.2 [25] excludes it. The machine certificate checks the complete group and all displayed equations, not just the value of κ.

The zero-spectrum test is the finite nine-index condition of [28, Theorem 2.14] in the form reproduced in [17, Section 3]. It is a necessary condition for an associative tensor lift, with no pivotality assumption. Its C++ scanner searches the nine indices on the full tensor and the Python checker independently recomputes every coefficient, positivity condition and vanishing contraction. The native arrays are sized for rank eight, and the checker has no rank-six restriction.

Every duality-closed, multiplication-closed based subset of a category gives a fusion subcategory. We compare every such proper subring of ranks two through six against the *complete positive list* in the corresponding lower-rank theorem. The lower-rank classifications are unrestricted for ranks two, three and four, and complete through multiplicity 20 for rank five and 2 for rank six [12, 13, 14, 16, 17]; every rank-eight subring in this historical multiplicity-two stage also has multiplicity at most two. A rank-seven subring is not excluded merely because it is absent from an incomplete rank-seven list.

The rank-seven analysis [18] provides four specific, fully checked negative multiplicity-two tensors that occur as rank-eight closed subrings. We transfer their exact signed-sphericalization witnesses by complete seven-dimensional based isomorphism and recheck all four proof trees. This separate stage gives exclusions only for rings actually containing one of those four proved negative subrings; it makes no claim that the rank-seven positive catalogue is exhaustive.

The involution-module theorem of [17, Section 5] says that an invertible involution fixing basis elements forces a finite anti-invariant module representation. Where its stated hypothesis is met, the script searches *all* bounded integer matrices and verifies multiplication, transpose, parity and entry bounds. The FP dimension quotient of a hypothetical fusion subcategory must likewise be an algebraic integer [22, Proposition 8.15]; a rational coefficient in its exact minimal polynomial excludes categorification.

4.2. **A rank-eight noncommutative codegree lemma.** For any fusion ring, write

$$\Omega = \sum_i b_i b_{i^*}, \qquad C_R = \sum_i N_i N_i^{\mathsf{T}}.$$

For an irreducible representation E of $R \otimes \mathbb{C}$ of degree e_E, the central element Ω acts as the scalar $e_E f_E$, where f_E is its ordinary formal codegree. The regular representation contains E with multiplicity e_E, so that scalar has multiplicity e_E^2 in the characteristic polynomial of C_R [24, 17].

Proposition 4.1 (Noncommutative rank-eight codegrees)**.** *Let R be a noncommutative fusion ring of rank eight. Its semisimple complex algebra has Wedderburn decomposition $\mathbb{C}^4 \oplus M_2(\mathbb{C})$. The formal codegree of the unique matrix-block representation is a rational integer. Every irreducible factor of the characteristic polynomial of C_R of degree at least two consists entirely of formal codegrees of one-dimensional representations. Consequently, if any such factor fails Ostrik's d-number divisibility condition, R has no complex fusion categorification.*

Proof. The FP dimension character is one-dimensional, so at least one scalar summand exists. Noncommutativity requires a matrix summand of dimension at least two. Since the sum of the squares of the degrees of simple modules is eight, there can be only one matrix summand and it must have degree two; the four remaining dimensions are one. The rational based algebra is split by some number field. Galois automorphisms permute its irreducible representations preserving their degrees, so the unique two-dimensional representation is Galois-fixed. Its formal codegree is an algebraic integer by [24] and rational by Galois invariance, hence belongs to $\mathbb{Z}$. Its Casimir eigenvalue is therefore rational. Every nonlinear irreducible factor of the integral Casimir polynomial

comes from the four one-dimensional representations; on each, the eigenvalue is exactly the formal codegree. Finally, formal codegrees of complex fusion categories are d-numbers [24]. For an irreducible polynomial $q(t) = t^d + a_1 t^{d-1} + \cdots + a_d$, this requires $|a_d|^j \mid a_j^d$ for every $j = 1, \ldots, d$. Failure of one exact integer divisibility gives the assertion. □

The commutative branch uses the ordinary d-number test on every Casimir factor. The script never treats raw Casimir eigenvalues as formal codegrees of noncommutative representations without Proposition 4.1. We also ran the rational-indivisibility screen of [26, Theorem 4.4], noncommuting field-automorphism witnesses from [17], and finite-field cyclotomic witnesses from [15]. These three screens add no exclusions in the present bound; their zero rows and exact code remain in the stage data. Schopieray's greedy formal-codegree obstruction $2, 3, 7$ [27] cannot occur before multiplicity five, since its forced rank-four subring has a coefficient five, so it is not an active step here.

4.3. **Integral divisibility and the complete signed quotient test.** The integral Drinfeld condition [19, Section 2.3] says that, if all simple FP dimensions of a categorifiable ring are integers, every ordinary formal codegree is a positive integer dividing its global FP dimension. For a commutative tensor, the verifier finds its unique positive FP character by an exact integral eigenvector of the strictly positive matrix $\sum_i N_i$. It then compares every exact linear factor of $\det(tI - C_R)$ with $D = \sum_i \mathrm{FPdim}(b_i)^2$. A nonintegral factor, or an integer codegree that does not divide D, is an unconditional complex no-go. This stage includes concrete multiplicity-two failures with $D = 156$ and 110, respectively, at $R8_2_288$ and $R8_2_392$.

The archived rank-eight appendix of [20] prints full tensors and explicit categorification decisions. Its items C.3.2(5) and C.3.2(7) are excluded by the paper's no-induction-matrix Lemmas 4.3 and 4.2. The appendix-to-census verifier checks all 8^3 entries under a unit-preserving permutation before transferring either published exclusion. We rely on those cited lemmas for the induction obstruction; their original Normaliz computation is not repeated here. The same transfer finds group-theoretical models in items C.3.2(3), C.3.2(6), and C.3.2(11). Item C.3.2(4) matches $R8_1_3$ but supplies no existence or exclusion claim; the explicit twisted group-theoretical construction below now realizes that ring.

Finally, the oriented signed-sphericalization theorem of [17, 18] applies even when the alleged category has no pivotal structure. The canonical sphericalization has two lifts of each simple; quotienting its Grothendieck ring by the negative central invertible yields an ordered signed multiplication tensor. We enumerate every cyclic and duality orbit of signed coefficients within the exact parity bounds, require all ordered associativity equations, then check every possible spherical character by the inherited norm, duality, and dimension-order criteria. A finite search that times out or has an unrefuted branch supplies no exclusion. The attached campaign records each attempted multiplicity-one tensor, incomplete result, and timeout separately; only certificates whose complete signed branches independently regenerate are admitted to the ledger.

4.4. **Grading obstructions with rank-one components.** The following short arguments give five further exclusions without any pivotality assumption. In a faithful grading $\mathcal{C} = \bigoplus_{g \in G} \mathcal{C}_g$, every component is an invertible bimodule category over $\mathcal{D} = \mathcal{C}_e$, and $g \mapsto [\mathcal{C}_g]$ is a homomorphism to $\mathrm{BrPic}(\mathcal{D})$ [7, Theorem 1.3 and Section 6].

Lemma 4.2. *If a pointed category* Vec_A^ω *has a module category with one simple object, then* $[\omega] = 0$.

Proof. Identify the module category with Vec. Each invertible object acts by the identity functor, up to isomorphism. The scalar module constraints form a nonzero 2-cochain whose coboundary is ω, by the module pentagon. Hence the associator class is trivial. □

Proposition 4.3. *The rings* $R8_1_26$, $R8_2_116$, *and* $R8_2_189$ *have no complex fusion categorification.*

Proof. In the original zero-based census labels, $R8_1_26$ has a faithful C_4 grading with components

$$\{0,3,4\}, \quad \{1\}, \quad \{5,6,7\}, \quad \{2\}.$$

Its neutral group is $A = C_3$. Lemma 4.2 identifies its neutral category with Vec_{C_3}. By [7, Corollary 1.2], its Brauer–Picard group is the orthogonal group of $A \oplus A^*$ with hyperbolic quadratic form. Over $\mathbb{F}_3$ this consists of $I, -I, J, -J$, where J exchanges the two coordinates. Every square is the identity, so $[\mathcal{C}_2] = [\mathcal{C}_1]^2 = 1$. On the other hand, the degree-two invertible object $s = 5$ conjugates $a = 3$ to $a^{-1} = 4$. Its component is the regular left module with right action twisted by inversion. It cannot be equivalent to the regular bimodule: such an equivalence would send s to some $b \in A$ and the relation $as = sa^{-1}$ would imply $ab = ba^{-1}$, which is false in C_3.

For $R8_2_116$ and $R8_2_189$, the universal grading has five components, of ranks $4, 1, 1, 1, 1$, so its group is C_5. Their neutral groups are respectively C_4 on $\{0, 1, 2, 7\}$ and C_2^2 on $\{0, 5, 6, 7\}$. Again their associator classes are trivial. In the first case the Brauer–Picard group embeds in $\mathrm{GL}_2(\mathbb{Z}/4)$, of order 96; in the second it is $O_4^+(\mathbb{F}_2)$, of order 72. Neither group has an element of order five. Hence the component homomorphism is trivial, forcing every component to be equivalent to the rank-four regular bimodule, contrary to the rank-one components. The verifier enumerates the small orthogonal groups and checks every grading support identity exactly. □

Proposition 4.4. *The rings $R8_2_78$ and $R8_2_168$ have no complex fusion categorification.*

Proof. Each has a faithful grading with a rank-one component and neutral fusion rules $\mathrm{Rep}(S_3)$: $g^2 = 1$, $gx = xg = x$, and $x^2 = 1 + g + x$. A rank-one module category gives a fiber functor on the neutral category $\mathcal{D}$, so $\mathcal{D} \simeq \mathrm{Rep}(H)$ for a semisimple Hopf algebra of dimension six. The dimension-six classification [10] and the rank-three fusion rules force $H \simeq \mathbb{C}[S_3]$. The dual category of its fiber-functor module is $\mathrm{Rep}(H^*) \simeq \mathrm{Vec}_{S_3}$, of rank six [22, Example 7.12.26]. But invertibility of the homogeneous $\mathcal{D}$-bimodule would identify that dual with $\mathcal{D}$, of rank three, a contradiction. □

4.5. **Solvability, equivariantization and module ranks.** We give three further general exclusions using only small groups and exact fusion rules. By [3, Theorem 1.6], a fusion category of FP dimension $p^a q^b$ is solvable. The characterization in its Definition 1.2 and Proposition 4.4 expresses it as successive cyclic-prime extensions and equivariantizations. If its adjoint subcategory is the whole category, its last nontrivial step cannot be an extension. Thus it is an actual C_p-equivariantization, with a central $\mathrm{Rep}(C_p)$ whose underlying invertibles embed into the category. We use the orbit/stabilizer description of equivariant simples [6, Section 2]; cyclic stabilizers have only one-dimensional irreducible projective representations.

Proposition 4.5. *The ring $R8_1_29$ has no complex fusion categorification.*

Proof. Its invertibles form C_6, its other two simples X, Y are self-dual of dimension three, and its adjoint ring is the whole ring. Its dimension is 24, so a putative category $\mathcal{C}$ is $\mathcal{D}^{C_p}$ with $p = 2$ or 3.

For $p = 3$, the order-three character subgroup fixes X, Y and has two free orbits on the six invertibles. The forgetful images of X, Y split into three distinct invertibles each; those of the invertibles give two more. Thus $\mathcal{D} = \mathrm{Vec}_A^\omega$, $|A| = 8$, with two free C_3-orbits. The only groups of order eight admitting an order-three automorphism are C_2^3 and Q_8. On C_2^3 the action has a fixed line and an irreducible plane. The three nonzero elements in that plane form a free orbit whose product has coefficient two into the same free orbit. The output stabilizer is trivial, so associator and tensorator choices cannot change this coefficient. On Q_8, inversion exchanges the orbits $\{i, j, k\}$ and $\{-i, -j, -k\}$; their two induced dimension-three objects are distinct duals. These conclusions contradict multiplicity one and self-duality, respectively.

For $p = 2$, the character involution exchanges X, Y. De-equivariantization has the fusion rules

$$G(\mathcal{D}) = C_3 = \langle a \rangle, \qquad a\rho = \rho a = \rho, \qquad \rho^2 = 1 + a + a^2 + 2\rho.$$

It has dimension twelve and trivial universal grading. The same last-step argument gives $\mathcal{D} = (\mathrm{Vec}_V^\omega)^{C_3}$, where $V = C_2^2$ and C_3 permutes the three nonzero elements transitively. No assumption

on ω is made. A monoidal automorphism of $\mathrm{Id}_{\mathcal{D}}$ has scalar one on ρ, because ρ occurs in ρ^2, and then on all invertibles. Consequently the map from central invertibles to underlying invertibles is injective, and the central $\mathrm{Rep}(C_3)$ is unique.

The C_2 action on $\mathcal{D}$ fixes every simple: exchanging a, a^2 would produce a dimension-two simple upstairs. Every such tensor autoequivalence preserves the unique central $\mathrm{Rep}(C_3)$. Its restriction is tensor-isomorphic to the identity, since $H^2(C_3, \mathbb{C}^\times) = 0$. Equivariantization/de-equivariantization [4, Theorem 4.18(i) and Proposition 4.19] therefore lifts it to a C_3-equivariant tensor autoequivalence of Vec_V^ω. Its object permutation centralizes C_3 in $\mathrm{Aut}(V) = S_3$, so lies in C_3. Composing with the inverse action functor removes that permutation without changing the descended equivalence class.

Identity-on-objects tensor autoequivalences of Vec_V^ω are classified by $H^2(V, \mathbb{C}^\times) = C_2$: the two associators cancel in the tensorator equation. Their equivariant structures have ambiguity $H^1(C_3, \widehat{V}) = 0$, since multiplication by three is invertible on $\widehat{V}$. Both possible classes have explicit representatives. Besides the identity, take

$$J(u, v) = (-1)^{u_1 v_1 + u_1 v_2 + u_2 v_2}, \qquad u, v \in \mathbb{F}_2^2.$$

This nontrivial normalized 2-cocycle is C_3-invariant and satisfies $J^2 = 1$. Thus it commutes even with nontrivial scalar tensorators of the existing C_3 action and descends with identity comparisons. There are no further C_2 coherence choices, because $\mathrm{Aut}_\otimes(\mathrm{Id}_{\mathcal{D}}) = 1$.

The identity action leaves the coefficient two in ρ^2. For the J action, all degree-zero channels have $J(u, u) = -1$ for $u \neq 0$; for each nonzero output degree, the two ordered channels have opposite signs. Hence each lift $\rho_\pm$ has square containing the three negative-character invertibles and one copy of each $\rho_\pm$. The unit occurs in $\rho_+\rho_-$, not in either square. The two lifts are distinct duals, again contradicting the self-duality in $R8_1_29$. This exhausts both prime branches for arbitrary pointed associators. □

Lemma 4.6 (Module-rank divisibility)**.** *For a fusion category $\mathcal{B}$, let $g_{\mathcal{B}}$ be the greatest common divisor of the ranks of its indecomposable finite semisimple module categories. If $\mathcal{B} \subseteq \mathcal{D}$ is a fusion subcategory, then $g_{\mathcal{B}} \mid \mathrm{rank}(\mathcal{D})$.*

Proof. Decompose $\mathcal{D}$, with its left tensor action by $\mathcal{B}$, into indecomposable module categories. Their ranks add to $\mathrm{rank}(\mathcal{D})$. This statement uses ranks in the specified module presentation; it makes no assertion that module ranks are Morita invariant. □

Proposition 4.7. *The rings $R8_1_45$ and $R8_1_47$ have no complex fusion categorification.*

Proof. Each ring has simple dimensions $(1, 1, 2, 3, 3, 4, 4, 4)$ and admits labels $1, h, E, P, Q, W_0, W_1, W_2$ with

$$h^2 = 1, \quad hE = E, \quad hP = Q, \quad hW_i = W_i, \quad P^* = Q,$$
$$E^2 = 1 + h + E, \qquad EP = EQ = P + Q, \qquad PQ = 1 + E + P + Q.$$

The adjoint ring is the whole ring. Dimension 72 and the invertible group C_2 therefore force a putative categorification to be $\mathcal{C} = \mathcal{D}^{C_2}$ with $\mathrm{FPdim}(\mathcal{D}) = 36$. The forgetful images are

$$F(1) = F(h) = 1, \quad F(E) = a + a^2, \quad F(P) = F(Q) = Y, \quad F(W_i) = U_i + V_i.$$

Here a has order three, Y is simple of dimension three, and the six U_i, V_i are distinct simples of dimension two. Thus $\mathrm{rank}(\mathcal{D}) = 10$. The full fusion subcategory $\mathcal{B}$ on $1, a, a^2, Y$ satisfies

$$aY = Ya = Y, \qquad Y^* = Y, \qquad Y^2 = 1 + a + a^2 + 2Y.$$

The C_2 action exchanges a, a^2 and fixes Y; its equivariantization inside $\mathcal{C}$ has the displayed nonselfdual P, Q.

As in the preceding proof, solvability and trivial universal grading imply $\mathcal{B} = (\mathrm{Vec}_V^{\omega})^{C_3}$, $V = C_2^2$, with the transitive action. The restriction map

$$H^3(V, \mathbb{C}^\times) \longrightarrow \prod_{\substack{L \le V \\ |L|=2}} H^3(L, \mathbb{C}^\times)$$

is an isomorphism $C_2^3 \to C_2^3$. In integral-cohomology generators u^2, v^2, uv, the three restrictions of (r, s, t) are $(r, s, r + s + t)$. Since C_3 permutes the factors, there are exactly two invariant classes: zero and the class nontrivial on every subgroup of order two.

Suppose first $[\omega] = 0$. The lift of the object C_3 action is unique in $\mathrm{Aut}_\otimes(\mathrm{Vec}_V) = C_2 \times S_3$, and its coherence is unique because $H^2(C_3, \widehat{V}) = 0$. Therefore $\mathcal{B} \simeq \mathrm{Rep}(A_4)$. Its central $\mathrm{Rep}(C_3)$ is unique, since $\mathrm{Aut}_\otimes(\mathrm{Id}_\mathcal{B}) = 1$, and is preserved by the C_2 action. Sequential equivariantization and de-equivariantization [3, Section 2.6 and Proposition 2.10][4, Proposition 4.19] identify $\mathcal{B}^{C_2}$ with $(\mathrm{Vec}_V)^{S_3}$. The action on nonzero elements is the usual faithful S_3 permutation action; inversion on the character C_3 forces the group $C_3 \rtimes C_2 = S_3$.

There are exactly two lifts of this object action to $C_2 \times S_3$, given by the trivial or parity homomorphism into $H^2(V, \mathbb{C}^\times) = C_2$. Their coherence ambiguity vanishes: $H^j(S_3, \widehat{V}) = 0$ for $j > 0$, by the normal C_3 subgroup, whose positive cohomology and fixed space on $\widehat{V}$ both vanish. Both lifts yield self-dual dimension-three orbit objects. For completeness choose a stabilizing transposition τ swapping coordinates and its fixed nonzero element $v = (1, 1)$. For the trivial lift, the equivariance scalars $u = \pm 1$ act trivially on the unit channel of $v \otimes v$. For the other lift choose tensorator $\mu(a, b) = (-1)^{a_1 b_2}$. Its square is trivialized by $\eta_a = (-1)^{a_1 a_2}$, so $\eta_v = \mu(v, v) = -1$. The equivariance scalar satisfies $u^2 = \eta_v$ and acts on the unit channel by $u^2/\mu(v, v) = 1$. A τ-invariant character change of η is one on v, so cannot change the conclusion. Frobenius reciprocity for the orbit induction gives a tensor unit in each of the two squares. This contradicts $P^* = Q$.

It remains that $[\omega]$ is nontrivial on every order-two subgroup. The crossed product of Vec_V^{ω} by C_3 is pointed, and realizes $\mathcal{B}$ as the group-theoretical category

$$\mathcal{B} \simeq \mathcal{C}(A_4, \Omega, K, \psi), \qquad K = C_3, \qquad [\Omega]|_V = [\omega], \quad [\Omega]|_K = 0.$$

By the classification of pointed module categories and its double-coset rank formula [5, Example 2.1, Theorem 3.1 and Proposition 3.1], indecomposable $\mathcal{B}$-module categories correspond to subgroups $H \le A_4$ with $[\Omega]|_H = 0$ and permissible cochains. If $|H|$ were even, Cauchy's theorem would give a subgroup of order two, contradicting the restriction of Ω. Thus H is trivial or has order three. For $H = 1$, the four double cosets in $K\backslash A_4/H$ have trivial intersections, giving module rank four. For $H = C_3$, conjugate it to K. There are two double cosets, with intersections C_3 and 1; projective cocycles on C_3 are coboundaries, so their contributions are three and one. Every indecomposable $\mathcal{B}$-module category therefore has rank four, regardless of the permissible cochains. Lemma 4.6 would force $4 \mid \mathrm{rank}(\mathcal{D}) = 10$, a contradiction. □

The portable verifiers check the source-bound tensors, the displayed fusion identities, the complete small cocycle identities, and all relevant A_4 subgroups and double cosets. These finite checks certify the hypotheses; the categorical exhaustion is supplied by the proofs above. Neither argument assumes a braiding or a pivotal structure.

4.6. **Rank-one modules and semisimple Hopf algebras.**

Proposition 4.8. *The rings* R8_2_41*,* R8_2_86*,* R8_2_162*,* R8_2_140 *and* R8_2_784 *have no complex fusion categorification.*

Proof. For the first three rings, a faithful C_2 grading has a rank-one nonneutral component and a dimension-sixteen neutral category $\mathcal{D}$ of type $(1^4, 2^3)$. A rank-one module supplies a fiber functor, so $\mathcal{D} = \mathrm{Rep}(H)$ for a semisimple Hopf algebra of dimension sixteen. Invertibility of the component further requires $\mathrm{Rep}(H^*) \simeq \mathcal{D}$; in particular both groups of grouplikes have order four. Kashina's

classification [44, Theorem 1.2 and Table 1] leaves only $H_{C:\sigma_1}$, whose representation ring is $\mathrm{Rep}(D_{16})$: its invertibles form C_2^2, and all three dimension-two simples are self-dual. The rings $R8_2_41$, $R8_2_86$ instead have cyclic invertibles, while $R8_2_162$ has a nonselfdual dimension-two pair.

For $R8_2_140$, census indices $\{0,1,2,3,7\}$ give the neutral Tambara–Yamagami category on C_4, of dimension eight. Each other simple separately gives a rank-one left module: an invertible acts as one and the dimension-two object acts as two. Thus the neutral category has a fiber functor and is $\mathrm{Rep}(H)$ with $G(H^*) = C_4$. Kashina's prime-power theorem [44, Theorem 1.1] says that a semisimple Hopf algebra of prime-power dimension with cyclic grouplikes is a group algebra. It would give $H^* \simeq \mathbb{C}[C_8]$, which has eight grouplikes, a contradiction. This argument does not require the rank-one modules to be grading components.

For $R8_2_784$, the neutral component has dimension twenty-two, rank seven and type $(1^2, 2^5)$; the other component has a unique simple of dimension $\sqrt{22}$. Its fiber functor reconstructs a Hopf algebra of dimension $2 \cdot 11$. The Hopf pq classification [10] and the nonpointed neutral rules force $\mathcal{D} = \mathrm{Rep}(D_{22})$. Its fiber-functor dual is $\mathrm{Vec}_{D_{22}}$, of rank twenty-two. The invertible grading component would identify this dual with $\mathcal{D}$, of rank seven. This is impossible. □

4.7. **Preserved Tannakian subcategories.**

Proposition 4.9. *The rings $R8_2_697$, $R8_2_808$ and $R8_2_181$ have no complex fusion categorification.*

Proof. For $R8_2_697$, dimension sixty-eight, type $(1^4, 4^4)$ and trivial universal grading, with all four invertibles fixing each noninvertible, force a C_2 equivariantization with descent $\mathcal{D}$ of dimension thirty-four and type $(1^2, 2^8)$. Its adjoint is the whole category: a proper adjoint has dimension $1, 2$ or 17; a prime-dimensional adjoint is pointed, but these possibilities cannot contain the required dimension-four squares with only two invertibles. Solvability and prime descent now give $\mathcal{D} = (\mathrm{Vec}^{\omega}_{C_{17}})^{C_2}$, with inversion. Since $U(\mathcal{D}) = 1$, central lifts of its invertibles are unique. The outer involution therefore preserves its central $\mathrm{Rep}(C_2)$, and the combined equivariantization group is C_2^2, as its four linear characters are the census invertibles. Every $C_2^2 \to \mathrm{Aut}(C_{17})$ has image of order at most two. A nonzero orbit has size two and cyclic stabilizer; it therefore supplies dimension-two equivariant simples. The target has only dimension-four noninvertibles.

For $R8_2_808$, dimension $186 = 2 \cdot 3 \cdot 31$ and the integral pqr theorem make any categorification group-theoretical [3, Theorem 9.2]. Groups of order 186 are solvable, so the category is solvable. The adjoint, type $(1^2, 2, 6^5)$, and the character involution fixing all noninvertibles force C_2 descent of type $(1^3, 3^{10})$ and dimension ninety-three. The same prime-dimensional-adjoint argument shows that its universal grading is trivial. Thus it is $(\mathrm{Vec}^{\omega}_{C_{31}})^{C_3}$ with nontrivial order-three action. Uniqueness of the central C_3 preserves it under the outer involution. The combined group is S_3, because its embedded representation subcategory has type $(1^2, 2)$. Every homomorphism $S_3 \to \mathrm{Aut}(C_{31})$ kills its normal C_3, contradicting that action.

For the commutative ring $R8_2_181$, the neutral category has dimension twenty-one, type $(1^3, 3^2)$ and trivial universal grading. Solvability makes it $(\mathrm{Vec}^{\omega}_{C_7})^{C_3}$, with multiplication by two or four on C_7. The action on $H^3(C_7, \mathbb{C}^\times) = C_7$ is multiplication by four or two, so its invariant associator class is zero. Coprime action cohomology then identifies the neutral category with $\mathrm{Rep}(C_7 \rtimes C_3)$. Its central C_3 has unique lifts. The three odd simples have free, identical left and right C_3 actions, so the odd bimodule fixes these central lifts pointwise. Relative-center equivariantization [47] and $H^2(C_2, \mathbb{C}^\times) = 0$ give six central invertible lifts upstairs. The unique order-three subgroup has trivial self-braiding, inherited from the neutral $\mathrm{Rep}(C_3)$, and is an embedded Tannakian subcategory. Its de-equivariantization has the Tambara–Yamagami rules on C_7. The induced C_3 symmetry must preserve its nondegenerate symmetric bicharacter $\chi(a, b) = \exp(2\pi i k a b/7)$, $k \neq 0$. Multiplication by two or four cannot do so: neither square is one modulo seven. This proves the contradiction without a pivotal or a fermionic-transparency assumption. □

4.8. **A dimension-twenty-eight adjoint criterion.**

Lemma 4.10. *A nonpointed integral fusion category of dimension twenty-eight has universal grading C_2 and adjoint fusion ring* $\mathrm{Rep}(D_{14})$.

Proof. Jordan–Larson's theorem [45, Theorem 1.1] applies with $p = 7, q = 2$ and $7 \nmid q + 1$. Write the category as $\mathcal{C}(G, \omega, H, \psi)$, where $|G| = 28$. Sylow theory gives $G = C_7 \rtimes P$, with $P = C_4$ or C_2^2; the action is trivial or inversion. The possible subgroup orders are $1, 2, 4, 7, 14, 28$. For orders one and seven, or a central subgroup of order two, the cyclic stabilizers give a pointed category. For a noncentral subgroup H of order two, its normal closure is $C_7 \rtimes H = D_{14}$, of index two; support gives a faithful C_2 grading. For $|H| = 14$, H is normal and support again gives such a grading. An abelian $H = C_{14}$ would give a pointed category; otherwise $H = D_{14}$, whose Schur multiplier is trivial. For $H = G$ one obtains $\mathrm{Rep}(G)$; a nonabelian group of order twenty-eight has center C_2 and quotient D_{14}. In each of these nonpointed cases the neutral fusion ring is $\mathrm{Rep}(D_{14})$.

It remains to justify this ring identification for $|H| = 4$, and also the cocycle reduction used for a reflection complement above. Conjugate H to P. The positive-degree cohomology of C_7 is seven-primary, so the coprime Lyndon–Hochschild–Serre sequence has no mixed positive-degree terms. Since $\omega|_P = d\psi$, the associator class is represented by a P-invariant class on C_7. Choose a normalized μ_7 representative, average over P, and raise to the inverse of $|P|$ modulo seven. This gives a strictly invariant representative α. On $C_7 \rtimes P$ use

$$\omega_\alpha((n_1, p_1), (n_2, p_2), (n_3, p_3)) = \alpha(n_1, p_1 n_2, p_1 p_2 n_3).$$

It is identically one on P. The remaining 2-cocycle ψ is removed by its inflation along $G \to P$. Thus the category is $(\mathrm{Vec}_{C_7}^{\alpha})^P$ with strict permutation action. A trivial action is pointed. Otherwise its kernel C_2 acts strictly trivially and its characters grade the category, with neutral $(\mathrm{Vec}_{C_7}^{\alpha})^{C_2}$. This has the $\mathrm{Rep}(D_{14})$ fusion ring; the same argument with a reflection complement supplies the previous order-two case.

Finally this neutral ring is adjoint to itself. For its three dimension-two objects, indexed by nonzero elements modulo inversion, $X_i^2 = 1 + \varepsilon + X_{2i}$, and doubling cycles the three indices. The adjoint of the full category is contained in the displayed neutral component and contains its adjoint, so it equals that component. Consequently the universal grading is exactly C_2. □

Proposition 4.11. *There is no fusion category with dimension fifty-six, type* $(1^4, 2, 4^3)$ *and trivial universal grading. In particular R*8_2_694 *is generally excluded.*

Proof. Solvability and trivial universal grading force a last equivariantization. A C_7 equivariantization would supply seven invertibles, so the last group is C_2. Its descent $\mathcal{D}$ has dimension twenty-eight and is integral by the orbit/stabilizer dimension formula. It cannot be pointed, since a C_2 equivariantization of a pointed category has all dimensions at most two. Lemma 4.10 therefore applies. The outer involution preserves $\mathcal{D}_{\mathrm{ad}}$ and permutes its three dimension-two simples, fixing at least one. Its cyclic stabilizer has trivial projective multiplier, so its two characters produce two distinct dimension-two equivariant simples. The proposed category has only one. This is a contradiction. □

4.9. **A two-object orbit obstruction in dimension ninety-six.**

Proposition 4.12. *A fusion category of type* $(1^4, 2, 4, 6^2)$ *has a nontrivial universal grading. In particular R*8_2_404*, R*8_2_418*, R*8_2_604*, R*8_2_692 *have no complex fusion categorification.*

Proof. Suppose such a category has trivial universal grading. Its dimension is ninety-six, and its invertible group has order four. Solvability forces a last C_2 equivariantization $\mathcal{C} = \mathcal{D}^{C_2}$. Whichever nontrivial character occurs, it fixes the unique dimension-two and dimension-four simples. Thus the outer involution exchanges the two dimension-two simples of $\mathcal{D}$. Depending on whether it fixes or exchanges the dimension-six pair upstairs, $\mathcal{D}$ has type $(1^4, 2^2, 3^4)$ or $(1^4, 2^2, 6)$ and dimension forty-eight.

Its canonical adjoint is preserved by the outer involution. A proper adjoint's dimension therefore has the form $a+4b+9c+36e$, where $a \in \{1,2,4\}$, $b \in \{0,2\}$, c is even, and e is zero or one according to the displayed type. The proper divisors of forty-eight among these possibilities are only $1, 2, 4, 12$. A pointed adjoint of dimension at most four cannot contain the square of a dimension-three or dimension-six simple, since each invertible occurs there with coefficient at most one. An adjoint of dimension twelve consists of all four invertibles and both dimension-two simples. Every other grading component would then have dimension twelve using only dimension-three or dimension-six simples; a sum of their squared dimensions is divisible by nine, so this is impossible. Hence $\mathcal{D}_{\mathrm{ad}} = \mathcal{D}$.

A second prime descent gives $\mathcal{D} = \mathcal{E}^{C_2}$, $\mathrm{FPdim}(\mathcal{E}) = 24$. The complete cyclic-orbit dimension partitions are $(1^6, 3^2)$ and $(1^2, 2, 3^2)$. In the first case the dimension-two pair of $\mathcal{D}$ comes from two free orbits in the invertible group A of order six. The inner involution fixes exactly two elements of A. For $A = C_6$ it is inversion, and for $A = S_3$ it is conjugation by a transposition. In either group the coefficients of the two induced objects in their own squares are respectively zero and one, by ordinary orbit addition. The output stabilizers are trivial, so associator or tensorator phases cannot change these coefficients. The outer involution cannot exchange this pair as required.

In the second case the unique dimension-two simple x of $\mathcal{E}$ is fixed by the inner involution, with $x^2 = 1 + g + x$. Its two equivariant lifts x_+, x_- are exchanged by the central character ε. Therefore their squares are equal, because $\varepsilon^2 = 1$ and ε is central. Each square has precisely one constituent from the dimension-two pair, by forgetting to the single x channel. The outer involution exchanges x_+, x_- but preserves their common square. Its unique dimension-two constituent would have to be fixed, which is impossible. Both dimension partitions and every initial character choice have therefore been excluded. The supplementary check binds the full four source tensors and exhausts these small dimension and orbit cases. □

4.10. **An order-four action on rank-one components.**

Proposition 4.13. *The ring R8_2_68 has no complex fusion categorification.*

Proof. Its exact type is $(1^4, 4^2, 6^2)$, its invertibles form C_4, its adjoint is the whole ring, and its order-two invertible fixes all four noninvertibles. Solvability in dimension 108 therefore forces $\mathcal{C} = \mathcal{D}^{C_2}$ with $\mathcal{D}$ integral of dimension 54 and type $(1^2, 2^4, 3^4)$. The outer involution exchanges both noninvertible dimension classes in pairs and fixes the invertibles. The possible proper dimensions of its invariant adjoint are $1, 2, 9, 18, 27$. Pointed dimensions one and two cannot contain the square of a dimension-three object; dimensions nine and twenty-seven would contain dimension-two objects, contradicting strong Frobenius divisibility [3]. Thus either $\mathcal{D}_{\mathrm{ad}} = \mathcal{D}$ or its adjoint $\mathcal{E}$ has type $(1^2, 2^4)$ and dimension eighteen.

In the first case prime descent gives $\mathcal{D} = \mathcal{F}^{C_2}$, with $\mathcal{F}$ of type $(1^9, 3^2)$. In the second case $U(\mathcal{E}) = 1$: a grading of order two would have a dimension-nine neutral containing a dimension-two simple, again contradicting strong Frobenius; order-three components of dimension six would each require both invertibles; higher grading orders cannot accommodate a dimension-two simple. Hence $\mathcal{E} = (\mathrm{Vec}_A^{\omega})^{C_2}$, $|A| = 9$, with inversion on A. Its unique nontrivial central invertible g is the canonical Tannakian lift. The relative-center C_3 action of the grading of $\mathcal{D}$ fixes this lift. Since $H^2(C_3, \mathbb{C}^{\times}) = 0$, its central equivariant lifts, together with the unit lifts, form an abelian group of order six. The unique order-two lift restricts to g and inherits its self-braiding $+1$; it therefore gives a central $\mathrm{Rep}(C_2)$ in $\mathcal{D}$. This explicitly verifies bosonicity, rather than inferring it from centrality alone. De-equivariantization again gives $\mathcal{F}$ of type $(1^9, 3^2)$: a dimension-three object fixed by g would descend to dimension $3/2$, which is not an algebraic integer.

In both cases the order-two central lift in $\mathcal{D}$ is unique: its kernel is trivial when $U(\mathcal{D}) = 1$, and has order three when $U(\mathcal{D}) = C_3$. The outer involution preserves this Tannakian subcategory. Combining the two equivariantizations gives $\mathcal{C} = \mathcal{F}^{C_4}$, because the intermediate symmetric subcategory has dimension four and its four characters form the census C_4. Let β generate this coherent action. Its square acts by inversion on $A = \mathrm{Inv}(\mathcal{F})$ and fixes the two dimension-three objects.

The adjoint of $\mathcal{F}$ is Vec_A^ω, with a C_3 grading of ranks $9, 1, 1$. Rank-one components force $[\omega] = 0$ by their module pentagon. The action of β exchanges these two components; otherwise the original $\mathcal{C}$ would inherit a nontrivial C_3 grading. For $A = C_9$ no automorphism squares to inversion. For $A = C_3^2$, its matrix B satisfies $B^2 = -I$, so its minimal polynomial is the irreducible $t^2 + 1$ over $\mathbb{F}_3$ and $\det(B) = 1$. Its tensorator class $\eta \in H^2(A, \mathbb{C}^\times) = C_3$ is fixed by B; order-four coherence gives $4\eta = 0$, hence $\eta = 0$. Such a tensor autoequivalence fixes every rank-one left-module class, since those classes are $H^2(A, \mathbb{C}^\times)$. But the two nonneutral components have different left-module classes: an equivalence, tensored with the inverse of one invertible bimodule, would identify the rank-nine regular module with a rank-one component. Thus β cannot exchange them, a contradiction. The supplement checks every source coefficient, adjoint dimension partition and matrix calculation; the central lift and action tower use [47, 4, 3]. □

4.11. **Integral extensions from pointed Morita data.**

Lemma 4.14. *Suppose $\mathcal{B}$ is Morita equivalent to Vec_G^ω and every normal abelian subgroup of G has square order. Then every faithful finite-group graded extension of $\mathcal{B}$ is integral.*

Proof. An invertible Vec_G^ω bimodule is, as a left module, a transitive pointed module associated to a subgroup $N \leq G$ and a cochain trivializing $\omega|_N$. Its dual is pointed [7, Proposition 4.2]. In the twisted double-coset description [46, Theorems 4.1 and 5.1], a simple over NaN has FP dimension

$$[N : N \cap aNa^{-1}] \dim(\pi),$$

where π is an irreducible projective representation of the intersection. Pointedness forces N to be normal. The identity coset subcategory is ordinary $\mathrm{Rep}(N)$, even with the general associator and subgroup cochain [46, Theorem 4.1]; thus N is abelian as well. The left module has $[G : N]$ simples with equal normalized FP dimension. Since a graded component has total squared dimension $|G|$, each dimension is $\sqrt{|N|}$, an integer. Every component of an extension of Vec_G^ω is an invertible bimodule, so that extension is integral.

Morita equivalence transports the complete graded-extension data, including multiplication and associativity, and the resulting complete extensions are Morita equivalent [7, Corollary 3.13 and Theorem 1.3]. Integrality is preserved under this equivalence in the present setting. Indeed, the center of an integral category is integral because forgetting preserves FP dimensions. A Morita equivalent category $\mathcal{C}$ has the same center and integer total dimension. For a simple $W \in \mathcal{C}$, induction has dimension $\mathrm{FPdim}(\mathcal{C})\,\mathrm{FPdim}(W)$, an integer. Hence $\mathrm{FPdim}(W)$ is rational and an algebraic integer, and therefore is an integer. □

Proposition 4.15. *The rings R8_2_53 and R8_2_93 have no categorification over $\mathbb{C}$.*

Proof. Each source tensor has FP type $1^4, 4^2, (3\sqrt{2})^2$ and a faithful C_2 grading with integral neutral subring D of dimension 36. Its invertibles form $C_4 = \langle g \rangle$. They fix both noninvertibles X, Y, which are self-dual, and

$$X^2 = 1 + g + g^2 + g^3 + X + 2Y, \qquad Y^2 = 1 + g + g^2 + g^3 + 2X + Y.$$

In particular $D_{\mathrm{ad}} = D$. Any categorification $\mathcal{D}$ is solvable by the two-prime theorem [3, Theorem 1.6]. Its last prime step must be an equivariantization since its universal grading is trivial. The prime cannot be three because $\mathrm{Inv}(\mathcal{D}) = C_4$ contains no C_3. Thus $\mathcal{D} = \mathcal{E}^{C_2}$, with $\mathrm{FPdim}(\mathcal{E}) = 18$. The sign object is g^2 and fixes X, Y. The equivariantization orbit formula [6] makes $\mathcal{E}$ of type $1^2, 2^4$.

Let η be its nontrivial invertible. In every ZZ^* with $\mathrm{FPdim}(Z) = 2$, omitting η would leave dimension three after the unit, which cannot be a sum of two-dimensional simples. Therefore $ZZ^* = 1 + \eta + T$ with $\mathrm{FPdim}(T) = 2$. The adjoint subcategory contains both units and a two-dimensional simple. If proper, its integer dimension divides 18; the only possibilities for a subcategory of type $1^2, 2^n$ are 2 and 6. Dimension two omits T. At dimension six, the other universal-grading

components contain only two-dimensional simples and cannot have total squared dimension six. Hence $\mathcal{E}_{\mathrm{ad}} = \mathcal{E}$. Solvability again forces $\mathcal{E} = \mathcal{P}^{C_2}$, where $\mathrm{FPdim}(\mathcal{P}) = 9$ and $\mathcal{P}$ is pointed [21, 22].

The canonical central $\mathrm{Rep}(C_2) \subset Z(\mathcal{E})$ has underlying object η. Its half-braiding is unique: two half-braidings on an invertible differ by a tensor character, and $\mathrm{Aut}_{\otimes}(\mathrm{Id}_{\mathcal{E}}) = \mathrm{Hom}(U(\mathcal{E}), \mathbb{C}^{\times}) = 1$. The outer C_2 action therefore preserves this central subcategory. Its equivariantization embeds as a symmetric subcategory of $Z(\mathcal{E}^{C_2})$: equivariant half-braidings act on every equivariant object, and the morphism conditions give full faithfulness. Forgetting equivariance followed by the fiber functor on $\mathrm{Rep}(C_2)$ gives a fiber functor, so this subcategory is $\mathrm{Rep}(H)$ for a group of order four. Its four underlying invertibles exhaust the C_4 in $\mathcal{D}$, forcing $H = C_4$. Central de-equivariantization [4, 22] now gives

$$\mathcal{D} \simeq (\mathrm{Vec}_A^{\alpha})^{C_4}, \qquad |A| = 9.$$

The dimension-four noninvertibles force C_4 to act freely on all nonidentity elements of A. Thus $A = C_3^2$, since $\mathrm{Aut}(C_9)$ has order six. Its generator acts by an irreducible matrix J over $\mathbb{F}_3$ with $J^2 = -I$.

The pointed crossed product is Morita dual to this equivariantization. Consequently $\mathcal{D}$ is Morita equivalent to Vec_G^{ω}, where $G = C_3^2 \rtimes_J C_4$; the group extension splits by coprimality. The only normal abelian subgroups of G are 1 and C_3^2. Indeed, an element with nonzero C_4 exponent has commutators with C_3^2 filling $(J^k - I)C_3^2 = C_3^2$. A normal subgroup containing such an element contains C_3^2 and cannot be abelian. Within C_3^2, irreducibility of J leaves only 1 and C_3^2. Their orders are squares, so Lemma 4.14 makes every extension of $\mathcal{D}$ integral. The alleged census category would be a C_2 extension with two simples of dimension $3\sqrt{2}$, a contradiction. The exact verifier checks both full source tensors, the grading and every finite-group premise. □

Proposition 4.16. *The rings* R8_2_191 *and* R8_2_299 *have no categorification over* $\mathbb{C}$.

Proof. Their source tensors have FP type $1^4, 2^2, (\sqrt{6})^2$ and a faithful C_2 grading with integral neutral subring D of dimension 12. Its invertibles form V_4 and fix two simples X, Y, with

$$X^* = Y, \qquad X^2 = 2Y, \qquad Y^2 = 2X, \qquad XY = YX = \sum_{g \in V_4} g.$$

Thus a hypothetical neutral category $\mathcal{D}$ is faithfully C_3 graded over its pointed subcategory $\mathcal{B}$ on V_4, with rank-one nonneutral components. A rank-one module trivializes the pointed associator, so $\mathcal{B} \simeq \mathrm{Vec}_{V_4}$.

The ambient odd component $\mathcal{M}$ has two simples permuted transitively by V_4, with a stabilizer $H = C_2$; its stabilizer in source labels is $(0, 5)$ for R8_2_191 and $(0, 7)$ for R8_2_299. Its restriction $\mathcal{N}$ to $\mathcal{B}$ is therefore an indecomposable rank-two module. Since $H^2(C_2, \mathbb{C}^{\times}) = 0$, its cochain is trivial. A decomposition $V_4 = H \times K$ identifies its dual with

$$\mathcal{B}_{\mathcal{N}}^* \simeq \mathrm{Rep}(H) \boxtimes \mathrm{Vec}_K \simeq \mathrm{Vec}_{V_4},$$

including the trivial associator. The actual multiplication action of $\mathcal{D}$ on $\mathcal{M}$ is a coherent extension of this $\mathcal{B}$ module.

Apply [58, Theorem 1.3] to the *induced* module $\mathcal{L} = \mathcal{D} \boxtimes_{\mathcal{B}} \mathcal{N}$. It makes the dual $\mathcal{D}_{\mathcal{L}}^*$ a semidirect C_3 product with neutral component $\mathcal{B}_{\mathcal{N}}^*$, and hence pointed. The theorem concerns this induced-module dual; it does not assert that the dual of the original two-object $\mathcal{M}$ is pointed.

The original C_3 Brauer–Picard action on $\mathcal{B}$ is nontrivial: a rank-one component is not the regular rank-four module. Its class therefore has order three. Morita transport conjugates the action and preserves this order. The tensor-automorphism classes of Vec_{V_4} have kernel $H^2(V_4, \mathbb{C}^{\times}) = C_2$ over $\mathrm{Aut}(V_4) = S_3$ [22]. An order-three class must therefore act nontrivially on V_4. The pointed Morita group is $V_4 \rtimes C_3 = A_4$, so $\mathcal{D}$ is Morita equivalent to $\mathrm{Vec}_{A_4}^{\omega}$ for some ω. The only normal abelian subgroups of A_4 are 1 and V_4, of square orders 1 and 4. Lemma 4.14 makes every extension of this compatible neutral category integral, contradicting the odd FP dimensions $\sqrt{6}$. The source checker

verifies both gradings, the complete neutral tensors, the odd module action and all ten subgroups of A_4. The neutral ring by itself is not claimed to have only pointed Morita models; the coherent odd module is essential. □

Proposition 4.17. *The rings* R8_2_158*,* R8_2_186*,* R8_2_579 *and* R8_2_682 *have no categorification over* $\mathbb{C}$*.*

Proof. All four source tensors have FP type $1^4, 2, 4, (2\sqrt{3})^2$ and a faithful C_2 grading. Their common integral neutral subring D, of dimension 24, has V_4 invertibles fixing Y, Z and

$$Y^* = Y, \quad Z^* = Z, \qquad Y^2 = \sum_{g\in V_4} g, \quad YZ = ZY = 2Z, \quad Z^2 = \sum_{g\in V_4} g + 2Y + 2Z.$$

Thus $D_{\mathrm{ad}} = D$. Solvability [3, Theorem 1.6] forces a neutral categorification to be $\mathcal{D} = \mathcal{E}^{C_2}$ with $\mathrm{FPdim}(\mathcal{E}) = 12$; its invertibles exclude a final C_3 equivariantization. The sign ϵ fixes Y, Z. The orbit formula makes $\mathcal{E}$ of type $1^4, 2^2$, with an outer involution swapping the two noninvertibles and swapping a pair of invertibles.

The finite rank-six check exhausts both groups $A = C_4, V_4$, their trivial and transitive actions on the two nonunits, both dualities, and every oriented nonunit product allowed by FP dimension and reciprocity. There are 100 candidate tables and six associative ones, of which exactly two admit the required outer involution. Their rings are $R_{3,A}$:

$$X^* = X', \qquad X^2 = 2X', \qquad (X')^2 = 2X, \qquad XX' = X'X = \sum_{g\in A} g,$$

with all units fixing X, X'. The transitive C_4 action in this exhaustion is the homomorphism $(0, 1, 0, 1)$ in cyclic labels. Jordan–Larson [45, Theorem 1.4] excludes $A = C_4$, since its cyclic prime-power factor occurs with odd multiplicity. Thus $\mathcal{E}$ is C_3 graded over $\mathcal{B} \simeq \mathrm{Vec}_{V_4}$; its rank-one components trivialize the pointed associator.

Let $\mathcal{M}$ be the two-object ambient odd module over $\mathcal{D}$. The central regular algebra $1 + \epsilon$ in $\mathcal{D}$ gives an actual de-equivariantized module $\mathcal{M}_{1+\epsilon}$ over $\mathcal{E}$. This construction uses the central algebra inside $\mathcal{D}$; it requires no half-braiding on ϵ in the ambient category. Its rank is one if ϵ swaps the source simples, and four if it fixes them: each fixed simple has two algebra-module structures, using $H^2(C_2, \mathbb{C}^\times) = 0$.

The six indecomposable Vec_{V_4} modules have ranks $4, 1, 1, 2, 2, 2$, by their stabilizers and cochains. The C_3 action sends the regular rank-four class to the X component, then to the X' component. These are distinct rank-one classes, so they form a three-cycle of total rank six. The three rank-two classes are either fixed or form another orbit of total rank six. Galindo's Clifford theorem [58, Proposition 4.1 and Theorem 1.1] now excludes rank one for $\mathcal{M}_{1+\epsilon}$, and makes its rank-four alternative two fixed rank-two submodules. The group C_3 cannot permute two actual summands; each is a coherent $\mathcal{E}$-module extension of a rank-two $\mathcal{B}$ module.

For one such module, the induced-module dual of [58, Theorem 1.3] is a semidirect C_3 product with neutral Vec_{V_4}. Its action class has order three, as in Proposition 4.16. Since the tensor-automorphism classes are $C_2 \times S_3$, this class is a pure order-three group action. The remaining action coherence lies in $H^2(C_3, \widehat{V_4}) = 0$ by coprimality [57, Proposition 3.2]. Hence the action is equivalent to the strict group action and the dual is the untwisted Vec_{A_4}. Its graded tensor section is part of Galindo's semidirect-product conclusion; no uncontrolled C_3 associator twist is discarded.

The equivariantization $\mathcal{D}$ is Morita equivalent to $\mathcal{E} \rtimes C_2$. Morita transport gives a C_2 extension $\mathcal{T}$ of Vec_{A_4} with the same center as $\mathcal{D}$. The center of Vec_{A_4} has exactly three Lagrangian subcategories, all in the orbit of the canonical $\mathrm{Rep}(A_4)$ [57, Proposition 7.6 and Section 8.3]. Consequently each has untwisted Vec_{A_4} as pointed Morita dual. The involution fixes at least one of these three. For its invariant pointed module $\mathcal{N}$, the dual of $\mathcal{T} \boxtimes_{\mathrm{Vec}_{A_4}} \mathcal{N}$ is a C_2 crossed product with pointed neutral

Vec_{A_4} [58, Proposition 5.3]; every component is pointed. This step requires module-class invariance, rather than a coherent extension of $\mathcal{N}$. It gives a pointed Morita group

$$1 \longrightarrow A_4 \longrightarrow G \longrightarrow C_2 \longrightarrow 1.$$

Because A_4 is centerless with $\mathrm{Out}(A_4) = C_2$, G is S_4 or $A_4 \times C_2$. The latter has six identity-supported central characters for every associator. But $\mathrm{Inv}(Z(\mathcal{D})) \to \mathrm{Inv}(\mathcal{D})$ is injective: its kernel is $\mathrm{Hom}(U(\mathcal{D}), \mathbb{C}^\times) = 1$. The four source invertibles therefore exclude $A_4 \times C_2$. Thus $\mathcal{D}$ is Morita equivalent to $\mathrm{Vec}^{\omega}_{S_4}$.

The only normal abelian subgroups of S_4 are 1 and its V_4. Lemma 4.14 makes every extension of $\mathcal{D}$ integral, contradicting the two odd dimensions $2\sqrt{3}$. The exact checker validates all four source tensors, the small-ring exhaustion, module-rank alternatives and all thirty subgroups of S_4. □

5. Further general obstructions

The following arguments exclude nine further source tensors without a pivotality assumption. The proofs use a conductor bound, a Frobenius–Perron index, cyclic-module divisibility, and changes of Morita representative.

Proposition 5.1. *The rings* R8_2_35, R8_2_38, R8_2_346 *and* R8_2_406 *have no categorification over* $\mathbb{C}$.

Proof. Each tensor has four invertibles and four simples of FP dimension $d = 3 + \sqrt{10}$. By the published [25, Lemma 3.1], existence of any categorification implies existence of a pseudo-unitary Galois conjugate, whose canonical spherical structure has global dimension

$$D = 4(1 + d^2) = 80 + 24\sqrt{10}, \qquad DD' = 640.$$

Its simple-dimension and global-dimension fields both equal $\mathbb{Q}(\sqrt{10})$. The nontrivial quadratic character has conductor 40, so [63, Theorem 5.1] requires

$$\left|\frac{1}{D} - \frac{1}{D'}\right|^2 \leq \frac{1}{40}.$$

The left side is $9/160 > 1/40$, a contradiction. □

Proposition 5.2. *The ring* R8_2_209 *has no categorification over* $\mathbb{C}$.

Proof. Put $d = (5 + \sqrt{29})/2$. The source dimension vector is $(1, 1, 1, d, d-1, d+1, d, d)$, and its first three basis elements form a pointed subcategory of dimension 3. Hence

$$\mathrm{FPdim}(\mathcal{C}) = 5(1 + d^2) = \frac{145 + 25\sqrt{29}}{2}.$$

By [21, Proposition 8.15], its quotient by 3 must be an algebraic integer. Its field trace is $145/3$, which is not an integer. □

Lemma 5.3 (Cyclic-module divisibility)**.** *Let $H \cong C_p$, with p prime, be a group of invertibles in a fusion category over $\mathbb{C}$. Suppose that simple objects X, Y, Z satisfy $hX \cong X$, $Yh \cong Y$, and $hZ \cong Z \cong Zh$ for every $h \in H$. If $\mathrm{FPdim}(Z)/p$ is not an algebraic integer, then $p \mid N_{XY}^{Z}$.*

Proof. The rank-one module category generated by X trivializes the pointed associator on H. Thus $A = \bigoplus_{h \in H} h$ admits its regular connected separable algebra structure. Since $H^2(C_p, \mathbb{C}^\times) = 0$, the stabilizer cocycles can be trivialized to give X a left A-module structure and Y a right A-module structure. Hence $X \otimes Y$ is an A-bimodule.

The A-bimodules supported on the fixed simple Z correspond to projective representations of C_p^2: the projective-representation dimension equals the underlying multiplicity of Z. A scalar two-cocycle on C_p^2 has either zero or nondegenerate commutator pairing. Its twisted group algebra is accordingly $\mathbb{C}^{p^2}$ or $M_p(\mathbb{C})$. The first possibility would give a simple bimodule of FP dimension

$\mathrm{FPdim}(Z)/\mathrm{FPdim}(A) = \mathrm{FPdim}(Z)/p$, contrary to algebraic integrality. In the second possibility every simple bimodule supported on Z has underlying object pZ. Decomposing the Z-isotypic bimodule summand of $X \otimes Y$ proves the assertion. This is the regular cyclic-algebra method of [64, Lemmas 6.1–6.2], with the stabilizer cocycle retained rather than assumed trivial. □

For later use, a free left C_p orbit whose right orbit agrees with it has cyclic left-right stabilizer of order p. Its block in A-*bimod*($\mathcal{C}$) therefore consists of p simples of the original FP dimension. A label fixed on both sides instead contributes either p^2 simples of dimension $\mathrm{FPdim}(X)/p$ or one simple of dimension $\mathrm{FPdim}(X)$, by the preceding proof.

Proposition 5.4. *The ring R*8_2_260 *has no categorification over* $\mathbb{C}$.

Proof. Put $d = 3+2\sqrt{3}$. Its dimension vector is $(1, 1, 1, d, d, d, d, 3)$, and the first three basis elements form C_3. They fix label 3 on both sides. Since $\mathrm{Norm}(d/3) = -1/3$, Lemma 5.3 requires $3 \mid N_{33}^3$. The source tensor has $N_{33}^3 = 2$. □

Proposition 5.5. *The rings R*8_2_380 *and R*8_2_673 *have no categorification over* $\mathbb{C}$.

Proof. Write $x = 4+\sqrt{6}$ and $y = 3+2\sqrt{6}$. Each ring has a subcategory $\mathcal{E}$ with the representation ring of the dihedral group D_{10} of order ten. It occupies labels $(0, 3, 4, 5)$ in *R*8_2_380 and $(0, 1, 6, 7)$ in *R*8_2_673. Its nontrivial invertible g fixes the two dimension-two labels, so the regular algebra $A = 1 + g$ exists. By [64, Lemma 6.2 and the proof of Theorem 6.3], A-*bimod*($\mathcal{E}$) is pointed with group D_{10}.

The outside labels form two free g-orbits, of dimensions x and y, with the same left and right actions. They are $(1, 2), (6, 7)$ in the first tensor and $(2, 5), (3, 4)$ in the second. The cyclic stabilizer calculation gives two x-dimensional and two y-dimensional simples in the dual category $\mathcal{B} = A$-*bimod*($\mathcal{C}$); all its other simples are the ten invertibles of D_{10}.

The rotation subgroup $H = C_5$ fixes every noninvertible on both sides, since each dimension fiber has only two elements. Moreover

$$\mathrm{Norm}(x/5) = \frac{2}{5}, \qquad \mathrm{Norm}(y/5) = -\frac{3}{5}.$$

For an x-dimensional simple X, Lemma 5.3 therefore makes every noninvertible multiplicity in XX^* divisible by 5. Write $x^2 = a + bx + cy$ for the resulting dimension equation, where a, b, c are nonnegative integers. The $\sqrt{6}$ coefficient gives $b + 2c = 8$, contradicting $5 \mid b, c$. □

Proposition 5.6. *The ring R*8_2_745 *has no categorification over* $\mathbb{C}$.

Proof. Put $d = 1+\sqrt{2}$ and let g be label 1. The dimension vector is $(1, 1, d, d, 2, 2, 2d, 2d)$. The subsets $\mathcal{D} = (0, 1, 2, 3)$ and $\mathcal{E} = (0, 1, 4, 5)$ are fusion subcategories, the latter with the representation ring of the dihedral group D_{10} of order 10. The element g exchanges labels $2, 3$ and fixes labels $4, 5, 6, 7$ on both sides. Thus $A = 1 + g$ exists.

In $\mathcal{B} = A$-*bimod*($\mathcal{C}$), the subcategory $\mathcal{E}' = A$-*bimod*($\mathcal{E}$) is pointed with group D_{10}, by [64, Lemma 6.2 and the proof of Theorem 6.3]. The subcategory $\mathcal{D}' = A$-*bimod*($\mathcal{D}$) has two invertibles and two simples of dimension d, by the cyclic-stabilizer calculation above. Each of labels $6, 7$ contributes either four simples of dimension d or one of dimension $2d$. These are all the simples of $\mathcal{B}$, so it has precisely ten invertibles and 2, 6 or 10 simples of dimension d.

For a d-dimensional simple X, the dimension of XX^* is $a + bd$ with $a, b \in \mathbb{Z}_{\geq 0}$, where a is its invertible stabilizer order. The identity $d^2 = 1 + 2d$ forces $a = 1$. Thus the D_{10} action on these simples is free, and their number must be 10. It follows that $\mathcal{B}$ is generalized near-group, with no $2d$-dimensional simples remaining.

By [65, Proposition 3.6(1)], the adjoint subcategory $\mathcal{B}_{\mathrm{ad}}$ is nonpointed and is contained in every nonpointed fusion subcategory, hence in $\mathcal{D}'$. There is no proper nonpointed fusion subcategory of $\mathcal{D}'$: one noninvertible would give the forbidden rank-two rule $X^2 = 1 + 2X$ [23], while two noninvertibles force the other invertible by reciprocity. Therefore $\mathcal{B}_{\mathrm{ad}} = \mathcal{D}'$. Its order-two invertible group must

be normal in D_{10}, since conjugation preserves the adjoint subcategory. No such normal subgroup exists. □

The source-bound verifiers in `evidence/theoretical_20261016/rank8` check all dimension characters and subcategory tensors used above, including the indicated left and right actions. The categorical implications are the preceding proofs and their explicitly cited dependencies.

6. An anti-invariant module obstruction

We give eight further exclusions that require no pivotal structure.

Lemma 6.1. *Let g be an invertible simple of order two in a fusion category, and let $x_1, \dots, x_f$ be all simples fixed by right tensoring by g, where $f > 0$. There is an integral f-dimensional representation T of the fusion ring such that, with $B_X = (N_{Xx_a}^{x_b})_{a,b}$,*

$$|(T_X)_{ab}| \le (B_X)_{ab}, \qquad (T_X)_{ab} \equiv (B_X)_{ab} \pmod 2, \qquad T_{X^*} = T_X^t.$$

In particular T_X is normal if $XX^* = X^*X$.

Proof. A fixed simple supplies a rank-one module category over the pointed category generated by g, trivializing its associator. Hence the regular connected separable algebra $A = \mathbf{1} + g$ exists; compare [64, Lemmas 6.1–6.2]. Each fixed simple has two simple right A-module structures m_a^+, m_a^-, whereas each free right orbit gives one simple module. Twisting the right action by the algebra automorphism that acts as -1 on g exchanges m_a^+ with m_a^- and fixes the free-orbit modules. It commutes with left tensoring by every object. Thus the anti-invariant subgroup has basis $v_a = [m_a^+] - [m_a^-]$. The two lift multiplicities have sum $(B_X)_{ab}$ and difference $(T_X)_{ab}$, proving the bound and parity. Rigidity gives the transpose relation, since the orthogonal difference vectors all have squared norm two. All fusion relations are preserved. □

Proposition 6.2. *The rings R8_2_318, R8_2_319, R8_2_320, R8_2_368, R8_2_370, R8_2_735, R8_2_758 and R8_2_760 have no categorification over $\mathbb{C}$.*

Proof. Set $q = z^2 - 6z + 1$, $r = z^2 + 2z - 1$, and $h = z^2 - 6z - 1$, and put

$$A = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 2 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 2 \end{pmatrix}, \quad C = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 2 \end{pmatrix}.$$

The following source identities use zero-based labels. Each indicated X commutes with X^*; the last column is a polynomial vanishing on X. The stated fixed labels are all the right-fixed simples of g.

suffix	g	fixed labels	X	B_X	polynomial
318	4	$(1,2,3,7)$	5	A	$(z^2-1)qr$
319	4	$(1,2,3,7)$	5	A	$(z^2-1)q$
320	6	$(1,2,5,7)$	3	A	$(z^2-1)(z^2+1)q$
368	3	$(4,5,6,7)$	1	C	$(z^2-1)(z^2+1)h$
370	3	$(4,5,6,7)$	1	B	$(z^2-1)(z^2+1)q$
735	1	$(4,5,6,7)$	2	B	$(z^2-1)qr$
758	1	$(4,5,6,7)$	2	C	$(z^2-1)h$
760	1	$(4,5,6,7)$	2	B	$(z^2-1)q$

Let $T = T_X$ from Lemma 6.1. Normality gives $\|T\|_F^2 = \sum_{j=1}^4 |\lambda_j|^2$. Integrality forces conjugate quadratic eigenvalues to occur with equal multiplicity. For bounds A, B, parity gives $\|T\|_F^2 \in \{12, 16\}$. The roots $3 \pm 2\sqrt{2}$ of q contribute 34 and so are absent. For suffixes $319, 320, 370, 760$, all remaining roots have modulus one, forcing $\|T\|_F^2 = 4$, a contradiction.

For $318,735$, the additional roots $-1 \pm \sqrt{2}$ contribute 6. The possible squared norms are therefore $4, 8, 12$, according to the number of copies of this pair. Only two copies are possible, giving norm squared 12 and trace -4. But norm squared 12 forces the last diagonal entry to vanish, while the other diagonal entries are two signs and a zero. Their trace lies in $[-2, 2]$, a contradiction.

For bound C, the squared norm is in $\{12, 16, 20\}$. The roots $3 \pm \sqrt{10}$ of h contribute 38 and are absent. All remaining roots for $368,758$ have modulus one, again giving the impossible squared norm 4. These arguments give unrestricted exclusions for $318,735$, in addition to their pivotal obstructions. $\square$

The portable verifier `evidence/theoretical_20261016/rank8/verify_anti_module_eight.py` checks the displayed source identities and spectral arithmetic directly; the proof does not rely on an enumeration of candidate matrices.

6.1. **Ordered exclusion ledger.** An excluded tensor is removed after the first verified certificate in the following order. For each row, "left" still counts rings subsequently recognized as positive, so this is an exclusion table rather than a table of outstanding questions.

Criterion	New $m=1$	New $m=2$	Left $m=1$	Left $m=2$
Near-group stabilizer	1	1	95	876
Rank-two closed subring	0	10	95	866
Zero-spectrum tensor condition	7	80	88	786
Proved rank 2–6 subring	3	121	85	665
Four proved rank-seven subrings	0	12	85	653
Commutative codegree d-number	23	462	62	191
Rank-eight noncommutative codegree	1	17	61	174
Integral formal-codegree divisibility	1	6	60	168
Published no-induction-matrix lemma	2	0	58	168
Involution-module obstruction	0	2	58	166
Field-automorphism obstruction	0	0	58	166
Fusion-subcategory dimension quotient	0	1	58	165
Primitive formal-codegree test	0	0	58	165
Cyclotomic eigenvalue test	0	0	58	165
Complete oriented signed quotient	3	71	55	94
Graded Brauer–Picard obstruction	1	2	54	92
Rank-one fiber-functor obstruction	0	2	54	90
Solvability and equivariantization	3	0	51	90
Prime descent and Hopf obstructions	0	9	51	81
Dimension-96 two-object orbit	0	4	51	77
Dimension-108 order-four action	0	1	51	76
Galois-invariant pointed associator	0	3	51	73
Normal abelian square-order kernels	0	8	51	65
Dimension-block conductor bound	0	4	51	61
Quadratic subcategory index	0	1	51	60
Cyclic-module divisibility (direct or Morita)	0	3	51	57
Morita adjoint normality	0	1	51	56
Conductor-32 signed Jacobi obstruction	0	1	51	55
Anti-invariant involution module	0	8	51	47
Algebraic $2D2$ graph-pair rigidity	0	4	51	43
Odd-involution Morita	0	1	51	42
Twisted half-braiding support	0	2	51	40
Quantum-group inverse Morita	0	1	51	39
Pivotal parity and inverse Morita	1	0	50	39
Fibonacci character parity	0	2	50	37
Conductor-seventeen indicator balance	0	5	50	32
Weighted Galois third indicator	0	1	50	31
Conductor-forty center trace	0	1	50	30
Cyclic induction phase multiplicity	1	0	49	30
Full center STS and fourth moment	1	0	48	30
Inner square and tensor-functor rigidity	0	1	48	29

The 77 tensors left by the exclusion ledger are exactly the positive models. Every negative stage is a theorem applied to tensor data or an exact based-tensor transfer of a published and proved no-go. Proposition 4.1 and the integral-divisibility stage both have certified first applications already at

multiplicity one; the latter also resolves two multiplicity-two cases not excluded by the preceding criteria. These *first appearances* refer only to the two multiplicities of the complete rank-eight census. Nothing in this calculation establishes a multiplicity from which no new criterion is needed.

7. Exact positive models

Each standard family in this section admits a unitary complex realization. A pointed group ring of order eight is $K_0(\mathrm{Vec}_G)$ for the group reconstructed from its tensor. The five groups are C_8, $C_4 \times C_2$, C_2^3, D_8 and Q_8. The near-group ring $C_7 + 0$ is realized by a Tambara–Yamagami category [41]. If A is a rank-four unitary model from the complete catalogue of [14] and B is either Vec_{C_2} or Fibonacci, the Deligne product $A \boxtimes B$ is a unitary rank-eight model with multiplication tensor $N_A \otimes N_B$. Its multiplicity is at most two precisely for the displayed matches.

The three group-theoretical models $\mathcal{C}(F_5, 1, D_5, 1)$, $\mathcal{C}(S_4, 1, A_4, 1)$, and $\mathcal{C}(\mathrm{PSL}(2,7), 1, C_7 \rtimes C_3, 1)$ are untwisted group-theoretical fusion categories, hence admit unitary realizations. Their exact appendix tensors are matched against the census in all 512 positions, with a stored basis permutation for every match [20, Appendix C.3.2(3), (6), (11)]. Some admit a second subgroup construction, also recorded in the model catalogue.

Proposition 7.1 (Representation and quantum-group templates)**.** *All three character-ring templates* $\mathrm{Rep}(D_{20})$, $\mathrm{Rep}(D_{26})$, *and* $\mathrm{Rep}(\mathrm{Dic}_5)$ *have rank eight and multiplicity one. The standard unitary* $SU(2)_7$ *category, and the even subcategories of* $SU(2)_{14}$ *and* $SU(2)_{15}$*, also have rank eight and multiplicity one. Their generated multiplication tensors occur in the supplied census and are matched by full based-ring isomorphisms.*

Proof. For $SU(2)_k$, the simples are the labels $0, \dots, k$ and

$$N_{ab}^c = 1 \quad \Longleftrightarrow \quad |a-b| \le c \le \min(a+b, 2k-a-b), \qquad a+b-c \equiv 0 \pmod 2;$$

all other coefficients are zero. The even labels close under multiplication and yield eight objects for $k = 14, 15$, while $k = 7$ already has eight labels [29]. The explicit tensor generator implements this formula directly.

For the dihedral group of order $2n$, take $n = 10$ or 13. Its two-dimensional characters ρ_j obey $\rho_i\rho_j = \rho_{i+j} + \rho_{|i-j|}$, with $\rho_0 = 1 + \beta$ and $\rho_{n-j} = \rho_j$. For $n = 10$, $\rho_5 = \alpha + \alpha\beta$, the four linear characters form C_2^2, $\beta\rho_j = \rho_j$ and $\alpha\rho_j = \rho_{5-j}$. For $n = 13$ there are two linear characters, with $\beta\rho_j = \rho_j$. Thus their ranks are respectively $4+4$ and $2+6$. For Dic_5 of order twenty the four linear characters χ_i form C_4, with

$$\rho_0 = \chi_0 + \chi_2, \quad \rho_5 = \chi_1 + \chi_3, \quad \rho_{10-j} = \rho_j, \quad \chi_i\rho_j = \begin{cases} \rho_j & i \text{ even}, \\ \rho_{5-j} & i \text{ odd}; \end{cases}$$

the same product $\rho_i\rho_j = \rho_{i+j} + \rho_{|i-j|}$ completes the ring. These are ordinary finite-group character identities [42]. The supplied generator expands each product and compares all 8^3 ordered coefficients after unit-fixing relabelling. A match is a ring isomorphism and the corresponding representation category supplies the construction. □

Multiplicity	Ring ID	One exact unitary model
1	`R8_1_1`	`Rep(Dic5)`
1	`R8_1_3`	$\mathcal{C}(C_5 \rtimes C_4, \omega_2, C_2, 1)$
1	`R8_1_4`	`Vec(G), |G|=8`
1	`R8_1_5`	`Vec(C4) x Fib`
1	`R8_1_6`	Bosonic quotient of $PSU(2)_6 \boxtimes \mathcal{C}(C_4, q)$
1	`R8_1_7`	Central square-root extension of pseudo-$PSU(2)_6$
1	`R8_1_10`	`Vec(G), |G|=8`
1	`R8_1_11`	Central-twisted flip extension of pseudo-$PSU(2)_6$

Multiplicity	Ring ID	One exact unitary model
1	R8_1_12	Isotropic orbifold of $NG(C_{16}, 16)$
1	R8_1_13	Vec(G), |G|=8
1	R8_1_14	Central-twisted flip extension of $PSU(2)_6$
1	R8_1_15	$\mathrm{HI}(C_4)$
1	R8_1_16	Vec(G), |G|=8
1	R8_1_19	$(\mathrm{Fib} \boxtimes \mathrm{Fib}) \rtimes C_2$
1	R8_1_20	TY(C7)
1	R8_1_22	Diagonal parity in $\mathrm{TY}(C_3) \boxtimes \mathrm{Vec}_{C_4}$
1	R8_1_23	$\mathrm{Vec}_{Q_8}^{C_3}$
1	R8_1_24	C(F5,1,D5,1)
1	R8_1_25	$PSU(2)_6 \rtimes C_2$ with the flip action
1	R8_1_27	TY(C3) x Vec(C2)
1	R8_1_30	$\mathrm{TY}(C_6)^{C_2}$ with odd coherence -1
1	R8_1_32	pseudo-PSU(2)_6 x Vec(C2)
1	R8_1_34	$\mathrm{TY}(C_9)^{C_2}$ with odd coherence -1
1	R8_1_35	$\mathrm{TY}(C_3^2)^{C_2}$ with odd coherence -1
1	R8_1_36	TY(C3) crossed C2
1	R8_1_39	TY(C3) x Fib
1	R8_1_42	Pseudo-$PSU(2)_6 \rtimes C_2$ with the flip action
1	R8_1_43	pseudo-PSU(2)_6 x Fib
1	R8_1_53	Isotropic orbifold of $NG(C_4^2, 16)$
1	R8_1_59	Rep(D10) x Vec(C2)
1	R8_1_61	One-half SO(12)_2
1	R8_1_65	Vec(G), |G|=8
1	R8_1_66	Vec(C2xC2) x Fib
1	R8_1_67	PSU(2)_6 x Vec(C2)
1	R8_1_69	$\mathrm{HI}(C_2^2)$
1	R8_1_70	Fib x Vec(C2) x Fib
1	R8_1_71	PSU(2)_6 x Fib
1	R8_1_72	PSU(2)_7 x Vec(C2)
1	R8_1_73	PSU(2)_14
1	R8_1_78	SO(9)_2
1	R8_1_79	Z(Vec(S3))
1	R8_1_83	Integral part of the $m = 0$ triality gauging
1	R8_1_85	Neutral part of S3-gauged SO(8)_1
1	R8_1_91	Rep(D_{26})
1	R8_1_92	PSU(2)_15
1	R8_1_93	Rep(D10) x Fib
1	R8_1_94	PSU(2)_7 x Fib
1	R8_1_95	Fib x Fib x Fib
2	R8_2_37	Nonsplit C_2 extension of the unitary $2D2$ category
2	R8_2_39	Asaeda–Haagerup AH_4
2	R8_2_143	$\mathrm{Vec}_{C_{17}}^{C_4}$
2	R8_2_154	$\mathrm{TY}(C_2^2)^{S_3}$ with parity coherence
2	R8_2_174	Rep(A4) x Vec(C2)
2	R8_2_179	$\mathrm{Vec}_{C_2^4}^{C_5}$
2	R8_2_202	C(S4,1,A4,1)
2	R8_2_212	Rep(A4) x Fib
2	R8_2_237	$\mathrm{Rep}(A\Gamma\mathrm{L}(1,8))$
2	R8_2_239	C_3-algebra Morita dual of an S_3 extension of Vec_{C_7}
2	R8_2_241	$\mathcal{C}(\mathrm{PSL}(2,7), 1, C_7 \rtimes C_3, 1)$
2	R8_2_244	$\mathrm{Vec}_{C_4^2}^{C_3}$

Multiplicity	Ring ID	One exact unitary model
2	R8_2_289	$\mathrm{Vec}_{C_2^4}^{C_3}$
2	R8_2_290	Charge-conjugation defects of $PSU(3)_4$
2	R8_2_313	Diagonal bosonic quotient of the two pseudo-$PSU(2)_6$ categories
2	R8_2_477	even 2D2 x Vec(C2)
2	R8_2_484	Asaeda–Haagerup AH_5
2	R8_2_491	even 2D2 x Fib
2	R8_2_493	Haagerup H1 x Vec(C2)
2	R8_2_535	Dual of the $1+\rho$ Q-system in $HI(C_6)$
2	R8_2_557	Diagonal bosonic quotient of $PSU(2)_6 \boxtimes PSU(2)_6$
2	R8_2_575	Davydov–Runkel extension of $\mathrm{Rep}(H_{C:\sigma_1})$
2	R8_2_611	$\mathrm{TY}(C_2^2)^{S_3}$ with strict action
2	R8_2_666	Pointed centralizer in $Z(NG(C_4,4))$
2	R8_2_820	$NG(C_5,5)$ center factor, auxiliary C_3^2
2	R8_2_839	Haagerup H1 x Fib
2	R8_2_855	Even D_{14} category
2	R8_2_868	Dual Haagerup–Izumi C_7 category
2	R8_2_869	$NG(C_5,5)$ center factor, auxiliary C_9

Five further multiplicity-one models are imported from exact AnyonWiki tensor matches: R8_1_36 is the crossed product of $\mathrm{TY}(C_3)$ by the order-two inversion action; R8_1_61 is the rank-eight fusion subcategory called one-half $SO(12)_2$; R8_1_78 is $SO(9)_2$; R8_1_79 is $Z(\mathrm{Vec}_{S_3}) \simeq \mathrm{Rep}(D(S_3))$; and R8_1_85 is the neutral component of the universal $\mathbb{Z}_2$ grading after gauging the full S_3 symmetry of $SO(8)_1$ [1, 2]. For the crossed product, the standard bicharacter $\chi(a,b) = \exp(2\pi iab/3)$ and positive $\tau = 1/\sqrt{3}$ are invariant under simultaneous inversion. The induced tensor action squares to the identity, and the crossed product is unitary. Its independently constructed tensor is matched in all 512 positions. The other four are standard unitary categories or fusion subcategories thereof; their complete sourced tensors and unit-preserving permutations are included in the positive catalogue. Existence of these models proves general complex categorifiability. No negative conclusion is imported merely from an AnyonWiki flag; R8_1_26 is excluded by the theoretical proof in Proposition 4.3.

7.1. **Twenty-two further constructions from extensions and subfactors.** The following constructions are verified by generating their multiplication tensors directly and comparing all 512 coefficients. No associator search is needed.

For $G = C_4$ and $G = C_2^2$, write $X_g = g\rho$ and use

$$gh = g + h, \qquad gX_h = X_{g+h}, \qquad X_g h = X_{g-h}, \qquad X_g X_h = (g-h) + m\sum_{a\in G} X_a.$$

At $m = 1$, Izumi's unitary generalized Haagerup constructions [8, Section 9] give R8_1_15 and R8_1_69, respectively. At $m = 2$, these are the Asaeda–Haagerup rings AH_4 and AH_5, giving R8_2_39 and R8_2_484. The C_4 construction is [9, Theorem 4.9]; existence of the C_2^2 category follows from its Lemma 3.15, Theorem 3.17 and Section 4. These are unitary subfactor categories.

The ring R8_1_19 is $(\mathrm{Fib} \boxtimes \mathrm{Fib}) \rtimes C_2$, using the canonical action exchanging the two factors. The ring R8_1_22 is the fusion subcategory of $\mathrm{TY}(C_3) \boxtimes \mathrm{Vec}_{C_4}$ whose simple objects have matching parity: (a, z^{2j}), (m, z) and (m, z^3). Both operations preserve unitarity.

For R8_1_30, take the standard unitary $\mathrm{TY}(C_6)$ with bicharacter $\chi(a,b) = \exp(2\pi iab/6)$ and positive parameter $\tau = 1/\sqrt{6}$. Inversion fixes m, preserves every standard associator coefficient, and gives a tensor functor T with identity tensorator. Equip the C_2 action with coherence $T^2 \Rightarrow \mathrm{Id}$ equal to $+1$ on invertibles and -1 on m. This is a T-invariant monoidal natural automorphism, since the TY grading is $\mathbb{Z}/2$. In the equivariantization, the two structures on m have eigenvalues $\pm i$. The tensor decomposition is obtained by splitting the permutation action on each fusion space into its

two characters; in particular two odd eigenvalues contribute the additional minus sign. The result is the stated full tensor, with four invertibles, two objects of dimension two, and two of dimension $\sqrt{6}$.

Let $F_{20} = C_5 \rtimes C_4$ with

$$(a,b)(c,d) = (a + 2^b c,\ b + d), \qquad \omega(x,y,z) = (-1)^{b(x)\lfloor (b(y)+b(z))/4 \rfloor},$$

where the two coordinates are reduced modulo five and four, respectively, and representatives $0 \le b < 4$ are used in the cocycle. For $H = \langle (0,2) \rangle$, the restriction $\omega|_H$ is trivial. The unitary group-theoretical category $\mathcal{C}(F_{20}, \omega, H, 1)$ realizes *R*8_1_3. The supplied code constructs the eight simple homogeneous $\mathbb{C}[H]$-bimodules and their balanced tensor products. All phases are fourth roots of unity and are stored as integer exponents modulo four; the cocycle identity, bimodule coherence, balanced quotient, decomposition multiplicities, and full census tensor match are checked exactly.

The ring *R*8_1_6 has a bosonic quotient construction. Let $\mathcal{D} = PSU(2)_6$, with simples $1, f, \rho, f\rho$, where f is a fermion and $\rho^2 = 1 + \rho + f\rho$. Let $\mathcal{P} = \mathcal{C}(C_4, q)$ be the unitary pointed modular category with $q(z^a) = \exp(\pi i a^2/4)$. Then $b = f \boxtimes z^2$ in $\mathcal{B} = \mathcal{D} \boxtimes \mathcal{P}$ is an order-two boson: its self-braiding is $+1$. It generates a Tannakian $\mathrm{Rep}(C_2)$, whose regular algebra is $A = 1 \oplus b$. The central inclusion supplied by the braiding gives a fusion category $\mathcal{B}_A$ of all right A-modules, with tensor free-module functor F [22, Theorems 8.4.9 and 8.23.3]. The unitary crossed-product construction preserves its positive $*$-structure [40]. The boson is not transparent in $\mathcal{B}$; thus this uses all modules, rather than only local modules, and asserts no descended braiding.

Tensoring by b acts freely on the sixteen simples by $(V_j, z^a) \mapsto (V_{6-j}, z^{a+2})$, with $j \in \{0,2,4,6\}$. By adjunction every free module $F(V_j \boxtimes z^a)$ is simple, and two are isomorphic exactly when their labels belong to the same orbit. Dominance of F gives precisely eight simples. With $g = F(1 \boxtimes z)$ and $r = F(\rho \boxtimes 1)$, their rules are

$$g^4 = 1, \qquad gr = rg, \qquad r^2 = 1 + r + g^2 r.$$

The ordered model basis $(1, g, g^2, g^3, r, gr, g^2 r, g^3 r)$ maps to census labels $(0,1,7,2,5,3,6,4)$. The verifier reconstructs all coefficients from the truncated $SU(2)_6$ rules and the eight free orbits.

For *R*8_1_7, use either of the two unitary pseudo-$PSU(2)_6$ categories $\mathcal{D}$ of [30, Theorem 4.1]. Their basis $1, g, x, y$ obeys $g^2 = 1$, $gx = y$, $x^2 = y^2 = g + x + y$ and $xy = 1 + x + y$. The two source-bound AnyonWiki associator tables, labelled $(4,1,2,4,1,0,1)$ and $(4,1,2,4,2,0,1)$, are included in the supplement. Over $\mathbb{Q}(\zeta)$, $\zeta^4 + 1 = 0$, the verifier checks all 224 coefficients, 136 invertible associator blocks, unit and rigidity conditions and 2688 pentagons for each table.

Write $a(i) = gi$ and let $F_\ell^{ijk}[e,f]$ take left intermediate e to right intermediate f. In the two tables respectively, the scalars

$$(\lambda_1, \lambda_g, \lambda_x, \lambda_y) = (1, \zeta^2, \zeta, \zeta^3), \qquad (1, -\zeta^2, -\zeta^3, -\zeta)$$

define a half-braiding of g. The exact check is the following identity for each of the 24 fusion channels $N_{ij}^k = 1$:

$$\lambda_i \lambda_j F^{igj}_{a(k)}[a(i), a(j)] = \lambda_k F^{gij}_{a(k)}[a(i), k] F^{ijg}_{a(k)}[k, a(j)].$$

Thus $\widetilde{g} = (g, c)$ is an invertible object of $Z(\mathcal{D})$. Its self-braiding satisfies $\lambda_g^2 = -1$, so a fermionic condensation argument would be inappropriate. The equivalence between the unitary and ordinary centers permits a unitary representative of this lift [32, corrected Remark 3.1].

Instead take the trivial homomorphism $C_2 \to \mathrm{BrPic}(\mathcal{D})$ and modify its product equivalences by the normalized central 2-cocycle $t(1,1) = \widetilde{g}$, with other normalized values the unit. The only nontrivial cocycle identity compares $\widetilde{g}$ with itself, regardless of its order in $\mathrm{Inv}\, Z(\mathcal{D})$. The remaining extension obstruction belongs to $H^4(C_2, \mathbb{C}^\times) = 0$ [7, Theorem 1.3]; this is also the trivial-extension zesting of [31, Section 6.2].

This extension is unitary. Choose regular unitary $\mathcal{D}$-bimodule components and unitary multiplication equivalences from the unitary lift. Preliminary associativity intertwiners can be chosen unitary by polar decomposition, so their normalized pentagon defect is $U(1)$-valued. A normalized

3-cochain on C_2 is determined by u at $(1,1,1)$, and its coboundary at $(1,1,1,1)$ is u^2. A unit-modulus square root therefore removes the defect while keeping every coherence map unitary. The resulting regular-component extension is a unitary fusion category; this argument uses no general assertion that arbitrary extensions of unitary categories are unitarizable.

Its simples are (X,a), $a \in C_2$, with

$$(X,a)(Y,b) = (XYg^{ab}, a+b).$$

For $z=(1,1)$, this gives $z^2 = g$, $zx = xz$ and $x^2 = z^2 + x + z^2 x$. The model basis $(1,g,x,y,z,zg,zx,zy)$ maps to census labels $(0,7,3,4,1,2,6,5)$, and all 512 coefficients agree. The exact associator and central-lift replay is invoked by the finite-update verifier.

The same two rank-four bases have nontrivial flip extensions. For $\mathcal{D} = PSU(2)_6 = \mathrm{Ad}(A_7)$, a tensor autoequivalence τ exchanges ρ and $f\rho$ [33, Lemma 4.10]. Its square is tensor-isomorphic to the identity. The fusion rules imply $\mathrm{Aut}_\otimes(\mathrm{Id}_\mathcal{D}) = 1$, so this square isomorphism is unique after normalization and obeys the C_2 action coherence. Reutter's unitarization and automatic-unitarity theorems give a unitary representative of the entire action [34, Theorems 1–2].

For the second pseudo-$PSU(2)_6$ source table, the permutation $\tau = (x\ y)$ also gives a coherent tensor action, now checked explicitly. Its tensorator J^k_{ij} is 1 when an input is the unit. On the seventeen other admissible channels, ordered lexicographically, write $J = (\sqrt{2})^a \zeta^b$; the exponent vectors are

$$a = (-1,-1,0,-1,-1,-1,-1,-1,0,-1,0,-1,0,-1,0,0,0),$$
$$b = (29,15,10,17,1,5,5,11,0,1,16,15,4,5,0,0,0).$$

All 224 tensor-functor identities hold over $\mathbb{Q}(\zeta)$. The natural isomorphism $\eta : \tau^2 \Rightarrow \mathrm{Id}$ has scalars

$$\eta = \left(1, \frac{1+\zeta^2}{2}, -\frac{1+\zeta^2}{2}, -\frac{1+\zeta^2}{2}\right)$$

in the order $(1,g,x,y)$. The verifier checks all 24 equations $\eta_i \eta_j = \eta_k J^k_{ij} J^{\tau(k)}_{\tau(i),\tau(j)}$ and $\eta_{\tau(i)} = \eta_i$. This is therefore a coherent action, not just a symmetry of the ring. Transport to unitary functors preserves its coherence; the transported monoidal natural isomorphism is automatically unitary by [34, Theorems 1–2].

For either rank-four base, its nontrivial invertible has a unique central lift: the kernel of $\mathrm{Inv}\, Z(\mathcal{D}) \to \mathrm{Inv}\, \mathcal{D}$ is $\mathrm{Aut}_\otimes(\mathrm{Id}_\mathcal{D}) = 1$, and a lift was already constructed above. The flip fixes this lift. Thus one may use either the crossed product or its central 2-cocycle modification. The remaining C_2 obstruction is removed by the unitary phase correction described above. With v the nontrivial invertible, their products are

$$(X,a)(Y,b) = (X\tau^a(Y) v^{\epsilon ab}, a+b), \qquad \epsilon \in \{0,1\}.$$

For $PSU(2)_6$, these give $R8_1_25$ at $\epsilon = 0$ and $R8_1_14$ at $\epsilon = 1$; for pseudo-$PSU(2)_6$, they give $R8_1_42$ and $R8_1_11$, respectively. The exact tensorator and coherent-square replay is `software/verify_pseudo_psu_swap.py`.

The ring $R8_1_23$ is $\mathrm{Vec}^{C_3}_{Q_8}$ for the strict automorphism $i \mapsto j \mapsto k \mapsto i$ of the quaternion group. The fixed objects $1, -1$ each have three equivariant structures, giving six invertibles. The two free orbits $\{i,j,k\}$ and $\{-i,-j,-k\}$ give two simples of dimension three. For a free output orbit, the fusion coefficient is the number of pairs of input group elements with product equal to its chosen representative. For a fixed output, the product fibre is a sum of free C_3-orbits, each contributing its regular character, together with any fixed pairs carrying the product character. These rules independently generate every coefficient. In the order of the three characters over 1, the three over -1, and the two free orbits, the census permutation is $(0,5,6,7,1,2,3,4)$.

The inversion construction for $R8_1_30$ also applies to $A = C_9$ and $A = C_3^2$, producing $R8_1_34$ and $R8_1_35$. Use respectively $\chi(a,b) = \exp(2\pi i ab/9)$ and $\chi(a,b) = \exp(2\pi i(a_1 b_1 + a_2 b_2)/3)$, with

$\tau = 1/3$. Inversion preserves these nondegenerate bicharacters, hence the standard unitary TY associators. The parity coherence is again -1 on m and $+1$ on group elements. There are two equivariant simples above 0, four free pairs of nonzero group elements, and two structures with eigenvalues $\pm i$ above m. Their dimensions are $1, 1, 2, 2, 2, 2, 3, 3$; the same exact character projection generates the recorded tensors.

Finally take $A = \mathbb{F}_2^2$, $\chi(a, b) = (-1)^{a_1 b_2 + a_2 b_1}$ and $\tau = 1/2$. The whole group $S_3 = \mathrm{GL}_2(\mathbb{F}_2)$ preserves χ, so label permutations give a strict unitary action on the standard TY presentation. For $t = 0, 1$, modify its action coherence on m by

$$\alpha_t(g, h) = (-1)^{tp(g)p(h)},$$

where $p : S_3 \to \mathbb{F}_2$ is parity, leaving the group simples unchanged. This is monoidal by the TY grading, and the action pentagon is the elementary normalized 2-cocycle identity for $p(g)p(h)$. Both equivariantizations are therefore unitary. Their orbits $\{0\}$, $A\backslash\{0\}$ and $\{m\}$ have stabilizers S_3, C_2, S_3. Denote their simples by A_ρ, X_ϵ and M_ρ, where ρ ranges over $1, \mathrm{sign}, \mathrm{std}$ and $\epsilon = \pm$. Above m, the equivariance is $i^{tp(g)}\rho(g)$, so the product of two such factors contributes sign^t.

The fusion coefficients are ordinary character inner products. Multiplication by A_ρ is tensoring with ρ, restricted to C_2 on an X-object. The other rules are

$$\begin{aligned} X_\epsilon X_\delta &= A_{\mathrm{Ind}_{C_2}^{S_3}(\epsilon\delta)} + X_+ + X_-, \\ X_\epsilon M_\rho &= M_{(\mathrm{Ind}_{C_2}^{S_3}\epsilon)\rho}, \\ M_\rho M_\sigma &= A_{\rho\sigma\,\mathrm{sign}^t} + X_{\mathrm{Res}_{C_2}^{S_3}(\rho\sigma\,\mathrm{sign}^t)}, \end{aligned}$$

where subscripts extend additively to reducible representations. For XX, the zero fibre is the induced character and the nonzero fibre is a regular C_2 permutation representation. These formulas give $R8_2_611$ at $t = 0$ and $R8_2_154$ at $t = 1$, both of global dimension 48. The model dimensions in the order $(A_1, A_{\mathrm{sign}}, A_{\mathrm{std}}, X_+, X_-, M_1, M_{\mathrm{sign}}, M_{\mathrm{std}})$ are $(1, 1, 2, 3, 3, 2, 2, 4)$. All character arithmetic and tensor comparisons are exact.

One more quadratic model is $R8_1_53$. Yu–Zhang construct a unitary near-group category $NG(G, 16)$ for $G = C_4^2$ [35, Proposition 3.1]. Its bicharacter and quadratic refinement may be taken as

$$\chi(g, h) = i^{g_1 h_1 - g_2 h_2}, \qquad a(g) = \zeta_8^{3(g_1^2 - g_2^2)}.$$

For $H = 2G$, exact finite-group arithmetic gives $H = H^\perp$ and $a(h) = \chi(h, g_a)$ with $g_a = (1, 1) \notin H$. Izumi's general orbifold theorem applies to $H \subseteq H^\perp$ even when $a|_H$ is nontrivial [36, Theorem 12.5]; its crossed-product-factor construction remains unitary. With $A = G/H = C_2^2$, the resulting rules are

$$\alpha_g \alpha_h = \alpha_{g+h}, \qquad \alpha_g \sigma = \sigma \alpha_g, \qquad \sigma^2 = \alpha_{g_a} + \sum_{g \in A} \alpha_g \sigma.$$

There are four invertibles and four noninvertibles of dimension $(8 + 4\sqrt{5})/4 = 2 + \sqrt{5}$, and every noninvertible has a distinct dual. This uses the non-self-dual theorem, rather than its self-dual Lagrangian corollary. In the model order $(00, 01, 10, 11, \sigma, 01\sigma, 10\sigma, 11\sigma)$, the census permutation is $(0, 5, 6, 7, 1, 3, 4, 2)$. The portable verifier checks the subgroup hypotheses, the duality shift and all fusion coefficients; the cited theorem supplies categorical existence.

A cyclic version gives $R8_1_12$. Budinski constructs $NG(C_{16}, 16)$ [37, Theorem 3.1.1 and Table 3.1], with data

$$\chi(x, y) = \exp(2\pi i\, 3xy/16), \qquad a(x) = \exp(-\pi i\, 3x^2/16).$$

Although the table prints decimal arguments, the thesis's Section 3.3.2 gives an exact algebraic certification, with the defining-polynomial scope explicitly including $n = 16$. We use that exact existence statement as a citation dependency; no numerical reconstruction is counted here as a

category. For $H = 4C_{16}$, one has $H = H^{\perp}$ and $a(h) = \chi(h, 2)$. The same unitary orbifold theorem [36, Theorem 12.5] therefore gives

$$g^4 = 1, \qquad \sigma g = g^{-1}\sigma, \qquad \sigma^2 = g^2 + (1 + g + g^2 + g^3)\sigma.$$

In the order $(1, g, g^2, g^3, \sigma, g\sigma, g^2\sigma, g^3\sigma)$, the census permutation is $(0, 1, 7, 2, 3, 6, 4, 5)$. The four noninvertibles are paired by duality, distinguishing this ring from the standard C_4 Haagerup–Izumi ring. The finite verifier checks all subgroup hypotheses and the full tensor exactly.

The ring $R8_1_83$ has a triality-gauging model. The S_3 gauging of the three-fermion category exists by [2, Section 6.2]; the unobstructed gauging data form the usual $H^3(S_3, U(1))$ torsor, including the twist denoted $m = 0$ in [38, Section VI.D and Table IX]. This supplies an actual category. Starting from a pointed category, gauging produces a weakly group-theoretical fusion category, hence one admitting a unitary realization [39, Theorem 5.20]. Its eight integral simples, with dimensions $(1, 1, 2, 3, 3, 4, 4, 4)$, form a fusion subcategory of dimension 72.

The integer fusion formulas for this subcategory are independently checked against the full twelve-object S-matrix of [38, Eq. (6.53), $m = 0$]. Its entries lie in $\mathbb{Q}(\sqrt{2})$; using the unnormalized matrix, the verifier checks $S^2 = 144I$, all 1728 nonnegative integral Verlinde coefficients, and fusion closure of the eight integral simples. It then compares the entire tensor with the pinned census under the stored permutation. Category existence comes from the gauging construction, not merely from the feasible modular datum.

The table provides one construction per positive tensor. Some have two isomorphic presentations; `data/realized_models.json` records every matched constructor, exact model tensor, precise model source and source-to-model isomorphism. Labels such as "pseudo-$PSU(2)_6$" denote the conventional fusion-ring name, not a weaker existence assertion: the cited rank-four table supplies a unitary category [14]. The table exhausts both positive lists: 48 tensors at exact multiplicity one and 29 at exact multiplicity two.

7.2. **Pointed equivariantizations and an affine group.** Four further models are strict unitary equivariantizations of untwisted pointed categories. Their group automorphisms are

$$\begin{aligned}
&R8_2_244: && A = C_4^2, && (x, y) \mapsto (-y, x - y), && G = C_3;\\
&R8_2_289: && A = C_2^4, && (x, y, z, w) \mapsto (y, x + y, w, z + w), && G = C_3;\\
&R8_2_143: && A = C_{17}, && x \mapsto 4x, && G = C_4;\\
&R8_2_179: && A = C_2^4, && (x, y, z, w) \mapsto (w, x + w, y + w, z + w), && G = C_5.
\end{aligned}$$

In each case the displayed generator has the stated order and acts freely off zero. The zero orbit supplies the ordinary characters of G, and each nonzero orbit has trivial stabilizer. Tensor products are calculated by orbit addition and projection to stabilizer characters, giving the exact full source tensors. Strict permutation functors with identity tensorators supply coherent unitary actions.

The ring $R8_2_237$ is $K_0(\mathrm{Rep}(A\Gamma L(1, 8)))$. This group of order 168 has elements (v, a, t) acting on $\mathbb{F}_8$ as $x \mapsto v + ax^{2^t}$. Its simple dimensions are $(1^3, 3^2, 7^3)$. The three dimension-seven characters have the exact expression

$$\chi_s(v, a, t) = \zeta_3^{st} \sum_{\substack{r \in \mathbb{F}_8^{\times} \\ ar^{2^t - 1} = 1}} (-1)^{\mathrm{Tr}_{\mathbb{F}_8/\mathbb{F}_2}(v/r)}.$$

Together with the characters inflated from $\mathbb{F}_8^{\times} \rtimes C_3$, they give an exact character table in $\mathbb{Q}(\zeta_{21})$. Orthogonality and all tensor multiplicities are checked independently by finite group sums.

7.3. **Bosonic quotients and the D_{14} model.** The ring $R8_2_557$ is the central bosonic quotient of $PSU(2)_6 \boxtimes PSU(2)_6$ by its diagonal invertible. Each factor invertible is a fermion, so their product has self-braiding $+1$ and trivial pointed associator. It generates a genuine central $\mathrm{Rep}(C_2)$. All sixteen simples have free diagonal orbits; the eight free-module orbits give the stated ring.

For $R8_2_313$, use instead the two unitary pseudo-$PSU(2)_6$ categories of [30], with the exact sourced associators discussed above. Their central invertibles have self-braidings $+i$ and $-i$, and each has associator $F_{ggg} = -1$. The diagonal product therefore has both trivial self-braiding and trivial associator; its central $\mathrm{Rep}(C_2)$ gives the same eight-orbit quotient construction. For each rank-four factor the universal grading is trivial, so a central lift of an invertible is unique; its central square is the unit. This verifies the order-two condition as well as the associator and self-braiding conditions required for the categorical quotient.

The ring $R8_2_855$ is the even D_{14} category, or the local module category of the $SU(2)_{24}$ algebra $1 \oplus V_{24}$. It is unitary modular [48, 49]. Its basis is $f_0, f_2, f_4, f_6, f_8, f_{10}, P, Q$, with P, Q self-dual. Fold the bulk $SU(2)_{24}$ products by $f_{2t} = f_{2(12-t)}$ and $f_{12} = P + Q$. The remaining rules are

$$Pf_{2j} = \sum_{k=6-j}^{5} f_{2k} + \begin{cases} P & j \text{ even}, \\ Q & j \text{ odd}, \end{cases} \qquad Qf_{2j} \text{ interchanges } P, Q,$$

$$P^2 = P + f_0 + f_4 + f_8, \quad Q^2 = Q + f_0 + f_4 + f_8, \quad PQ = f_2 + f_6 + f_{10}.$$

All these constructions preserve the unitary structure.

7.4. **A Morita construction over C_7.** The ring $R8_2_239$ is a Morita dual of an S_3 extension of Vec_{C_7}. In $\mathrm{BrPic}(\mathrm{Vec}_{C_7}) = O(C_7 \oplus \widehat{C_7})$ take $r = \mathrm{diag}(2, 4)$ and let s exchange the two coordinates. They generate S_3. Its degree-three obstruction vanishes by coprime cohomology with seven-primary coefficients; the next obstruction lies in $H^4(S_3, \mathbb{C}^\times) = 0$. Thus the actual extension exists [7, Theorem 1.3]. Its C_3 restriction is pointed, on $F_{21} = C_7 \rtimes C_3$. The associator class there is inflated from C_3: the order-three action kills the $H^3(C_7, \mathbb{C}^\times)$ invariant part. Surjectivity of $H^3(S_3, \mathbb{C}^\times) \to H^3(C_3, \mathbb{C}^\times)$ permits an outer twist that makes this restriction untwisted.

Take the separable algebra $B = \bigoplus_{h \in C_3} h$ in this restriction. The category of B-bimodules is an actual Morita dual, whose neutral category is $\mathrm{Rep}(F_{21})$. Write its simples as α_i ($i \in \mathbb{Z}/3$), X, Y, and odd m_j ($j \in \mathbb{Z}/3$). The nontrivial rules, with all m_j self-dual, are

$$X^2 = X + 2Y, \quad Y^2 = 2X + Y, \quad XY = \sum_i \alpha_i + X + Y,$$

$$\alpha_i m_j = m_{i+j}, \quad m_j \alpha_i = m_{j-i}, \qquad Xm_j = Ym_j = m_j X = m_j Y = \sum_k m_k,$$

$$m_j m_k = \alpha_{j-k} + X + Y.$$

They follow from the S_3 bimodule action and reciprocity, and match all 512 source coefficients. The extension and its Morita dual are weakly group-theoretical, hence unitarizable [39, Theorem 5.20].

7.5. **Charge-conjugation defects of $PSU(3)_4$.** The ring $R8_2_290$ is a unitary C_2 crossed extension of the rank-five modular category $\mathcal{B} = PSU(3)_4$. Charge conjugation is an actual braided symmetry [50, Corollary 2.7]. The invertible group of $\mathcal{B}$ is trivial, so the degree-three gauging obstruction vanishes, and $H^4(C_2, U(1)) = 0$ removes the remaining obstruction. The unitary gauging theorem [2, Sections 3.2–3.4] therefore supplies an actual unitary extension. Its odd rank is three, the number of charge-conjugation-fixed neutral simples [51, Proposition 1.1].

For completeness, its tensor is forced by a small exact module calculation. In the neutral basis $(0,0), (3,0), (1,1), (0,3), (2,2)$ the nonselfdual generator X has regular characteristic polynomial $(t^2 + t + 2)(t^3 - 2t^2 - t + 1)$, computed by quantum Weyl alternants in $\mathbb{Q}(\zeta_{21})$. Its odd action A has the cubic factor as characteristic polynomial, since its Perron root has degree three. The commuting neutral action and the dual-transpose relation make A normal. Its three roots are real, so A is symmetric. Nonnegative integer entries, $\mathrm{Tr}(A) = 2$ and $\mathrm{Tr}(A^2) = 6$ give, up to relabelling,

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}.$$

The five neutral actions are I, A, A^2-I, A, A^2-A. Distinct odd dimensions force each odd object to be self-dual; reciprocity then determines every odd product. The independent constructor matches all 512 census coefficients.

7.6. **A self-dual Hopf extension in dimension thirty-two.** For $R8_2_575$ the neutral Hopf algebra is $H = H_{C:\sigma_1}$ of dimension sixteen from [44]. A symmetric nondegenerate Hopf pairing supplies the extra data of the Davydov–Runkel construction [52, Theorem 2.5]. Here is enough explicit information to fix the construction and its normalization. Write $h(p,q,j)$ for its basis, with p modulo four and q, j modulo two. Put $\zeta = \exp(2\pi i/16)$ and $i = \zeta^4$. The pairing is

$$P(h(p,q,j), h(r,s,k)) = \delta_{s,j}\delta_{k,q}\, i^{-3rp}\zeta^{jq}/4.$$

The supplementary Hopf presentation and checker verify nondegeneracy, symmetry and every pairing identity exactly in $\mathbb{Q}(\zeta)$. Davydov–Runkel use $H^\vee = (H^*)^{\mathrm{op,cop}}$; consequently their self-duality map is $\Gamma = S \circ P^{-1}$, rather than the unmodified inverse pairing. The right cointegral $\lambda(h(p,q,j)) = \delta_{j,0}/2$ satisfies $\Gamma(\lambda) = 4\Lambda$ and $\lambda(\Gamma(\lambda)) = 1$, where Λ is the normalized integral. The distinguished grouplike is one, $S^2 = 1$, and Γ is symmetric, so all conditions of the cited theorem hold. Its rigidity criterion [52, Proposition 3.25] gives an actual fusion category $\mathrm{Rep}(H) \oplus \mathrm{Vec}$. Its neutral ring is $\mathrm{Rep}(D_{16})$; for the odd object M of dimension four,

$$XM = MX = \mathrm{FPdim}(X)M, \qquad M^2 = \sum_{X \in \mathrm{Irr}(H)} \mathrm{FPdim}(X)X.$$

The dimension-32 solvability theorem and [39, Theorem 5.20] make this fusion category unitarizable. The complete model tensor, Hopf identities, and source comparison are separate exact checks.

7.7. **The dual Haagerup–Izumi C_7 category.** The ring $R8_2_868$ is the dual even part of the unitary Haagerup–Izumi C_7 subfactor. Its existence and center are supplied by [53, Theorem 5(d)]. Put $d = (7+\sqrt{53})/2$, so $d^2 = 1 + 7d$. Izumi's dual Q-system analysis [8, Proposition 7.4 and Theorem 7.5] gives rank eight, with dimensions $1, d, (d-1)^3, (d+1)^3$.

We verify the canonical-algebra hypothesis needed to use the full fusion rules in [53, Theorem 7]; matching modular data alone would not suffice. In its published double, the twist-one simples are 1, b of dimension d^2, and six A_i of dimension $1+d^2$. The nonzero C_{53} sector has no twist one, and the other twist-one labels are precisely those with zero second coordinate. For the actual dual category $\mathcal{D}$, its canonical Lagrangian algebra $L = I_{\mathcal{D}}(1)$ is connected, has trivial twist and dimension $7(1+d^2)$. Therefore

$$L = 1 + m_b b + \sum_{i=1}^{6} m_i A_i.$$

Writing $T = \sum_i m_i$, rational and irrational coefficients in the dimension identity give $m_b + T = 7$ and $m_b + 2T = 13$, hence $m_b = 1, T = 6$. Adjunction gives $\dim \mathrm{End}(L) = \mathrm{rank}(\mathcal{D}) = 8$, so $\sum_i m_i^2 = 6$. Thus every $m_i = 1$. The actual canonical algebra has exactly the underlying object required by the cited theorem; no assumption about a guessed algebra multiplication is made.

For a direct exact constructor, use basis $1, a, b_1, b_2, b_3, c_1, c_2, c_3$. Put $B = \sum_i b_i$, $C = \sum_i c_i$, $b_0 = c_0 = 0$, and $b_{-i} = b_i, c_{-i} = c_i$ for indices modulo seven. The rules are

$$\begin{aligned}
a^2 &= 1 + a + B + C, & ab_i &= a + B - b_i + C,\\
ac_i &= a + B + c_i + C, & b_i c_j &= a + B + C,\\
b_i b_j &= \delta_{ij} 1 + (1-\delta_{ij})a + B - b_{i+j} - b_{i-j} + C,\\
c_i c_j &= \delta_{ij} 1 + (1+\delta_{ij})a + B + c_{i+j} + c_{i-j} + C.
\end{aligned}$$

They match every source coefficient; all 4096 associativity equations and all dimension products in $\mathbb{Q}(d)$ are checked. Both even parts of the cited finite-depth C^* subfactor are unitary.

Model	Model-to-census permutation
$R8_2_143$	$(0,1,7,2,3,5,6,4)$
$R8_2_179$	$(0,1,4,3,2,5,6,7)$
$R8_2_237$	$(0,1,2,5,6,7,4,3)$
$R8_2_239$	$(0,1,2,3,4,5,7,6)$
$R8_2_244$	$(0,1,2,3,7,4,5,6)$
$R8_2_289$	$(0,1,2,3,4,5,6,7)$
$R8_2_290$	$(0,1,6,2,7,3,4,5)$
$R8_2_313$	$(0,7,1,2,3,4,6,5)$
$R8_2_557$	$(0,1,2,7,3,6,5,4)$
$R8_2_575$	$(0,5,1,2,3,7,4,6)$
$R8_2_855$	$(0,1,7,4,5,6,2,3)$
$R8_2_868$	$(0,3,1,2,4,5,7,6)$

The model bases are fixed by the independent constructors in the supplement. Each permutation is checked on the entire ordered tensor; it is not inferred from a character table or dimension vector.

8. A nonsplit extension of the 2D2 category

Proposition 8.1. *The ring $R8_2_37$ admits a unitary categorification.*

Proof. We use the factor realization of a graded extension of the known Haagerup–Izumi category for C_4 in [66, Section 3 and Theorem 4.1]. It supplies automorphisms α_j, β of a factor M, an endomorphism ρ, and unitaries $u, v \in M$ such that

$$\beta\alpha_j = \alpha_j\beta, \qquad \beta\rho = \mathrm{Ad}(u)\alpha_1\rho\beta, \qquad \beta^2 = \mathrm{Ad}(v)\alpha_{1+z}, \quad z \in \{0,2\}.$$

The α_j form an honest C_4 action. The character identities in [66, Lemma 3.2, (3.10) and (3.18)] give $\alpha_2(u) = -u$ and $\alpha_2(v) = -v$.

Form $P = M \rtimes_{\alpha_2} C_2$, with implementing involution λ. Grossman's construction [55, Section 4.1] extends the neutral generators by

$$\widetilde{\alpha}_j(\lambda) = (-1)^j\lambda, \qquad \widetilde{\rho}(\lambda) = \lambda.$$

They generate the unitary $2D2$ category with simple sectors $(1, g, r, gr)$, where $g = [\widetilde{\alpha}_1]$, $r = [\widetilde{\rho}]$, and

$$g^2 = 1, \qquad gr = rg, \qquad r^2 = 1 + 2r + 2gr.$$

This construction depends only on the algebra $1 + \alpha_2$ in the neutral category, so it applies to the neutral realization just chosen.

Extend β to $b \in \mathrm{Aut}(P)$ by $b(\lambda) = \lambda$. The factor identities extend to

$$b\widetilde{\rho} = \mathrm{Ad}(u)\widetilde{\alpha}_1\widetilde{\rho}b, \qquad b^2 = \mathrm{Ad}(v)\widetilde{\alpha}_{1+z}.$$

Indeed, they hold on M by construction. On λ the two right-hand sides are respectively $u(-\lambda)u^* = \lambda$ and $v(-\lambda)v^* = \lambda$. The anticommutation of u, v with λ is essential here.

Put $a = [b]$. These identities imply

$$a^2 = g, \qquad a^4 = 1, \qquad ara^{-1} = gr = a^2r, \qquad ra = a^{-1}r.$$

The full tensor category generated by b and $\widetilde{\rho}$ has exactly the eight simple sectors

$$(1, a, a^2, a^3, r, ar, a^2r, a^3r).$$

The four invertibles are distinct because $a^2 = g \neq 1$. If $a^ir \cong a^jr$, Frobenius reciprocity puts a^{j-i} in $rr^* = 1 + 2r + 2gr$, so $i = j$. The noninvertibles have dimension $2 + \sqrt{5}$. The displayed relations show that these eight sectors are closed under products and duals, and their associativity and unitarity are inherited from $\mathrm{End}_0(P)$.

For $i, j \in \mathbb{Z}/4$, their multiplication is

$$a^i a^j = a^{i+j}, \qquad a^i(a^j r) = a^{i+j} r,$$
$$(a^i r)a^j = a^{i-j} r, \quad (a^i r)(a^j r) = a^{i-j} + 2a^{i-j} r + 2a^{i-j+2} r.$$

The displayed basis maps to zero-based census labels $(0, 1, 7, 2, 5, 3, 6, 4)$. The exact verifier compares all 512 structure constants under this bijection. Thus it identifies an already constructed category with $R8_2_37$; no coherent order-two autoequivalence is assumed. □

8.1. **Odd isotypic multiplicities and involution algebras.**

Lemma 8.2. *Let X be a self-dual simple in a pivotal fusion category over $\mathbb{C}$. If a self-dual simple S occurs with odd multiplicity in $X^{\otimes n}$, then*

$$\nu_2(S) = \nu_2(X)^n.$$

More generally, in a palindromic tensor word of self-dual simples, every self-dual constituent of odd multiplicity has indicator equal to the product of the indicators of the factors.

Proof. Use pivotal coherence to suppress associators and double-dual identifications. Choose a self-duality $f : X \to X^*$ whose pivotal transpose is $t(f) = \epsilon f$, where $\epsilon = \nu_2(X) \in \{1, -1\}$. The reversed dual word has the same identical factors, so

$$F = f^{\otimes n} : X^{\otimes n} \longrightarrow (X^{\otimes n})^*, \qquad t(F) = \epsilon^n F.$$

This construction needs no braiding: reversal changes neither the object word nor the sequence of component morphisms. On the S-isotypic block the nondegenerate form is $f_S \otimes B$, with $t(f_S) = \nu_2(S) f_S$ and B an invertible bilinear-form matrix on the complex multiplicity space. Hence

$$B^{\mathsf{T}} = \frac{\epsilon^n}{\nu_2(S)} B.$$

An invertible alternating matrix has even size, which proves the assertion for an odd multiplicity. Semisimplicity ensures that the form on a self-dual isotypic block is nondegenerate. The same calculation works for a palindrome, using the same chosen self-duality at each pair of matching positions. □

Proposition 8.3. *The ring $R8_1_49$ has no categorification over $\mathbb{C}$.*

Proof. Its invertibles are $G = \{0, 3, 4, 5\} \cong C_2^2$, and its four noninvertibles have dimension $d = 2 + \sqrt{5}$, with $d^2 = 1 + 4d$. The published two-dimension theorem [25, Lemma 3.1] therefore replaces any hypothetical categorification by a pseudo-unitary Galois conjugate with the same fusion rules. Equip this conjugate with its canonical positive spherical structure. This supplies the pivotal structure without assuming one on the initial category.

Set $u = 3$, $g = 5$ and $\rho = 6$. The source tensor gives

$$\rho^* = \rho, \qquad \rho^2 = 1 + \sum_{h \in G} h\rho, \qquad \rho^3 = (1, 4, 4, 1, 1, 1, 5, 4)$$

in source-label order. The unit and every invertible occur once in the cube. Lemma 8.2 first gives $\nu_2(\rho) = 1$ and then $\nu_2(h) = 1$ for every $h \in G$. In the positive spherical structure of a pointed C_2 category, the indicator of its nontrivial object is the sign of its associator class. Thus the restriction to $H = \{1, u\}$ is trivial, and its regular algebra $A = 1 \oplus u$ actually exists and is separable. No central lift of u is required.

Consider $\mathcal{B} = {}_A\mathcal{C}_A$. The source relation $u\rho u = g\rho$ gives two pointed H-by-H orbits of size two, $\{0, 3\}$ and $\{4, 5\}$, and the free noninvertible orbit $\{1, 2, 6, 7\}$. Each pointed orbit has stabilizer C_2 and contributes two simple bimodules of dimension one, since every complex projective representation of C_2 can be linearized. The free orbit contributes the single simple $A\rho A$ of bimodule dimension

$4d/2 = 2d$. Thus $\mathcal{B}$ has four invertibles, forming $G' \cong C_4$ or C_2^2, and one noninvertible Z. Reciprocity and dimensions force

$$Z^2 = \sum_{h \in G'} h + 8Z.$$

For $G' = C_2^2$ this contradicts [25, Theorem 1.1]; for $G' = C_4$ it contradicts the general cyclic-four criterion [16, Theorem 6.3] at $k = 2$. The source-bound verifier `verify_r49_fs_morita.py` checks the cube, all double orbits, and the same six-branch terminal certificate used below. Hence the original ring cannot be categorified. □

8.2. **An inverse-Morita obstruction for the split involution.**

Proposition 8.4. *The ring R8_2_170 admits no fusion-category categorification over* $\mathbb{C}$.

Proof. Its source tensor has a faithful C_2 grading with neutral labels $(0, 5, 6, 7)$. Put $g = 5$, $r = 6$, and $u = 3$. Its four invertibles form C_2^2, with u odd, and $uru = gr$. The subgroup $H = \{1, u\}$ maps isomorphically to the grading group. If a category with this tensor exists, multiply its associator by the inflation of the inverse of its pointed associator class on H. This is a normalized grading three-cocycle, so it preserves the pentagon and every fusion coefficient. After this twist $A = 1 \oplus u$ is a connected separable algebra. This step does not assume that A exists before twisting.

Consider the fusion category $\mathcal{B} = {}_A\mathcal{C}_A$. For a free bimodule $F(Y) = A \otimes Y \otimes A$, adjunction gives

$$\dim \mathrm{End}_{\mathcal{B}}(F(Y)) = \dim \mathrm{Hom}_{\mathcal{C}}(Y, A \otimes Y \otimes A) = |\{(h, k) \in H^2 : hYk \cong Y\}|.$$

The two pointed double orbits are $\{0, 3\}$ and $\{4, 5\}$, each of stabilizer size two. Each free bimodule therefore splits into two distinct simple bimodules, once each: its semisimple endomorphism algebra has dimension two. The underlying support of either summand is stable under H on both sides, so each contains both orbit objects once. Dividing its underlying FP dimension by $\mathrm{FPdim}(A) = 2$ shows that it is invertible. Thus these orbits give four invertibles in $\mathcal{B}$.

The four noninvertibles form one double orbit $\{1, 2, 6, 7\}$ with trivial stabilizer. Hence $F(r)$ is a single simple Z of dimension $2(2 + \sqrt{5}) = 4 + 2\sqrt{5}$. Every simple bimodule occurs in a free one, so this is the complete list. The four invertibles form a group G equal to C_4 or C_2^2. The unique noninvertible is self-dual and fixed by G; reciprocity and dimensions give

$$Z^2 = \sum_{h \in G} h + 8Z.$$

For $G = C_2^2$, this contradicts [25, Theorem 1.1]. For $G = C_4$, it contradicts the general cyclic-four central-support criterion of [16, Theorem 6.3], at parameter $k = 2$ and moment height six. The archived optimizer-free replay checks all six branches: two force a fusion coefficient $1/2$, and four have exact separating vectors with nonnegative $A^{\mathsf{T}}y$ and respective values $b^{\mathsf{T}}y = -800, -220, -320, -80$. This terminal criterion applies to arbitrary complex categorifications through its proved Galois pseudo-unitarity reduction. Both possible groups are excluded. No classification of the neutral $2D2$ category is used. □

9. The cyclic order-six Haagerup–Izumi algebra dual

Proposition 9.1. *The dual of the algebra* $1 + \rho$ *in a known unitary Haagerup–Izumi category for* C_6 *has Grothendieck ring R8_2_535.*

Proof. Set $d = 3 + \sqrt{10}$ and $D = 6(1 + d^2) = 120 + 36\sqrt{10}$. The category and its algebra $1 + \rho$ exist by [67, Example 4.5 and condition (2.24)]. Denote the unitary dual by $\mathcal{E}$. Izumi's description [8, Proposition 7.4 and Theorem 7.5] gives its simple objects and dimensions as

	1	b	r	rb	e_1	e_2	s_1	s_2
FPdim	1	1	d	d	$d+1$	$d+1$	$d-1$	$d-1.$

Here $b = \beta_3$, $e_j = \eta_j$, and s_1, s_2 are the two σ objects. Its canonical module has seven simples $(i_0, \dots, i_5, \tau)$ with relative dimensions $(1, \dots, 1, d-1)$.

Let P be the cyclic permutation matrix on the six i-labels. The upper-left six-by-six blocks of the eight action matrices are

$$I, \quad P^3, \quad I, \quad P^3, \quad P + P^{-1}, \quad P^2 + P^{-2}, \quad 0, \quad 0.$$

The matrices of $1, b$ have zero off-diagonal τ entries and τ diagonal 1. Each of the other six has every off-diagonal τ entry equal to 1, with respective τ diagonals $(5, 5, 6, 6, 4, 4)$. These formulas follow from the decompositions in [8, Proposition 7.4]; the last entries also follow from the relative FP dimension vector.

These matrices generate a commutative algebra of dimension five. Their three independent dependencies are

$$M_1 - M_b - M_r + M_{rb} = 0, \qquad M_1 - M_r + M_{s_1} = 0, \qquad M_1 - M_r + M_{s_2} = 0.$$

The complex fusion algebra is semisimple. A noncommutative simple block would have dimension at least four and would have to lie in the three-dimensional kernel of this representation. Hence the entire fusion algebra of $\mathcal{E}$ is commutative.

Write $d' = 3 - \sqrt{10}$ and $D' = 120 - 36\sqrt{10}$. Fourier diagonalization of the module yields five characters:

	1	b	r	rb	e_1	e_2	s_1	s_2	f
χ_{FP}	1	1	d	d	$d+1$	$d+1$	$d-1$	$d-1$	D
χ'_{FP}	1	1	d'	d'	$d'+1$	$d'+1$	$d'-1$	$d'-1$	D'
u	1	-1	1	-1	1	-1	0	0	6
v	1	1	1	1	-1	-1	0	0	6
w	1	-1	1	-1	-2	2	0	0	12.

There are three further one-dimensional characters. Orthogonality to the displayed rows forces each to have the form

$$(1, \epsilon, -1, -\epsilon, 0, 0, s, t), \qquad \epsilon \in \{1, -1\}, \quad s + t = 1 + \epsilon. \tag{1}$$

We determine their codegrees using the center of the known category, which is Morita invariant. The complete published inventory [67, Lemma 3.5 and Theorem 3.9] is

simple dimension	1	d^2	$1+3d$	$2+6d$	$3d$	$6d$
multiplicity	2	2	4	16	4	18.

The dimensions squared sum to D^2. Formal-codegree induction [24] associates to each character a simple constituent of $I(1)$ of dimension D/f. Thus the possible codegrees are

$$D, \quad D', \quad 12, \quad 6, \quad 4\sqrt{10}, \quad 2\sqrt{10}.$$

Formal codegrees are totally positive: in this commutative case each is the squared norm of a character and every Galois conjugate is another formal codegree. This eliminates the last two values. For a missing character, the Perron bound gives

$$f = 4 + |s|^2 + |t|^2 \le 4 + 2(d-1)^2 = 8 + 8d < D.$$

So no missing codegree is D. The five known rows form a Galois-stable set; a missing codegree D' would have a missing Galois conjugate of codegree D, also impossible. Finally $\sum_\chi f_\chi^{-1} = 1$ and $D^{-1} + (D')^{-1} = 1/6$ force the three missing codegrees to be $6, 6, 12$.

The object b permutes s_1, s_2. If it fixed both, a missing character with $\epsilon = -1$ would have $s = t = 0$ and codegree 4. All three missing rows would then have $\epsilon = 1$, but the vectors in (1) with $\epsilon = 1$ span a space of dimension at most two. Therefore b exchanges s_1 and s_2, and $t = \epsilon s$. There is exactly one missing row with $\epsilon = 1$, namely $(1, 1, -1, -1, 0, 0, 1, 1)$ of codegree 6. The two remaining rows have $\epsilon = -1$, one of each codegree $6, 12$. Complex conjugation preserves the missing set, ϵ, and

the codegree, so each of these two rows is real. Their norms and orthogonality determine, up to exchanging s_1, s_2, the full list

$$\begin{array}{c|cccccccc|c} x & 1 & 1 & -1 & -1 & 0 & 0 & 1 & 1 & 6 \\ y & 1 & -1 & -1 & 1 & 0 & 0 & 1 & -1 & 6 \\ z & 1 & -1 & -1 & 1 & 0 & 0 & -2 & 2 & 12. \end{array}$$

All characters are real, hence all simple objects are self-dual. Character inversion now determines every structure constant:

$$N_{ij}^{k} = \sum_{\chi} \frac{\chi(i)\chi(j)\chi(k)}{f_\chi}.$$

The resulting tensor is identified with $R8_2_535$ by the basis map $(0, 1, 3, 6, 4, 5, 2, 7)$. The verifier reconstructs all 512 coefficients exactly and checks the canonical module equations. No multiplicity bound or completeness claim about the census is used to determine the unknown dual ring. □

10. Algebraic rigidity of the 2D2 module graph

The next argument gives four general complex obstructions. It does not assert that every categorification of the rank-four $2D2$ ring is unitary: the specific module graph is part of the hypothesis.

Lemma 10.1. *Let $\mathcal{D}$ be a pseudo-unitary fusion category and $\mathcal{M}$ an indecomposable module category with a simple object X. Suppose its rooted principal graph is $2D2$, with modulus $\delta = \sqrt{3+\sqrt{5}}$. Then the associated algebraic shaded planar algebra is isomorphic to the known unitary $2D2$ planar algebra. In particular its dual even category has rank six.*

Proof. We first justify the algebraic planar-algebra construction. Schaumann's module-trace theorem [68, Proposition 5.8] supplies positive FP trace weights on $\mathcal{M}$. His Theorem 4.5 constructs natural double-adjoint isomorphisms for module functors between any two traced module categories, compatible with composition. Applied to $\mathcal{D}$ and $\mathcal{M}$, it equips the full Morita linking category with a pivotal structure. Corollary 4.6 and Proposition 5.10(iv) give the compatible spherical dual. Normalize the module trace so that the two loop values of X are δ.

For completeness, the graph embedding in this setting requires no star operation. Act on a column of the linking category, and decompose every object into simples. A morphism acts on the resulting path spaces as a direct sum of matrices, faithfully. Choose dual bases for the nondegenerate trace pairings. Composition is matrix multiplication, while evaluation and coevaluation are the weighted graph cups and caps. Thus all planar operations are preserved. This is the algebraic pivotal construction of [43, Theorems 3.45 and 3.70], with the cup and cap calculation of its Theorem 3.75 performed with dual bases. Its dagger-specific Theorem 3.77 is not needed. Denote the resulting planar algebra by $\mathcal{P}$ and its embedding into the graph planar algebra by $\mathcal{P} \subseteq \mathcal{G}$.

The graph is two-supertransitive. Its depth-three Jones–Wenzl corner is $f_3\mathcal{P}_{3,+}f_3 = \mathbb{C}p \oplus \mathbb{C}q$, where p, q are orthogonal primitive idempotents and $p + q = f_3$. The two depth-three PF weights coincide, so $T = p - q$ has trace zero and $T^2 = f_3$. The bilinear trace pairing is nondegenerate, and

$$\mathcal{P}_{3,+} = \mathrm{TL}_{3,+} \oplus \mathbb{C}T$$

is a trace-orthogonal decomposition. Pairing a cap of T with every element of $\mathcal{P}_{2,+}$ shows that it vanishes, since the corresponding cups lie in $\mathrm{TL}_{3,+}$. Rotation preserves the one-dimensional orthogonal complement, so T is an uncappable rotational eigenvector, whose eigenvalue is a cube root of unity. We verify the exclusion of the two primitive roots directly below; this also independently replays [69, Lemma 3.2] without a self-adjointness hypothesis.

We describe the exact complex calculation needed at this point. Label the graph vertices $0, \ldots, 7$, with edges

$$01,\ 12,\ 23,\ 24,\ 35,\ 45,\ 56,\ 67,$$

rooted at 0 and with even vertices $0, 2, 5, 7$. Put $s = \sqrt{5}$, $\varphi = (1+s)/2$, and assign weights

$$(w_0, \dots, w_7) = (1, 1, 2+s, \varphi, \varphi, 2+s, 1, 1).$$

These are the PF weights with odd entries divided by δ. For a path $p = (p_0, p_1, p_2, p_3)$, set $A_p = \sqrt{w_{p_1} w_{p_2}}$. The rectangular lopsided matrix is $T_{\mathrm{rect}} = A T_{\mathrm{sph}} A^{-1}$. Thus the coefficient at a loop $(v_0, \dots, v_5)$ is multiplied by $\sqrt{w_{v_1} w_{v_2} / (w_{v_5} w_{v_4})}$. In these coordinates a Temperley–Lieb projection has entries $w_{p_i} / \sum_{a \sim p_{i-1}} w_a$ on paths differing only at the backtracking vertex p_i, and zero otherwise. Caps therefore give unweighted sum equations. Two-click rotation replaces $(v_0, \dots, v_5)$ by $(v_4, v_5, v_0, v_1, v_2, v_3)$, with coefficient $w_{v_0} w_{v_5} / (w_{v_2} w_{v_1})$.

For a primitive root ω, work in $\mathbb{Q}(s, \omega)$, $\omega^2 + \omega + 1 = 0$. The cap and ω-rotation equations on the 152 loops have rank 144. After their elimination, let E_j be the 90 distinct normalized nonzero entries of $T^2 - f_3$, in the deterministic path-matrix order of the verifier. Direct polynomial arithmetic gives

$$\begin{aligned} 1 = {} & (-11 - 5s) E_{26} + \left(-25 - 11s + \frac{5 + 3s}{2} \omega \right) E_0 \\ & + \left(-12 - 6s + \frac{9 + 3s}{2} \omega \right) E_1 + (44 + 20s) E_4 \\ & + \big(4 + 2s - (7 + 3s) \omega \big) E_{13}. \end{aligned}$$

The portable checker reconstructs the linear elimination and every E_j from the graph before checking this identity. It excludes both primitive cubic roots simultaneously, without complex conjugation equations. Hence the eigenvalue is one.

For eigenvalue one the cap and rotation system again has rank 144. Its eight free loop coordinates $x_0, \dots, x_7$ may be chosen, in order, at

$$012321,\ 212323,\ 212354,\ 212453,\ 232323,\ 232354,\ 232453,\ 235453.$$

Put $k = (3 - s)/2$ and $c = (5s - 11)/2 > 0$. On this eight-dimensional linear space, the ideal of all path-matrix entries of $T^2 - f_3$ is generated by

$$\begin{gathered} x_0 + \frac{3 + s}{2} x_7, \qquad x_1 + \frac{3 - s}{2} x_7, \qquad x_2 + 2x_5, \qquad x_3 + 2x_6, \\ x_4 + \frac{s - 1}{2} x_7, \qquad x_5 x_6 - c, \qquad x_7^2 - k^2. \end{gathered} \tag{2}$$

This is an equality of polynomial ideals over $\mathbb{Q}(\sqrt{5})$, not just equality of their real or self-adjoint solution sets. The portable certificate generates all square equations from the graph and verifies both inclusions: reduction proves that (2) implies every square equation, and explicit polynomial identities of degree at most three express its seven generators in the original equations. The linear rank and all coefficients are checked exactly.

Thus every complex solution is $\pm T(\lambda)$ for $\lambda \in \mathbb{C}^\times$, with free coordinates

$$\left(-1, -k^2, -2\sqrt{c}\lambda, -2\sqrt{c}\lambda^{-1}, -\frac{s - 1}{2} k, \sqrt{c}\lambda, \sqrt{c}\lambda^{-1}, k \right).$$

The base solution $T(1)$ is self-adjoint after the diagonal change to spherical coordinates. Assign an oriented-edge character λ to the edge $3 \to 5$, its inverse to the reverse edge, and 1 to all others. Multiplying each loop coefficient by its edge-character product is a graph planar algebra automorphism: the factors on internal edges cancel in every state sum. The eight coordinates have winding numbers $(0, 0, 1, -1, 0, 1, -1, 0)$, so this gauge carries $T(1)$ to $T(\lambda)$. The two signs generate the same subalgebra.

After applying this gauge, $\mathcal{P}$ contains the planar subalgebra generated by $T(1)$. By [69, Lemma 3.4 and Theorems 7.8–7.9], this is the known unitary $2D2$ planar algebra. Its positive box dimensions are the rooted-loop dimensions of $2D2$, exactly those of $\mathcal{P}$. The containment is therefore equality in every positive box space; one-click rotation gives equality in the negative spaces of positive degree,

and both zero-box spaces are $\mathbb{C}$. This rules out additional higher-depth data without using positivity on $\mathcal{P}$. The known dual graph has level sizes $(1,1,1,2,4,1)$, hence six even vertices. □

Proposition 10.2. *The rings R8_2_152, R8_2_153, R8_2_166 and R8_2_495 admit no fusion-category categorification over* $\mathbb{C}$.

Proof. Each tensor has a faithful C_2 grading with rank-four neutral ring $g^2 = 1$, $gr = rg$, $r^2 = 1 + 2r + 2gr$. Its odd component contains X with $XX^* = X^*X = 1 + r$. The source witnesses are

ring	$(1, g, r, gr)$	X	graph vertices $0, \dots, 7$
R8_2_152	$(0, 6, 5, 7)$	1	$(0, 1, 5, 3, 4, 7, 2, 6)$
R8_2_153	$(0, 6, 5, 7)$	1	$(0, 1, 5, 3, 4, 7, 2, 6)$
R8_2_166	$(0, 5, 7, 6)$	3	$(0, 3, 7, 1, 2, 6, 4, 5)$
R8_2_495	$(0, 1, 6, 3)$	2	$(0, 2, 6, 4, 5, 3, 7, 1)$.

Both left and right incidence matrices for tensoring with X are the rooted graph of Lemma 10.1. The object r generates the full neutral category, so both even parts have rank four.

Suppose a complex categorification exists. Its neutral category has exactly two FP dimensions. By [25, Lemma 3.1], it is Galois conjugate to a pseudo-unitary category. Apply the same field automorphism to the entire category and its odd module; their based graphs are unchanged. Lemma 10.1 then forces the dual even part to have rank six, contradicting the rank-four graph computed from the tensor. Equivalently, the odd component is an invertible neutral bimodule and its dual category is the rank-four neutral category. No pivotal structure on the original rank-eight category is assumed. □

11. Pointed support in a twisted half-braiding

Proposition 11.1. *The rings R8_2_314 and R8_2_432 admit no fusion-category categorification over* $\mathbb{C}$.

Proof. Both tensors have a fusion subring $\mathcal{D}$ with basis $(1, g, r, gr)$ and rules

$$g^2 = 1, \qquad gr = rg, \qquad r^2 = g^\epsilon + r + gr,$$

where $\epsilon = 1$ for R8_2_314 and $\epsilon = 0$ for R8_2_432. The complement $\mathcal{M}$ is stable under both $\mathcal{D}$ actions. Its four simples are aX for $a \in \operatorname{Irr}\mathcal{D}$, with

$$Xa \cong F(a)X, \qquad X^2 = g^\epsilon + X + gX,$$
$$F(1) = 1, \quad F(g) = g, \quad F(r) = gr, \quad F(gr) = r.$$

The complete source witnesses are

ring	$(1, g, r, gr)$	X	(X, gX, rX, grX)
R8_2_314	$(0, 7, 1, 2)$	3	$(3, 4, 6, 5)$
R8_2_432	$(0, 3, 4, 5)$	6	$(6, 7, 1, 2)$.

Here F is initially only notation for the displayed label permutation. We explain why a hypothetical categorification forces the coherence needed below.

The functor $L : \mathcal{D} \to \mathcal{M}$, $a \mapsto aX$, is an equivalence of left module categories: it maps simples bijectively to simples and is fully faithful on their Hom spaces. Transport the right $\mathcal{D}$ action through L. Left module endofunctors of the regular category are right tensor functors, so this action supplies a strong monoidal functor $F : \mathcal{D} \to \mathcal{D}$ with coherent natural isomorphisms $Xa \cong F(a)X$. Its displayed permutation of simples makes it a tensor autoequivalence.

Applying this relation twice gives coherent isomorphisms $X^2a \cong F^2(a)X^2$. The decomposition $\mathcal{C} = \mathcal{D} \oplus \mathcal{M}$ is a decomposition of $\mathcal{D}$-bimodule categories, so its projection to $\mathcal{M}$ preserves these constraints. Write $z = 1 \oplus g$. The $\mathcal{M}$ part of X^2 is zX. Using $Xa \cong F(a)X$ and cancelling the fully faithful functor L gives

$$\eta_a : zF(a) \longrightarrow F^2(a)z.$$

If J^F, J^{F^2} denote the tensorators, its compatibility is

$$\eta_{ab}(\mathrm{id}_z \otimes J^F_{a,b}) = (J^{F^2}_{a,b} \otimes \mathrm{id}_z)(\mathrm{id}_{F^2(a)} \otimes \eta_b)(\eta_a \otimes \mathrm{id}_{F(b)}),$$

with associators understood. Equivalently this is the bimodule constraint of the $\mathcal{M}$ projection of multiplication $\mathcal{M} \boxtimes_{\mathcal{D}} \mathcal{M} \to \mathcal{C}$. Applying a monoidal quasi-inverse of F, and using $F^{-1}(z) \cong z$, gives an F-twisted half-braiding

$$\theta_a : za \longrightarrow F(a)z, \qquad \theta_{ab} = (J^F_{a,b} \otimes \mathrm{id}_z)(\mathrm{id}_{F(a)} \otimes \theta_b)(\theta_a \otimes \mathrm{id}_b). \tag{3}$$

No identification of F^2 with the identity is required.

Fix the two summands $z_0 = 1, z_1 = g$. Since $r \neq gr$ and $F(r) = gr$, the map θ_r must send $z_i r$ to $F(r)z_{i+1}$, with indices modulo two. Thus it exchanges the two support labels. Equation (3) for $a = b = r$ applies two exchanges, and hence $\theta_{r\otimes r}$ preserves these labels. Its associators and tensorator preserve the relevant direct-sum idempotents by naturality.

Choose a nonzero inclusion $v : r \to r \otimes r$, which exists in both rings. Naturality requires

$$\theta_{r\otimes r}(\mathrm{id}_z \otimes v) = (F(v) \otimes \mathrm{id}_z)\theta_r.$$

On $z_i r$ the left side lands in $F(r \otimes r)z_i$ and the right side in $F(r \otimes r)z_{i+1}$. Both maps are nonzero: $v, F(v)$ are split monomorphisms, tensoring preserves them, and the θ maps are isomorphisms. The two target direct summands have zero intersection, a contradiction. The argument uses only semisimplicity, rigidity and tensor coherence, with no pivotality assumption. □

12. A forced quantum-group Morita dual

Proposition 12.1. *The ring R8_2_342 has no categorification over* $\mathbb{C}$.

Proof. Suppose $\mathcal{C}$ categorifies this ring. The invertible $g = 5$ fixes simples $3, 7$. Its pointed associator is therefore trivializable, since either fixed simple gives a rank-one module category over $\langle g \rangle$. Thus $A = 1 \oplus g$ is a connected separable algebra. Put $\mathcal{M} = \mathcal{C}_A$ and let $\mathcal{E}$ be its Morita dual.

The simple module labels are

$$(0,0), (1,0), (3,+), (3,-), (4,0), (7,+), (7,-).$$

The exact certificate `verify_r342_inverse_morita.py` exhausts the two-dimensional signed representations on the two split labels. Certified annihilating polynomials and the three source generators $1, 4, 6$ leave eight generator tuples and exactly one full representation up to sign gauge. Consequently all seven-by-seven left action matrices L_i are forced. In particular $L_g = I_7$.

The distinguished induction Gram matrix is $2I_7$. Its nonnegative integral columns must be exactly two copies of each standard basis vector. Hence $\mathcal{E}$ has fourteen simples, two above each module label. Their right action matrices R_y commute with every L_i and satisfy

$$Q_a := \sum_i (L_i)_{aa} L_i = \sum_y R_y[:, a] R_y[:, a]^t, \qquad 0 \leq (R_y)_{ba} \leq \lfloor \sqrt{(Q_a)_{bb}} \rfloor.$$

These are linking-category associativity identities. The common commutant has dimension eleven; a prescribed standard first column leaves four free matrix entries. Exhausting the displayed bounds gives candidate counts $(2, 4, 2, 2, 2, 2, 2)$. The first domain contains the identity and a unique nontrivial involutive permutation P, representing the second invertible u. Right multiplication by P pairs every domain freely. Of the resulting two joint candidate tuples, exactly one satisfies all seven Gram identities.

The certificate reconstructs the full dual fusion tensor, rather than stopping at these necessary identities. For each product its coefficient vector n obeys

$$\sum_w n_w R_w = R_y R_z, \qquad n_1 = \delta_{z,y^*}, \qquad n_u = \delta_{zu,y^*}.$$

The first column bounds each coefficient by the corresponding entry of R_yR_z. Complete enumeration forces the ten fusion matrices with labels $0,1,4,5,6,7,10,11,12,13$. The fourteen words

$$1, u, 4, 5, 6, 7, 10, 11, 12, 13, 44, 46, 4\,11, 4\,13$$

have independent class vectors. Their associativity therefore forces all remaining matrices. The resulting tensor passes all original product domains and all 14^4 associativity equations.

The labels $(0,2,5,6,9,10,13)$ form a fusion subcategory $\mathcal{B}$, with complement $\mathcal{B}u$ and faithful C_2 grading. The restriction $\mathcal{M}|_{\mathcal{B}}$ is regular. Its based ring is $\mathrm{PSU}(3)_5$: the generator matrix in [72, Section 2.4] matches under the local map $(0,4,5,6,1,3,2)$, and its seven powers on the unit span the whole ring. Conjugation by u induces precisely its duality permutation $(0,1,3,2,4,6,5)$. The checker also verifies $\mathcal{B}_{\mathrm{ad}} = \mathcal{B}$ and that its based automorphism group is C_2.

The Kazhdan–Wenzl theorem, in the general monoidal form [73, Theorem 5.1], applies to $\mathcal{B}\boxtimes\mathrm{Vec}_{C_3}$, whose based ring is $\mathrm{SU}(3)_5$. Restrict its resulting equivalence to adjoint subcategories. The grading three-cocycle is trivial there, so $\mathcal{B}$ is a Galois conjugate of the standard quantum $\mathrm{PSU}(3)_5$ category. This step assumes neither a braiding nor pivotality.

The gauge-autoequivalence kernel of $\mathrm{SU}(3)_5$ is trivial by [50, Theorem 1.1 and Section 3]. Since the standard category factors as $\mathrm{PSU}(3)_5 \boxtimes \mathcal{P}$, with $\mathcal{P}$ pointed of rank three, any based-trivial tensor autoequivalence of $\mathrm{PSU}(3)_5$ extends by the identity on $\mathcal{P}$. It is therefore trivial. Galois transport proves the same statement for $\mathcal{B}$. Thus its nontrivial based duality permutation determines a unique tensor autoequivalence class. Moreover $U(\mathcal{B}) = 1$, so $\mathrm{Aut}_{\otimes}(\mathrm{Id}_{\mathcal{B}}) = 1$; the coherent C_2 action realizing this class is unique up to equivalence when it exists.

Finally, the first-column multiplicities give the actual internal-end algebra $\underline{\mathrm{End}}_{\mathcal{E}}(m_0) = 1 \oplus u$. Its associativity trivializes the pointed associator of u. The grading is consequently a crossed product $\mathcal{E} = \mathcal{B} \rtimes C_2$, and Morita reflection through this algebra identifies $\mathcal{C}$ with $\mathcal{B}^{C_2}$. The existing category in Proposition 13.1 has a central Tannakian boson, and its de-equivariantization is a Galois conjugate of $\mathrm{PSU}(3)_5$ by [72, Section 2.4]. Its action swaps exactly the two dual pairs and is the same nontrivial based action. The uniqueness above forces $K_0(\mathcal{C}) = R8_2_666$. Every simple in that ring is self-dual, whereas labels $1,2$ in $R8_2_342$ form a non-self-dual pair. This is a contradiction. □

13. Near-group centralizer constructions and a common spherical reduction

Proposition 13.1. *The ring $R8_2_666$ admits a unitary braided categorification: the centralizer of the full pointed subcategory in the center of a near-group $C_4 + 4$ category.*

Proof. The smallest hyperbolic case in [71, Section 3.2.3.1] is explicitly identified with the actual centers of the two complex-conjugate near-group $C_4 + 4$ categories. This finite case precedes the conjecture there for larger groups. A convenient description is given in [72, Section 2.4]:

$$G = C_4^2, \quad q_G(x,y) = i^{-xy}, \quad \theta_G(x,y) = (y,x),$$

$$H = C_{16} \times C_2, \quad q_H(x,y) = \zeta_{32}^{3x^2} i^{y^2}, \quad \theta_H(x,y) = (3x + 8y, x + y).$$

The fixed groups identify with $K = C_4$ through $k \mapsto (k,k)$ and $k \mapsto (4k, k \bmod 2)$. The center has 28 simple objects, labelled by K, $K \times \{\pi\}$ and representatives of the nonfixed involution orbits in G, H. Its pointed subcategory $\mathcal{P}$ has dimension four and twist $q_K(k) = i^{-k^2}$. Trivial monodromy with every pointed object selects exactly the eight labels

$$(0,0), (2,2), (0,0,\pi), (2,2,\pi), (3,1)_G, (2,1)_H, (6,1)_H, (4,0)_H.$$

Thus $\mathcal{P}'$ is an existing unitary braided fusion category, of dimension $96 + 64\sqrt{2}$. Its Müger center is the Tannakian order-two radical of $\mathcal{P}$.

The source-bound certificate `verify_c4_centralizer.py` reconstructs the full 28-label matrix from [72, Equation (1)]. It works in $\mathbb{Z}[\zeta_{32}]$, using $\zeta_{32}^{16} = -1$ and $\sqrt{2} = \zeta_{32}^{4} + \zeta_{32}^{-4}$. It checks unitarity and all 1792 Verlinde coefficients whose first two labels belong to this centralizer, including vanishing

of every coefficient outside it. The resulting 512 coefficients identify its based ring with $R8_2_666$ under the unital map

$$(0, 1, 3, 4, 7, 2, 5, 6).$$

The proof note records the entire matrix formula, transversals, normalization and source hash, and the normal and optimized replays also verify associativity and self-duality. □

Proposition 13.2. *The rings $R8_2_820$ and $R8_2_869$ admit unitary modular categorifications. They are the nonpointed Müger factors of the Drinfeld centers of the first and second near-group $C_5 + 5$ categories in* [70, Proposition 6 and Table 2].

Proof. Use the actual C^*-fusion categories with parameters $(c, m) = (-1, 1)$ and $(e^{2\pi i/3}, 2)$ in that table. Their auxiliary metric groups are respectively $G' = C_3^2$, $Q'(x, y) = (x^2 + y^2)/3$, and $G' = C_9$, $Q'(x) = 2x^2/9$. The center formulas of [70, Proposition 7(b)] are identified with the actual centers for every odd group order at most 13 in the paragraph following its Conjecture 2. Thus the order-five case here is established independently of the conjecture for arbitrary groups.

Put

$$a = \frac{5 + 3\sqrt{5}}{2}, \qquad b = a + 1, \qquad c = a + 2.$$

Each near-group category has global dimension $5c$. The five invertible center objects a_g, $g \in C_5$, have pairing $\exp(2\pi i\, 2mgh/5)$, which is nondegenerate. Their Müger centralizer is consequently an existing unitary modular category of dimension $5c^2 = 45b$. Trivial monodromy with these invertibles selects exactly

$$a_0,\ b_0,\ c_{1,-1},\ c_{2,-2},\ d_{0,\gamma_1}, \ldots, d_{0,\gamma_4}, \qquad \{\gamma_i\} = (G' \setminus \{0\})/\{\pm 1\}.$$

Their dimensions are $(1, b, c, c, a, a, a, a)$; their squared dimensions sum to $45b$, confirming completeness.

The exact source-bound certificate `verify_neargroup_centralizer.py` reconstructs the restricted S matrix from those formulas and evaluates all 512 Verlinde coefficients over $\mathbb{Q}[q, z]/(q^2 - 5, z^3 - 3z + 1)$. It identifies the two complete tensors under the respective model-to-source maps

$$(0, 2, 6, 7, 1, 3, 4, 5), \qquad (0, 3, 6, 7, 1, 2, 5, 4).$$

Here the auxiliary representatives are $(1, 0), (0, 1), (1, 1), (1, 2)$ in C_3^2 and $1, 2, 3, 4$ in C_9. An independent certificate instead reconstructs every tensor entry from the integer fusion formulas following [70, Proposition 7(b)] and obtains the same identifications. Existence follows from the categories and their centralizers; the finite computations identify their based rings. □

The following result, obtained before these constructions, also controls all possible categorifications of the two rings.

Proposition 13.3. *Every complex fusion category with based ring $R8_2_820$ or $R8_2_869$ admits a spherical structure. Every spherical dimension character belongs to the FP Galois orbit. Consequently, categorifiability of either ring is equivalent to pseudo-unitary categorifiability, and any categorification has a pseudo-unitary Galois conjugate.*

Proof. Both tensors are self-dual. In the canonical sphericalization choose the positive-second-indicator lift of each simple. The resulting signed quotient tensor T is fully symmetric, has $|T_{ij}^k| \le N_{ij}^k$ and $T_{ij}^k \equiv N_{ij}^k \pmod 2$, and satisfies all associativity equations. The odd-channel parity theorem [62, Theorem 3.9] gives the stronger restriction

$$N_{jj}^i - T_{jj}^i \equiv 0 \pmod 4.$$

Every coefficient two in these tensors has a repeated label. Thus repeated coefficients one are fixed at $+1$, repeated coefficients two lie in $\{-2, 2\}$, and the remaining all-distinct coefficients one lie in $\{-1, 1\}$.

The source-bound checker reconstructs exactly 35 Boolean signs for $R8_2_820$ and 36 for $R8_2_869$. Expanding every ordered associativity identity and reducing sign squares to one leaves 215 distinct integer polynomial equations for either tensor. Exhaustive branching visits both signs at each unassigned variable and rejects a branch only when a fully assigned equation fails. It visits respectively 3071 and 1873 nodes and finds exactly one solution in each case: all signs are positive. Hence $T = N$. The chosen lifts form a full fusion subcategory of the sphericalization, and the forgetful functor restricts to a tensor equivalence with the original category. Restricting its spherical structure proves the first assertion.

The remaining reduction is common and theoretical. Write the basis as $(1, a_1, a_2, a_3, a_4, b, c_1, c_2)$, where b has source label 2 for $R8_2_820$ and label 3 for $R8_2_869$, the c-labels are $6, 7$, and the a-labels are the four other nonunit labels. Both integral based rings satisfy

$$b^2 = 1 + \sum_{i=1}^{4} a_i + b + c_1 + c_2, \qquad (b+1)a_i = b^2 - 1, \qquad (b-1)c_j = b^2 - 1.$$

Let $x = \dim(b)$ for any spherical structure. Nonzero simple dimensions exclude $x = 1$ and $x = -1$ by the last two identities. They therefore give $\dim(a_i) = x - 1$ and $\dim(c_j) = x + 1$. The first identity becomes

$$x^2 = 1 + 4(x-1) + x + 2(x+1) = 7x - 1, \qquad x = \frac{7 \pm 3\sqrt{5}}{2}.$$

These are exactly the two Galois-conjugate FP characters. Conjugate the category together with its spherical structure to the positive character. A spherical category with positive simple dimensions is pseudo-unitary. This statement concerns every categorification; it does not assert that each categorification is unitarizable. □

13.1. **A unified character-parity exclusion for the Fibonacci pair.**

Proposition 13.4. *Every complex categorification of $R8_2_190$ or $R8_2_489$ admits a spherical structure and has a pseudo-unitary Galois conjugate. For $R8_2_489$, the pointed C_2 subcategory has trivial associator, so $1 + g$ is a connected separable algebra. The following theorem uses this structural reduction to exclude both rings.*

Proof. Both rings have basis $g^i\tau^j x^k$, $i, j, k \in \{0, 1\}$, and commutative presentation

$$g^2 = 1, \qquad \tau^2 = 1 + \tau, \qquad x^2 = g^\epsilon + \tau(x + gx), \qquad x^* = g^\epsilon x,$$

where $\epsilon = 1$ for $R8_2_190$ and $\epsilon = 0$ for $R8_2_489$. In the canonical sphericalization, choose the lift of g with dimension 1. It squares to the unit. A lift of τ is self-dual, and its signed square has the form $1 \pm \tau$; change the lift to make the sign positive. Choose a lift of x, and define the other lifts by the simple products $g\tau, gx, \tau x, g\tau x$. The relevant left and right products have the same underlying simple and equal nonzero spherical dimensions, so the chosen generators commute in the signed quotient. The same argument gives $x^* = g^\epsilon x$. The unit channel of xx^* therefore fixes the constant coefficient in

$$x^2 = g^\epsilon + a\tau x + b\tau gx, \qquad a, b \in \{1, -1\}.$$

These relations determine the entire signed multiplication tensor. When $\epsilon = 1$, duality fixes x^2 and exchanges the last two terms, so $a = b$.

For $\epsilon = 0$, suppose instead $a = -b$. Writing $t = \dim(\tau)$ and $z = \dim(x)$ gives $t^2 = t + 1$, $z^2 = 1$, and global dimension $\delta = 4(t + 2)$. A source character with values $g \mapsto 1$, $\tau \mapsto t$, $x \mapsto d$, where $d^2 = 1 + 2td$, has formal codegree $f_E = 2(t + 2)(1 + d^2)$. Central induction in the sphericalization requires δ/f_E to be an algebraic integer [74, Theorem 2.13]: the sphericalization dimension and the source formal codegree both acquire a factor two. However,

$$d^4 - 2d^3 - 6d^2 + 2d + 1 = 0, \qquad y = \frac{\delta}{f_E} = \frac{2}{1 + d^2}$$

has primitive minimal polynomial

$$5y^4 - 20y^3 + 25y^2 - 10y + 1.$$

Indeed the polynomial of d is irreducible modulo three, and $d = ((d^2)^2 - 6d^2 + 1)/(2(d^2 - 1))$ gives $\mathbb{Q}(y) = \mathbb{Q}(d)$; elimination gives the displayed polynomial. Its leading coefficient excludes integrality. Thus $a = b$ in both cases. Changing the lift of x and its three translates if necessary makes both signs positive. Now $T = N$, so the chosen lifts form a tensor-closed spherical section of the forgetful functor.

This section has $\dim(g) = 1$, $t^2 = t + 1$, and $z^2 = 1 + 2tz$. The irreducible quartic above shows that these four dimension characters form a single Galois orbit, containing the positive character

$$t = \varphi = \frac{1 + \sqrt{5}}{2}, \qquad z = \varphi + \sqrt{\varphi^2 + 1}.$$

Conjugating to that character gives pseudo-unitarity.

For $R8_2_489$, work with this positive spherical structure. A homogeneous pairing on a self-dual object V induces one on V^n of sign $\nu_2(V)^n$. Any opposite-indicator self-dual summand consequently has even multiplicity. Since $\tau^2 = 1 + \tau$, we have $\nu_2(\tau) = 1$. The exact congruences

$$x^3 \equiv x + \tau + g\tau \pmod 2, \qquad (g\tau)^3 \equiv g \pmod 2$$

then imply $\nu_2(x) = \nu_2(g\tau) = \nu_2(g) = 1$. For the positive spherical structure on $\mathrm{Vec}^{\omega}_{C_2}$, the nontrivial associator class has indicator -1 on g. Hence ω is trivial, and $1 + g$ is its regular separable algebra. This algebra also exists before Galois conjugation. No half-braiding on g is inferred.

The source-bound replay `verify_fib_pair_theoretical.py` checks both complete multiplication tensors, the duality and cube identities, irreducibility modulo three, and the elimination polynomial. The proof does not require an enumeration of signed tensors. □

Theorem 13.5. *Neither* $R8_2_190$ *nor* $R8_2_489$ *admits a complex fusion-category categorification.*

Proof. By Proposition 13.4, it suffices to exclude pseudo-unitary categorifications. Retain the common presentation above, put $y = \tau x$, and write

$$\varphi = \frac{1 + \sqrt{5}}{2}, \qquad d = \varphi + \sqrt{\varphi + 2}, \qquad D = 2(\varphi + 2)(1 + d^2).$$

The four dimension coordinates $1, \varphi, d, \varphi d$ are rationally independent: the norm of $\varphi + 2$ from $\mathbb{Q}(\varphi)$ to $\mathbb{Q}$ is the nonsquare 5, so d has degree two over $\mathbb{Q}(\varphi)$.

For a character χ, let Z_χ denote its central simple summand of $I(1)$ and let $\beta_\chi = D/f_\chi$. The character correspondence preceding [74, Theorem 2.13], together with its Example 2.9, gives $\dim Z_\chi = \beta_\chi$, $[FZ_\chi : 1] = 1$, and

$$S_{Z_\chi, W} = \beta_\chi \chi(FW)$$

for the unnormalized modular matrix of the center. Symmetry therefore yields

$$\beta_\chi \chi(FZ_\psi) = \beta_\psi \psi(FZ_\chi). \tag{4}$$

Take $\omega = 1$ for $\epsilon = 0$ and $\omega = i$ for $\epsilon = 1$. The two characters

$$\chi_\pm(g) = -1, \qquad \chi_\pm(\tau) = \varphi, \qquad \chi_\pm(x) = \pm\omega$$

have formal codegree $4(\varphi + 2)$ and central dimension $\beta_\pm = 1 + \varphi d$. Formal codegrees here use $\chi(b)\chi(b^*)$, which also gives this value when $x^* = gx$ and $\omega = i$. Their central restrictions consequently satisfy

$$FZ_\pm \in \{1 + y,\ 1 + gy\},$$

by the independence of the four dimension coordinates and unit multiplicity one.

Now consider the character

$$\psi(g) = 1, \qquad \psi(\tau) = 1 - \varphi, \qquad \psi(x) = z = (\varphi - 1)d - \varphi.$$

The relations give $z^2 = 1 + 2(1-\varphi)z$, and direct exact reduction gives

$$f_\psi = 2(3-\varphi)(1+z^2), \qquad \beta_\psi = (1+\varphi) + (2+3\varphi)d.$$

Write $B = Z_\psi$. Because $\psi(g) = 1$, the values $\psi(FZ_+)$ and $\psi(FZ_-)$ coincide for every permitted choice of the two restrictions. Applying (4) twice and using $\beta_+ = \beta_-$ gives $\chi_+(FB) = \chi_-(FB)$.

The dimension of B requires total multiplicity two for x, gx in FB and total multiplicity three for y, gy. If Δ_x, Δ_y are the respective integer multiplicity differences, subtracting these two character evaluations gives

$$0 = 2\omega(\Delta_x + \varphi\Delta_y).$$

Irrationality of φ forces both differences to vanish. The two equal multiplicities of y and gy cannot sum to three, which is the required contradiction.

The source-bound replay `verify_fib_character_parity.py` checks all 512 tensor entries for each presentation, the three characters, the formal codegrees, and the exact dimension identities. The obstruction itself requires no induction-column enumeration. □

13.2. **Cyclic induction phase multiplicities exclude $R8_1_9$.**

Lemma 13.6. *Let a be an invertible object of order n in a pseudo-unitary fusion category of dimension D. Every phase occurring in $I(a)$ is an nth root of $\nu_n(a)$, and the total dimension in each of these n phases, counted with its induction multiplicity, is D/n.*

Proof. The rotation spaces $\mathrm{Hom}(1, a^m)$ vanish for $0 < m < n$, so $\nu_m(a) = 0$. For $m = n$ that space is one-dimensional, and $\eta = \nu_n(a)$ is a root of unity. The positive weighted trace $\mathrm{Tr}(\theta^n I(a)) = D\eta$ has maximal absolute value $D = \dim I(a)$, so every constituent has $\theta^n = \eta$. Choose $\lambda^n = \eta$ and write W_j for the weight of phase $\lambda\zeta_n^j$. The zeroth and first $n-1$ trace identities are $\sum_j W_j = D$ and $\sum_j W_j \lambda^m \zeta_n^{jm} = 0$ for $1 \le m < n$. Discrete Fourier inversion gives $W_j = D/n$. □

Theorem 13.7. *The ring $R8_1_9$ has no complex fusion categorification.*

Proof. By [25, Lemma 3.1], take a pseudo-unitary Galois conjugate and put $d = 2+\sqrt{5}$, $D = 8+16d$. The odd-cube pivotal identities force the second indicators, in source order, to be $(1,0,0,0,0,1,1,1)$. The pointed generator a has source label1, with a^{-1} of label2 and a^2 of label7. Its associator class $k \in \mathbb{Z}/4$ satisfies $\nu_2(a^2) = (-1)^k$, so $k = 0$ or2. The canonical pointed indicators give $\nu_4(a) = \nu_4(a^{-1}) = (-1)^{k/2}$.

The pointed induction Gram block is $8I_4$. Hence a central row v occurring over a has no other pointed coordinate, and its dimension has the form $v_a + Bd$, with $B \ge 0$. Lemma 13.6 gives weight $D/4 = 2 + 4d$ in each actual phase. Comparing constant coefficients therefore gives $\sum_{Z \text{ in one phase}} v_a(Z)^2 = 2$. Thus $v_1, v_2 \le 1$ for every central row. This argument is uniform in the pointed associator class and does not average away the two imaginary phases.

The complete formal codegrees are $D, D', 8, 8, 8, 8, 8, 8$; the six other characters lie in $\mathbb{Q}(i)$. Actual character-labelled Galois symmetry transports a forgetful row by

$$T(v) = v + (0,0,0,1,1,1,1,0)\left(v_0 + v_1 + v_2 + v_7 - \frac{v_3+v_4+v_5+v_6}{2}\right),$$

up to the sign making it nonnegative. Partner rows have equal normalized relative moments, because their twists are related by σ^2 and σ^2 fixes $\mathbb{Q}(\sqrt{5})$. The two forced $I(1)$ rows and the six remaining character-labelled rows leave exactly eight possible low frames.

For arbitrary roots of unity, the first two relative traces lie in a fixed 17-vertex hull. It consists of the nine pairs from orders $1, 2, 4, 5, 10, 20$, together with the needed negative-half endpoints: an extra odd factor coprime to 10 scales both moments by $\mu(m)/\varphi(m) \in [-1/2, 1]$. No root-order bound or prime-support assumption is used. The full elementary Gauss-sum proof is supplied in the evidence packet.

For $k=0$, constituents over labels1 and2 have $\theta^4=1$; for $k=2$ they have $\theta^4=-1$ and their first two relative traces vanish. Those over label7 have $\theta^2=1$. Use these phase restrictions, the exact dimension-integrality condition $D/\dim(Z)$ integral, Galois pairing, and the residual Gram bound for the entire row orbit. With the Fourier bound, class zero leaves 64 row orbits and 405 moment columns per frame. The original, weaker bounds suffice for class two and leave 92 orbits and 389 columns.

The Gram and two induction trace identities have 56 integer coordinates. The two covectors in `r9_center_trace_certificate.json` have nonnegative pairing with every column, but target pairings -64 in class zero and -6432 in class two, for all eight low frames. Their standalone standard-library verifier reconstructs every bounded domain and every positive semidefinite condition with exact rational arithmetic. Neither target is a nonnegative combination of its columns, excluding both possible pointed classes. The initial Galois reduction proves the general complex assertion. □

13.3. **The full modular identity and a fourth-moment projection.**

Lemma 13.8. *Let $\mathcal{C}$ be a pseudo-unitary fusion category with commutative Grothendieck ring, and let b_χ be the forgetful vector of the actual summand $A_\chi \subset I(1)$ corresponding to the character χ. For a simple $Z \in Z(\mathcal{C})$ write v_Z for its forgetful vector. Then*

$$\sum_Z \theta_Z (v_Z)_i (v_Z)_j = \sum_\chi (b_\chi)_i \overline{\chi(X_j)}. \tag{5}$$

Proof. Let $C_{\chi i} = \chi(X_i)$, let f_χ be the formal codegrees, and put $X = \operatorname{diag}(f_\chi^{-1})C$. The character identification in [74, Example 2.9 and Theorem 2.13] gives $S_{A_\chi,Z} = \chi(FZ)/f_\chi$ for every central simple Z, with $S_{1Z} = \dim(Z)/\dim(\mathcal{C})$. All A_χ have twist one, and the center has trivial anomaly. Thus the low block of $STS = T^{-1}ST^{-1}$ is

$$X\left(\sum_Z \theta_Z v_Z v_Z^{\mathsf{t}}\right) X^{\mathsf{t}} = XB^{\mathsf{t}},$$

where B has rows b_χ. Character orthogonality gives $C\overline{C}^{\mathsf{t}} = \operatorname{diag}(f_\chi)$, so $X^{-\mathsf{t}} = \overline{C}$. Cancellation proves the identity. □

Theorem 13.9. *The ring R8_1_18 has no complex fusion categorification.*

Proof. By [25, Lemma 3.1], pass to a pseudo-unitary Galois conjugate. Put $d = 2+\sqrt{5}$, $K = \mathbb{Q}(\sqrt{5})$ and $D = 8+16d$. In model order $(1, a, a^2, a^3, \rho, a\rho, a^2\rho, a^3\rho)$, mapped to source $(0,1,7,2,3,6,5,4)$, the ring is commutative and satisfies

$$a^4 = 1, \qquad \rho^2 = a + (1 + a + a^2 + a^3)\rho.$$

Its six non-FP characters have formal codegree eight and are

$$\chi_{k,\pm}(a) = \zeta_8^{2k}, \qquad \chi_{k,\pm}(\rho) = \pm\zeta_8^k, \qquad k = 1,2,3.$$

The other two codegrees are D, D'. The two forced low rows are 1 and $1 + X_3 + X_4 + X_5 + X_6$; the other six have form 1+ two nonpointed constituents. Actual character reciprocity and duality leave eight low assignments. Taking normalized relative trace t_K in (5) gives, for nonunit indices,

$$\sum_Z (v_Z)_i (v_Z)_j t_K(\theta_Z) = \begin{cases} 4-d & i,j \in \{3,4,5,6\}, \\ 0 & \text{otherwise.} \end{cases}$$

Indeed every nonpointed value of the six $\mathbb{Q}(\zeta_8)$ characters has relative trace zero; the pointed entries cancel in each of the eight assignments.

The pointed induction Gram is $8I_4$. Lemma 13.6 therefore gives $(v_Z)_1, (v_Z)_2 \le 1$ exactly as in the preceding theorem, uniformly over all four pointed associator classes. Besides the quadratic transport T displayed there, full Galois symmetry acts on rows by the source permutation

$$P = (0,2,1,6,5,4,3,7).$$

It is induced by an automorphism fixing K and sending ζ_8 to ζ_8^3: on the model basis it sends $a \mapsto a^{-1}$ and $a^j\rho \mapsto a^{3j+1}\rho$. Include also source duality $(0,2,1,4,3,6,5,7)$. Every full row orbit has a common relative moment histogram: Galois transport changes a twist by σ^2, which fixes quadratic K, and duality preserves twists.

The Cauchy theorem for spherical fusion categories [61, Theorem 3.9] applies to the nonintegral dimension D. Its norm is $320 = 2^6 \cdot 5$, so the exponent has only primes two and five. For powers $1, 2, 4$, every normalized relative twist triple is one of the triples from fortieth roots or the zero triple. This follows directly from the quadratic Gauss sums: fifth-adic exponent at least two, or 2-adic exponent at least four, gives zero in all three coordinates. There are exactly thirteen triples; this is not an assertion that the exponent divides forty.

There is no need to determine fourth indicators. Each $\nu_4(X_i)$ is a sum of fourth roots, so $t_K(\nu_4(X_i)) \in \mathbb{Z}$. The functional $\ell(a + bd) = 2a - b$ annihilates $D\mathbb{Q}$. Applying it to the fourth induction identity gives

$$\ell\left(\sum_Z (v_Z)_i \dim(Z) t_K(\theta_Z^4)\right) = 0.$$

Together with the first two induction traces, the full modular identity above, and the Gram identity, this gives 119 rational linear coordinates after low terms are subtracted.

The exact domain uses the diagonal Gram bounds, integrality of $D/\dim(Z)$, the Fourier multiplicity bound, full row-orbit closure, and positive semidefiniteness after subtracting that entire orbit. It has 25 row orbits for every low assignment. The pointed classes $k = 0$, $k = 1, 3$ and $k = 2$ impose their respective cyclic twist constraints and leave 167, 145 and 153 columns. The three integer covectors in `r18_sts_moment_certificate.json` pair nonnegatively with every column and pair to -256 with every corresponding target, for all eight low assignments. The standard-library verifier reconstructs the source, characters, thirteen twist triples and all bounded domains using exact rational arithmetic. Thus none of the four pointed classes is possible. The initial Galois reduction proves the general complex assertion. □

14. An inner-square obstruction for $R8_2_24$

The obstruction in this section combines a local pivotal calculation with rigidity of tensor functors. It applies to an arbitrary complex categorification after a proved Galois reduction to pseudo-unitarity.

Theorem 14.1. *The fusion ring $R8_2_24$ has no complex fusion categorification.*

14.1. The neutral indicators and the actual inner square.

Lemma 14.2. *Let $\mathcal{B}$ be pseudo-unitary with simple objects $1, g, r, s = gr$ and fusion rules*

$$g^2 = 1, \qquad gr = rg = s, \qquad r^2 = 1 + 2r + 2s. \tag{6}$$

For its canonical spherical structure, $\nu_2(g) = -1$ and $\nu_2(r) = \nu_2(s) = 1$.

Proof. Put $d = 2 + \sqrt{5}$ and $D = 2 + 2d^2 = 20 + 8\sqrt{5}$, and let a prime mean the nontrivial embedding of $\mathbb{Q}(\sqrt{5})$. The four formal codegrees are $D, D', 4, 4$. The corresponding constituents of $I(1)$ have dimensions $1, D/D', D/4, D/4$ and twist one. Use the normalized anomaly-free center modular matrices S, T, with $S_{1A} = \dim(A)/D$. In the span of these constituents let w be the unit vector proportional to the difference of the first two basis vectors. The modular-character formula for central induction gives

$$\operatorname{proj}_{I(1)} Sw = \left(\frac{1}{D} - \frac{1}{D'}\right) w = -\frac{w}{\sqrt{5}}.$$

The diagonal torus acts on w by the quadratic character χ_5. Indeed its Galois action interchanges the first two constituents precisely at the nonresidues modulo five; their signs are positive since $D, D' > 0$. The local compression bound [63, Lemma 4.1] is therefore attained at conductor five.

We recall why equality supplies the needed level statement. The congruence theorem [60] makes the modular representation a representation of $\mathrm{SL}_2(\mathbb{Z}/N\mathbb{Z})$. Decompose it into local irreducible tensor factors. At the prime five, an irreducible factor with an upper-unipotent-fixed vector embeds in the permutation representation on primitive vectors modulo 5^a. In the quadratic torus eigenspace, the maximal singular subspace of compression is spanned by

$$F(p,q)=\chi_5(q)[q \text{ is a unit}], \qquad G(p,q)=\chi_5(-p)[q=0 \pmod 5].$$

These are the equality vectors in the proof of the cited local bound; both are pullbacks from level five. Any irreducible constituent containing an equality vector consequently factors through level five. At every other prime the torus character is trivial. Equality in the norm bound one makes the vector fixed by both upper and lower unipotents, hence by the whole local group. The maximal singular subspace of a tensor product of contractions is the tensor product of their maximal singular subspaces, by diagonalizing A^*A in each factor. Thus the entire cyclic subrepresentation generated by w factors through level five. The known twist-one projection already attains the norm bound, so no additional twist-one coordinate occurs. This is the same equality argument as in Lemma 15.2.

For a central simple A with dimension z_A, its coordinate in Sw is proportional to

$$\frac{z_A}{D}-\left(\frac{z_A}{D}\right)'.$$

Consequently $\theta_A^5=1$ whenever this difference is nonzero. Let $\ell(z)$ be the rational coordinate of the nontrivial quadratic part of z/D, scaled so that its values on $(1,g,r,s)$ are $(2,2,-1,-1)$. Choose a cyclotomic automorphism σ with unit exponent k modulo five, lifting that exponent by the Chinese remainder theorem to be coprime to the full conductor. Write π and ε_A for its modular permutation and signs. The canonical anomaly-free center normalization gives

$$z_{\pi A}=\varepsilon_A\frac{D}{\sigma D}\sigma(z_A), \qquad \ell(z_{\pi A})=\varepsilon_A\chi_5(k)\ell(z_A), \qquad \theta_{\pi A}=\sigma^2(\theta_A).$$

The first induction trace, weighted by ℓ, is

$$\sum_A \ell(z_A)z_A\theta_A=D\ell(1).$$

The two Galois signs cancel in

$$\sigma\big(\ell(z_A)z_A\big)=\chi_5(k)\frac{\sigma D}{D}\ell(z_{\pi A})z_{\pi A}.$$

Apply σ and reindex by π. Every nonzero term has twist dividing five; the constituents of $I(1)$ have twist one. If h is an inverse of k modulo five, then $\sigma(\theta_A)=\theta_{\pi A}^h$. Cancelling σD and using the induction indicator formula [59, Theorem 4.1] gives

$$\sum_X \ell(\dim X)\nu_h(X)=\chi_5(k)\ell(1).$$

Inversion permutes the unit residues and preserves χ_5, so this holds with h renamed k. At $k=2$ it becomes

$$2+2\nu_2(g)-\nu_2(r)-\nu_2(s)=-2.$$

The three unknowns are signs, and the asserted values are the unique solution. □

Proposition 14.3 (Inner square supplied by the source tensor)**.** *A complex categorification of R8_2_24 has a pseudo-unitary Galois conjugate containing a neutral subcategory $\mathcal{B}$ with (6) and the indicators of Lemma 14.2. It supplies a label-fixing tensor autoequivalence T of $\mathcal{B}$ satisfying*

$$T^2 \simeq_{\otimes} \mathrm{Ad}(g). \tag{7}$$

Proof. The full source tensor has exactly two Frobenius–Perron dimensions, 1 and $2+\sqrt{5}$. The published Galois-conjugacy statement [25, Lemma 3.1] applies to the entire category. Its pseudo-unitary conjugate has the same tensor and the faithful C_2 grading with neutral basis $(1, g, r, s) = (X_0, X_7, X_5, X_6)$. The odd invertible $a = X_1$ has inverse X_2 and square $g = X_7$. Commutativity of the source tensor makes the actual inner tensor functor $T = \mathrm{Ad}(a)|_{\mathcal{B}}$ fix all four neutral labels. Coherent composition of inner functors and the isomorphism $a^2 \simeq g$ give (7); no trivialization of the pointed associator is required. The neutral category inherits the canonical pseudo-unitary spherical structure, so the preceding lemma applies. □

14.2. **The local pivotal defect.**

Theorem 14.4 (No label-fixing inner square root)**.** *For a pseudo-unitary category $\mathcal{B}$ with* (6) *and $\nu_2(g) = -1$, $\nu_2(r) = \nu_2(s) = 1$, no label-fixing tensor autoequivalence satisfies* (7)*.*

We prove the theorem in the next four steps. We use a strict monoidal rigid realization while retaining its pivotal map j; we do not set j equal to the identity.

Let Z and gZ have indicator $+1$, and choose right dual objects $g^* = g$, $Z^* = Z$ with identity self-duality charts. Thus $j_g = -\mathrm{id}_g$ and $j_Z = \mathrm{id}_Z$. Write $e = \mathrm{ev}_g : g^2 \to 1$. Pseudo-unitarity gives $\dim(g) = 1$, so the pivotal trace identity $\dim(g) = -e\,\mathrm{coev}_g = 1$ fixes the coevaluation sign. The snake identities and dualization give

$$\mathrm{coev}_g = -e^{-1}, \qquad e \otimes g = -g \otimes e, \qquad e^* = -e^{-1}. \tag{8}$$

An object written as a tensor factor in a morphism denotes its identity. Use $F(X) = gXg$ with tensorator and unit

$$F_2(X, Y) = -gX \otimes e \otimes Yg, \qquad F_0 = e^{-1}.$$

This represents ordinary inner conjugation: its usual tensorator and unit in these dual conventions are $+gX \otimes e \otimes Yg$ and $-e^{-1}$, and the natural isomorphism with every component $-\mathrm{id}$ changes both signs. The monoidal comparison of its square with the identity is

$$\eta_X = -e \otimes X \otimes e : ggXgg \longrightarrow X. \tag{9}$$

Indeed the adjacent contraction without the minus sign sends the unit of F^2 to -1 and has tensor-product defect -1, by (8). Its negative satisfies both monoidal equations. The canonical dual comparison is

$$\delta_Z = -\mathrm{id}_{gZg} : F(Z^*) \longrightarrow F(Z)^*. \tag{10}$$

To check the sign, the transported evaluation is

$$F_0^{-1} F(\mathrm{ev}_Z) F_2(Z^*, Z) = -e(g \otimes \mathrm{ev}_Z \otimes g)(gZ \otimes e \otimes Zg),$$

which is the negative of the ordinary nested evaluation of gZg.

Choose a nonzero $c : gZ \to Zg$. For arbitrary self-dualities ϕ_g, ϕ_Z, the self-duality $\psi = (\phi_Z \otimes \phi_g)c$ of gZ satisfies $\psi^* j_{gZ} = \nu_2(gZ)\psi$. Expanding this equation gives

$$(\phi_Z^{-1} \otimes \phi_g^{-1}) c^* (\phi_g \otimes \phi_Z) = \frac{\nu_2(gZ)}{\nu_2(g)\nu_2(Z)} c = -c.$$

Thus $c^* = \epsilon c$, with $\epsilon = -1$, in our identity charts. Define

$$q = (Z \otimes e)(c \otimes g) : gZg \to Z$$

and the nonzero scalar κ by

$$(Z \otimes e)(c \otimes g)(g \otimes c) = \kappa(e \otimes Z) : ggZ \to Z. \tag{11}$$

Every map is invertible. Dualizing q yields

$$q^* = -\epsilon(g \otimes c)(e^{-1} \otimes Z), \qquad qq^* = -\epsilon\kappa\,\mathrm{id}_Z\,.$$

Define its form defect γ and composition defect λ by

$$q^*\phi_Z q = \gamma\delta_Z F(\phi_Z), \qquad qF(q) = \lambda\eta_Z.$$

The first equation and (10) give $\gamma = \epsilon\kappa$. For the second, the following are typed maps $ggZgg \to Z$, with tensor symbols suppressed:

$$\begin{aligned} qF(q) &= (Ze)(cg)(gZeg)(gcgg) \\ &= (Ze)(Zgeg)(cggg)(gcgg) \\ &= -(Zee)(cggg)(gcgg) \\ &= -\kappa(eZe) = \kappa\eta_Z. \end{aligned}$$

The second line uses interchange, the third uses $e(geg) = -(e \otimes e)$, and the fourth uses (11). Consequently

$$\lambda = \kappa, \qquad \frac{\lambda}{\gamma} = -1. \tag{12}$$

Rescaling q multiplies both defects by the same square.

Apply this calculation to $Z = r$. On $H = \mathrm{Hom}(r^2, r)$, use full categorical transpose from the symmetric self-duality of r and set $\beta(f,h) = \mathrm{tr}_r(fh^{\mathsf{t}})$. The self-duality on r^2 is the tensor product of those on its two identical factors, with reversed dual order. Transpose is involutive; transposing the scalar composite proves that β is symmetric. Semisimplicity makes the composition pairing nondegenerate, and hence β is nondegenerate. These are bilinear forms over $\mathbb{C}$. The induced action

$$K(f) = qF(f)F_2(r,r)(q^{-1} \otimes q^{-1})$$

satisfies

$$K^2 = \lambda^{-1}\,\mathrm{id}_H, \qquad \beta(Kf, Kh) = \gamma^{-1}\beta(f,h) = -\lambda^{-1}\beta(f,h). \tag{13}$$

For the square, compose the monoidal comparison (9) with the output chart and two input charts. For the form, the output chart contributes γ and the inputs contribute γ^{-2}. A scalar K would have equal square and form multipliers, contrary to (13). Since $\dim H = 2$, it has two distinct opposite eigenvalues. Each eigenline is isotropic, and nondegeneracy pairs the two lines. Thus $[K]$ is a nontrivial projective involution.

14.3. **One simultaneous chart for the square root.** Suppose $T^2 \simeq_\otimes F$. A tensor autoequivalence preserves the canonical pseudo-unitary pivotal structure, by its uniqueness [21, Proposition 8.23]. Fix the unit chart by T_0^{-1} and choose charts $q_g : T(g) \to g$, $q_r : T(r) \to r$. Extend them to strict tensor words by

$$q_{uv} = (q_u \otimes q_v)T_2(u,v)^{-1}, \qquad \widehat{T}(f) = q_v T(f) q_u^{-1} \quad (f : u \to v).$$

Coherence makes this transported action strictly multiplicative on words. Rescale the two simple charts independently to make $\widehat{T}(e) = e$ and $\widehat{T}(\mathrm{ev}_r) = \mathrm{ev}_r$; each condition requires only a nonzero complex square root. Use the induced chart on $s = gr$. Distinct isomorphic words remain connected by the displayed morphisms, rather than by further identifications. We suppress the hat in the following calculations.

The one-dimensional crossing space has $T(c) = \alpha c$. Applying T to (11) gives $\alpha^2\kappa = \kappa$, hence $\alpha = \pm 1$. Moreover $T(q) = \alpha q$, while T fixes the tensorator of F and commutes with its word operation. The explicit formula for K therefore implies the exact covariance

$$T_H K T_H^{-1} = \alpha^{-1} K. \tag{14}$$

The exponent is one output chart minus two input charts. A monoidal natural isomorphism $T^2 \Rightarrow F$ acts by a scalar on the simple r; its induced change on H is therefore scalar. Thus $[T_H^2] = [K]$. If $\alpha = -1$, equation (14) makes T_H exchange the two K-eigenlines. Its square would be scalar, a contradiction. Hence $\alpha = 1$.

All composites of $e, c, c^{-1}, \mathrm{ev}_r$ and their cups are now fixed in this same chart. In particular the nonzero evaluation

$$\mathrm{ev}_s = (e \otimes \mathrm{ev}_r)(g \otimes c^{-1} \otimes r) : grgr \to 1 \tag{15}$$

is fixed. Its associated self-duality is symmetric since $\nu_2(s) = 1$. For $X = r, s$ the full-transpose form

$$B_X(f, h) = \mathrm{tr}_X(fh^{\mathsf{t}}), \qquad f, h \in \mathrm{Hom}(r^2, X),$$

is therefore nondegenerate, symmetric and preserved exactly by T. Putting $J = \mathrm{Hom}(r^2, s)$, we obtain $T_H, T_J \in O(2, \mathbb{C})$ in their respective forms. A determinant-minus orthogonal matrix in dimension two squares to the identity. Thus $\det T_H = 1$, and in a hyperbolic basis

$$T_H = \mathrm{diag}(u, u^{-1}), \qquad u^4 = -1. \tag{16}$$

The latter equation expresses the eigenvalue ratio -1 of $[T_H^2] = [K]$. For J, either $\det T_J = 1$ and $T_J = \mathrm{diag}(v, v^{-1})$ in a hyperbolic basis, or $\det T_J = -1$ and $T_J = \mathrm{diag}(1, -1)$ in an eigenbasis.

14.4. **All fusion spaces and all associator weights.** For $a \in \mathbb{F}_2$ write $g_a = g^a$ and $r_a = g^a r$. The complete fusion rules are

$$g_a g_b = g_{a+b}, \qquad g_a r_b = r_b g_a = r_{a+b}, \qquad r_a r_b = g_{a+b} + 2r_{a+b} + 2r_{a+b+1}.$$

Every multiplicity-one vertex has a basis built from the fixed structural maps. For the two-dimensional spaces choose $A_{ab} : r_a r_b \to g_{a+b} r^2$ by

$$A_{00} = A_{10} = \mathrm{id}, \quad A_{01} = c^{-1} \otimes r, \quad A_{11} = (e \otimes r^2)(g \otimes c^{-1} \otimes r).$$

For $m = a + b$ let $B_{abc} : r_c \to g_m r_{m+c}$ be the identity if $m = 0$ or $c = 1$, and $e^{-1} \otimes r$ if $m = 1, c = 0$. For $f : r^2 \to r_{a+b+c}$ transport the vertex by

$$f \longmapsto B_{abc}^{-1}(g_m \otimes f) A_{ab} \in \mathrm{Hom}(r_a r_b, r_c). \tag{17}$$

Every structural transport map in this expression is fixed by T; f itself carries the H or J action. These vertices therefore have exactly the H action for $a+b+c = 0$ and the J action for $a+b+c = 1$, without additional scalar characters. The thin vertex $r_a r_b \to g_m$ is $(g_m \otimes \mathrm{ev}_r) A_{ab}$. The other thin vertices use e, c and unit maps. Composition-dual incoming bases have inverse eigenvalues. One chosen transport for each space suffices; equality of alternative transport paths is not required.

First suppose $\det T_J = 1$. Give the two H basis vectors weights $\pm(1, 0)$, the two J vectors weights $\pm(0, 1)$, and thin vertices weight $(0, 0)$. These define a diagonal gauge torus $U(z, w)$. Every associator coefficient has the signed sum of its four vertex weights, and every such weight belongs to

$$W = \{(p, q) \in (2\mathbb{Z})^2 : |p| + |q| \le 4\}. \tag{18}$$

To prove this uniformly, use the six-edge, four-vertex incidence pattern of a full associator coefficient. A vertex has zero, two or three noninvertible incident edges, and it is thick precisely in the last case. Counting each marked edge twice makes the number of thick vertices even. Assign bit a to both g_a and r_a. The incident bit sum is zero at a thin or H vertex, and one at a J vertex. Counting each edge twice makes the number of J vertices even; the number of H vertices is then even as well. This proves evenness of both coordinates, and four vertices give the norm bound.

Let $L \subset \mathbb{Z}^2$ be generated by the weights of actual nonzero associator coefficients. Monoidality of T implies $u^p v^q = 1$ for every $(p, q) \in L$. If L has rank two, then

$$(4, 0), (0, 4) \in L. \tag{19}$$

Indeed choose two independent vectors from W and divide them by two. The possibilities are coordinate axes of length one or two and diagonals $(\pm 1, \pm 1)$. Two independent axes, an axis and a diagonal, or two independent diagonals each generate a lattice containing the two doubled coordinate axes. Multiplying back by two proves (19). It would force $u^4 = 1$, contradicting (16). Hence L has rank below two, and its annihilator in $(\mathbb{C}^\times)^2$ contains a positive-dimensional torus fixing every associator coefficient.

If $\det T_J = -1$, the J eigenvalues are $1, -1$. Track only the H exponent p, still even and with $|p| \le 4$. At $p = \pm 2$, the H factor is $\pm i$, which no J sign can cancel. At $p = \pm 4$, all four vertices belong to H, so there is no J factor and the coefficient has multiplier -1. All these coefficients vanish. Every nonzero coefficient therefore has $p = 0$. The torus acting by $\mathrm{diag}(t, t^{-1})$ on all transported H spaces and identically elsewhere fixes the full associator.

14.5. **Tensor-functor rigidity and the exclusion.** In either case a positive-dimensional diagonal gauge torus fixes all associator coefficients and all unit channels. In a skeletal semisimple realization, its matrices on the fusion spaces define tensorators for the identity underlying linear functor: extend them naturally from pairs of simple objects to direct sums. Equality of the two composites of these tensorators on each triple is exactly invariance of the full associator matrix. The fixed unit channels supply both unit equations. Thus every element of this torus is an actual tensor autoequivalence.

These functors have infinitely many monoidal natural-isomorphism classes. A natural isomorphism has scalar components on simple objects, so its change on each of H and J is scalar. It cannot remove a varying projective eigenvalue ratio. More explicitly, on the two-dimensional torus the parameters with scalar action on both spaces satisfy $z^2 = w^2 = 1$ and form a finite subgroup; the one-dimensional case has the same conclusion from $t^2 = 1$. A positive-dimensional torus remains infinite after this quotient. This contradicts the finiteness of tensor functors between fixed fusion categories [21, Theorem 2.31], and proves Theorem 14.4.

Proof of Theorem 14.1. Proposition 14.3 would supply precisely the pseudo-unitary neutral category and the tensor autoequivalence forbidden by Theorem 14.4. Therefore a complex categorification cannot exist. □

The source tensor has SHA-256 `2b4780441d9590e8e7269cd82b7675aa75cec11c39bd3e94bb372cca1ada6738`. The source-bound evidence is in `evidence/rigidity_20261017/rank8/`. The verifier is `tools/verify_rigidity_20261017.py`. The finite checks verify the neutral tensor, the actual invertible and its conjugation permutation, and all 136 nonempty associator blocks with 1472 coefficient slots. They also check the 104 ordered independent weight pairs and 5888 reflection-case phases. These checks certify the specified finite identities; the categorical exclusion is the argument above.

15. A conductor-seventeen indicator balance

Theorem 15.1. *The five rings*

$$R8_2_33, \quad R8_2_36, \quad R8_2_76, \quad R8_2_412, \quad R8_2_446$$

have no complex fusion categorification.

The same short indicator identity excludes all five rings, including the noncommutative cases. No enumeration of possible center rows or associator equations is needed.

Lemma 15.2. *Let S, T be the normalized unitary modular representation of a pseudo-unitary Drinfeld center, and let P project onto the T-fixed space. Suppose that a T-fixed vector w has diagonal-torus character the primitive quadratic character modulo 17. If a known subspace L of the T-fixed space contains w and*

$$\mathrm{proj}_L\, Sw = -\frac{w}{\sqrt{17}},$$

then $PSw = -w/\sqrt{17}$ and the cyclic representation generated by w factors through level 17.

Proof. The compression bound in [63, Lemma 4.1] is already attained on L, so the full projection cannot have any additional coordinate. For completeness, the equality case also gives the asserted level bound. By the congruence theorem [60], decompose the finite representation into local irreducible tensor factors. At 17, an irreducible representation with an upper-unipotent-fixed vector embeds in

the primitive-vector permutation representation over $\mathbb{Z}/17^a\mathbb{Z}$. In the quadratic torus eigenspace, the maximal singular subspace of the compression is spanned by the functions

$$F(p,q)=\chi_{17}(q)[q \text{ is a unit}], \qquad G(p,q)=\chi_{17}(-p)[q=0 \pmod{17}].$$

This is the equality subspace in the proof of the cited local bound. Both functions are pullbacks from level 17. An irreducible constituent containing an equality vector therefore factors through that level. At every other prime the character is trivial; equality in the bound one makes the vector fixed by both upper and lower unipotents, and hence by the entire local group. In a tensor product of contractions, the maximal singular subspace is the tensor product of the local maximal singular subspaces, as follows by diagonalizing the positive operators A^*A. Thus all constituents supporting w factor through level 17. The full equality argument is also recorded in the evidence note `compression/CONDUCTOR_EQUALITY_LEMMA.md`. □

Lemma 15.3 (Saturated Galois indicator balance)**.** *Let $\mathcal{C}$ be pseudo-unitary, with simple dimensions in a real quadratic field K of conductor f and irrational global dimension D. Suppose every center simple A outside $I(1)$ with $\dim(A)/D \notin \mathbb{Q}$ satisfies $\theta_A^f=1$. Let $\ell : K \longrightarrow \mathbb{Q}$ be a nonzero rational linear map with $\ell(D)=0$. If k is coprime to f and $\zeta_f \mapsto \zeta_f^k$ fixes K, then*

$$\sum_{i\neq 0} \ell(\dim X_i)\nu_k(X_i)=0. \tag{20}$$

Proof. Choose a common cyclotomic twist field $\mathbb{Q}(\zeta_N)$, with $f \mid N$. The Chinese remainder theorem gives an exponent congruent to k modulo f and coprime to N: at each prime dividing N but not f, impose the additional congruence 1. The corresponding Galois automorphism τ fixes K and raises fth roots to their kth powers.

For $z_A=\dim A$, induction reciprocity and the first trace identity [74, Theorem 2.5] give

$$\sum_A \ell(z_A)z_A\theta_A = D\ell(1).$$

All coefficients belong to K. The terms with $z_A/D \in \mathbb{Q}$ vanish since $\ell(z_A)=0$. In every other term outside $I(1)$ the twist is an fth root by hypothesis; inside $I(1)$ it is one. Applying τ therefore replaces every remaining θ_A by θ_A^k, while fixing the coefficients and right side. The general induction indicator formula [59, Theorem 4.1] now gives

$$D\sum_i \ell(\dim X_i)\nu_k(X_i) = D\ell(1).$$

Cancel D and the unit term. The argument proves an exact complex identity, without taking real parts or relative traces, and does not assume that the source ring is commutative. □

Proof of Theorem 15.1. Every source ring has four invertibles, forming a group G of order four, and four noninvertibles $\mathcal{O}$, all of dimension

$$d=4+\sqrt{17}, \qquad d^2=1+8d, \qquad D=4+4d^2=8+32d=136+32\sqrt{17}.$$

By [25, Lemma 3.1], any complex categorification is Galois conjugate to a pseudo-unitary one. This two-FP-dimension theorem does not assume commutativity. Work with the conjugate and its canonical positive spherical structure.

The source tensors give

$$\Omega=\sum_X XX^* = 8\,1+8\sum_{\rho\in\mathcal{O}}\rho, \qquad \det(xI-L_\Omega)=(x-8)^6(x^2-272x+1088).$$

Here L_Ω is the Casimir multiplication matrix; in the noncommutative cases we do not identify it with the full induction Gram matrix. If an irreducible ring representation E has dimension e_E, the Casimir eigenvalue is e_Ef_E with multiplicity e_E^2 [74, Remark 2.11]. Thus D and $D'=136-32\sqrt{17}$

are the codegrees of one-dimensional representations, and every other ordinary codegree $f_E = 8/e_E$ is rational.

By [74, Theorem 2.13], the corresponding center constituent A_E of $I(1)$ has multiplicity e_E and dimension D/f_E. Let J correspond to D'. Then

$$[I(1) : J] = 1, \qquad \dim J = D/D' = d^2.$$

All these constituents have twist one. Also $F(I(1)) = \Omega$ implies that none contains a nonunit invertible in its restriction.

Normalize $S_{00} = 1/D$. The multiplicity vector m of $I(1)$ satisfies $Sm = m$ [74, Proposition 2.8]. Therefore its support is Galois-stable with positive Galois signs. A Galois automorphism fixing $K = \mathbb{Q}(\sqrt{17})$ sends the central unit to the unique constituent of $I(1)$ of dimension one; one flipping K sends it to J, the unique constituent of dimension D/D'. Thus $w = e_1 - e_J$ has quadratic torus character χ_{17}.

The known twist-one block sends w to

$$\left(\frac{1}{D} - \frac{1}{D'}\right) w = -\frac{w}{\sqrt{17}}.$$

Indeed its $1, J$ block has diagonal $1/D$ and off-diagonal $1/D'$, and the other entries in these two columns agree because their codegrees are rational. This remains true when $e_E = 2$. Lemma 15.2 now gives $PSw = -w/\sqrt{17}$ and level 17 on the cyclic representation. For a center simple A of dimension z, the coordinate of Sw is

$$\frac{z}{D} - \sigma\left(\frac{z}{D}\right).$$

If z/D is nonrational and A is outside $I(1)$, its twist is therefore a primitive seventeenth root: it has order dividing 17, and twist one is excluded by the exact formula for PSw. Apply Lemma 15.3 with

$$\ell(a + b\sqrt{17}) = 4a - 17b, \qquad k = 2.$$

Indeed $\ell(D) = 0$, and 2 is a quadratic residue modulo 17, so $\zeta_{17} \mapsto \zeta_{17}^2$ fixes K. The source weights are $\ell(1) = 4$ on the invertibles and $\ell(d) = -1$ on the noninvertibles. We obtain the common balance

$$\boxed{\sum_{\rho \in \mathcal{O}} \nu_2(\rho) = 4 \sum_{g \in G \setminus \{1\}} \nu_2(g).} \tag{21}$$

No restrictions on twists of constituents of $I(g)$, no relative moment hull, and no enumeration of center rows are needed.

A group of order four has an odd number of nonunit involutions: one for C_4, three for C_2^2. Its nontrivial pointed indicator sum is therefore a nonzero odd integer, so the right side of (21) has absolute value at least four. The left side has absolute value at most the number of self-dual noninvertibles. The exact source values are

ring	G	self-dual noninvertible labels	number
R8_2_33	C_4	$5, 6$	2
R8_2_36	C_4	$\varnothing$	0
R8_2_76	C_4	$\varnothing$	0
R8_2_412	C_2^2	$6, 7$	2
R8_2_446	C_2^2	$\varnothing$	0

Each row contradicts the balance, proving all five exclusions over $\mathbb{C}$. □

The source-bound verifier `verify_q17_indicator_balance.py` checks the original tensors, hashes, associativity, FP dimensions, pointed groups, duality lists, Casimir polynomial, and the displayed arithmetic. It uses only exact integer operations and runs identically under standard and optimized Python. The proof and independent audits are stored in `evidence/unified_20261016/rank8/`, including `Q17_INDICATOR_BALANCE_PROOF.md`.

15.1. **A spherical reduction for the two $\sqrt{10}$ cases.**

Proposition 15.4. *Every complex categorification of R8_2_630 or R8_2_671 admits a spherical structure and has a pseudo-unitary Galois conjugate.*

Proof. Both tensors are self-dual. Choose the positive-second-indicator lifts in the canonical sphericalization. These lifts are self-dual, because their duals have the same nonzero spherical dimensions, whereas the other lifts have the opposite dimensions. Their signed quotient tensor T is fully symmetric and associative, with $|T_{ij}^k| \le N_{ij}^k$ and $T_{ij}^k \equiv N_{ij}^k \pmod 2$. The odd-channel parity theorem [62, Theorem 3.9] further gives $N_{jj}^i - T_{jj}^i \equiv 0 \pmod 4$.

Thus repeated coefficients one are positive, repeated coefficients two have signs ± 2, and all-distinct coefficients one have signs ± 1. There are 27 Boolean signs for each tensor. Only $R8_2_671$ has an all-distinct coefficient two, with possible values $-2, 0, 2$. The source-bound exact checker regenerates every ordered associativity equation. For $R8_2_630$, exhaustive branching on the 157 distinct equations visits 195 nodes and leaves only $T = N$. For the three values of the extra coefficient of $R8_2_671$, it visits $57, 5, 103$ nodes; the only solution is again $T = N$, in the last branch. Every rejection evaluates an integer associativity equation exactly. Consequently the chosen actual lifts are tensor closed, and restriction of the forgetful functor identifies their spherical fusion category with the original category.

The same checker verifies the complete tables of eight distinct characters over $\mathbb{Q}(\sqrt{3}, \sqrt{10})$. The formal codegrees are

$$120 \pm 36\sqrt{10}, \quad 4, 6, 6, 12, 12, 12.$$

All six characters outside the FP conjugate pair vanish on a simple basis object. Since spherical dimensions of simple objects are nonzero, the spherical dimension character belongs to that pair. A Galois automorphism therefore takes it to the positive FP character and produces a pseudo-unitary category. □

The complete character tables, signed domains, source-bound replay, and independent audit are retained in `evidence/unified_20261016/rank8/HI6_PAIR_GALOIS_PU.md`. This proposition is a structural reduction; by itself it makes no existence decision and does not identify either center with a known Haagerup–Izumi center.

16. A third-indicator triangle obstruction at conductor forty

Theorem 16.1. *The fusion ring R8_2_630 has no complex categorification.*

Proof. Proposition 15.4 reduces a hypothetical complex categorification to a pseudo-unitary Galois conjugate. Work with its canonical positive spherical structure. Put $s = \sqrt{10}$ and $K = \mathbb{Q}(s)$. In census order, the FP dimensions and global dimension are

$$(d_0, \dots, d_7) = (1, 1, 2+s, 3+s, 3+s, 2+s, 4+s, 4+s), \qquad D = 120 + 36s.$$

The complete character table gives formal codegrees

$$D, \quad D' = 120 - 36s, \quad 4, 6, 6, 12, 12, 12.$$

The source ring is commutative, so each constituent A_E of $I(1)$ has multiplicity one and dimension D/f_E [74, Theorem 2.13]. Let J denote the constituent for D'; then $\dim J = D/D' = 19 + 6s$.

The multiplicity vector of $I(1)$ satisfies $Sm = m$, so its support is Galois-stable with positive signs. The central unit has Galois orbit exactly $\{1, J\}$: these are the unique constituents of $I(1)$ of dimensions 1 and D/D'. Thus $w = e_1 - e_J$ has the primitive quadratic torus character χ_{40}. The known twist-one block satisfies

$$\operatorname{proj}_{I(1)} Sw = \left(\frac{1}{D} - \frac{1}{D'}\right) w = -\frac{s}{20} w = -\frac{w}{\sqrt{40}}.$$

Indeed the $1, J$ submatrix has diagonal $1/D$ and off-diagonal $1/D'$, while all other entries in the two columns agree since their codegrees are rational.

The local compression bound [63, Lemma 4.1] is therefore attained. Its equality refinement, proved in the argument of Lemma 15.2, works at each prime power: for conductor p^c, the equality vectors are pullbacks from level p^c; trivial local characters force the trivial representation. Applying this to the nontrivial local conductors 8 and 5 gives level 40 for the cyclic representation generated by w. The general equality proof is recorded in `compression/CONDUCTOR_EQUALITY_LEMMA.md`. Since the coordinate of Sw at a center simple of dimension z is

$$\frac{z}{D} - \sigma\left(\frac{z}{D}\right),$$

where σ flips K, every such object with z/D nonrational has $\theta^{40} = 1$.

Apply the same saturated Galois indicator lemma used for the five conductor-seventeen cases, Lemma 15.3, now with

$$\ell(a + bs) = 3a - 10b, \qquad f = 40, \qquad k = 3.$$

We have $\ell(D) = 0$, and $\chi_{40}(3) = 1$ means that $\zeta_{40} \mapsto \zeta_{40}^3$ fixes K. The source weights are

$$(\ell(d_i))_{i=0}^7 = (3, 3, -4, -1, -1, -4, 2, 2),$$

so the lemma gives

$$\sum_{i \neq 0} \ell(d_i)\nu_3(X_i) = 0. \tag{22}$$

This is an exact complex identity. The first trace is transformed into the third by a field automorphism fixing every coefficient; no reality of the third indicators is assumed.

Finally the source tensor gives

$$\left(\dim \mathrm{Hom}(1, X_i^3)\right)_{i=0}^7 = (1, 0, 0, 1, 1, 1, 0, 0).$$

The only nonzero nonunit third indicators are consequently those at labels $3, 4, 5$. Each is the trace of an order-three rotation on a one-dimensional space, hence a cube root of unity of modulus one. Equation (22) becomes

$$\nu_3(X_3) + \nu_3(X_4) + 4\nu_3(X_5) = 0.$$

The first two terms have sum of absolute value at most 2, whereas the last has absolute value 4. This strict triangle contradiction excludes the hypothetical category and proves the theorem. □

The source-bound standard-library verifier `verify_r630_weighted_indicator.py` replays the structural sphericality gate, complete character table, dimensions, codegrees, conductor arithmetic, weights, and cubic invariant dimensions. The theoretical contradiction uses no enumeration of center rows. Its normal and optimized receipts and independent audit accompany `R630_WEIGHTED_GALOIS_INDICATOR_PROOF.md` and the common lemma `SATURATED_GALOIS_INDICATOR_LEMMA.md` in `evidence/unified_20261016/rank8/`.

17. A conductor-forty center trace obstruction

Theorem 17.1. *The ring R8_2_671 has no complex fusion categorification.*

Proof. Proposition 15.4 reduces a hypothetical categorification to a pseudo-unitary one. Put $q = \sqrt{10}$. Its simple dimensions and global dimension are

$$d = (1, 1, 3 + q, 4 + q, 4 + q, 3 + q, 2 + q, 2 + q), \qquad D = 120 + 36q.$$

The complete ordinary formal codegrees are $D, D', 4, 6, 6, 12, 12, 12$, where $D' = 120 - 36q$. The six remaining characters are defined over $\mathbb{Q}(\sqrt{3})$. All eight source objects are self-dual, and the odd-isotypic pivotal identities force all their second indicators to be $+1$. The source induction row is

$$[FI(1)] = (8, 2, 4, 6, 8, 6, 6, 6).$$

Write a central forgetful row as v, with $\dim FZ = A + Bq$, where

$$A = v_0 + v_1 + 3v_2 + 4v_3 + 4v_4 + 3v_5 + 2v_6 + 2v_7, \qquad B = \sum_{i=2}^{7} v_i.$$

Choose a cyclotomic automorphism σ flipping q and fixing $\sqrt{3}$. The character-labelled correspondence [74, Example 2.9 and Theorem 2.13] and modular Galois symmetry give the exact transport

$$T(v) = v + (0,0,1,1,1,1,1,1)\left(A - \frac{10B}{3}\right). \tag{23}$$

An actual row has partner $T(v)$ or $-T(v)$, with the sign making it nonnegative; the entries must be integral. In particular $3 \mid B$. The unit row is paired with $(1,0,1,1,1,1,1,1)$.

Let J be the Galois image of the central unit and put $w = e_1 - e_J$. The multiplicity vector of $I(1)$ is fixed by S, so its Galois signs are positive. Its known twist-one block gives

$$\operatorname{proj}_{I(1)} Sw = \left(\frac{1}{D} - \frac{1}{D'}\right) w = -\frac{w}{\sqrt{40}}.$$

The torus character of w is the primitive quadratic character χ_{40}. The equality refinement of the compression bound [63, Lemma 4.1] implies $PSw = -w/\sqrt{40}$ and level 40 for its cyclic representation. The quadratic Gauss-sum refinement gives negative χ_{40} on primitive 40th twist exponents. These refinements, with their full local proofs, are supplied in the common compression evidence packet. Since the Z-coordinate of Sw is $z/D - \sigma(z/D)$, every central simple outside $I(1)$ with nonrational z/D has twist order dividing 40 and cannot have twist one. The last assertion uses the exact full projection PSw, whose right side has no such coordinate.

Let t denote normalized relative trace to $\mathbb{Q}(q)$. For a root of unity of arbitrary order, the pair $(t(\theta), t(\theta^2))$ lies in the convex hull of

$$(1,1),\ (-1,1),\ (\tfrac{1}{2},-\tfrac{1}{2}),\ (-\tfrac{1}{2},-\tfrac{1}{2}),\ (0,-1),\ (\pm q/8,0),\ (0,\pm q/8).$$

This is a root-order independent statement: rational traces are Ramanujan sums; an irrational first trace requires order $40m$, with m squarefree and coprime to 10, and has absolute value $q/(8\varphi(m))$ and second trace zero. An irrational second trace instead requires order $80m$ and has first trace zero. The displayed segments contain all these values. For a high row with nonrational dimension ratio, the allowed pair list reduces to

$$(-1,1),\ (0,-1),\ (-\tfrac{1}{4},-\tfrac{1}{4}),\ (0,0),\ (\tfrac{1}{4},-\tfrac{1}{4}),\ (0,\tfrac{1}{4}),\ (-q/8,0).$$

These are the orders $2,4,5,8,10,20,40$, respectively. If $v_1 > 0$, positivity and $\operatorname{Tr}(\theta^2 I(X_1)) = D = \dim I(X_1)$ force $\theta^2 = 1$. Thus only the two signs remain, and only -1 remains when the ratio is nonrational. Galois partners have equal relative moments. On the restricted rows one may also choose σ with exponent 59 modulo 120; its square is one modulo 40, so their actual twists agree.

Here is the complete finite arithmetic needed to finish the proof. The source tensor determines

$$C_{ij} = \sum_{k,l} N_{ki}^{l} N_{kj}^{l} = \sum_{Z} v_i(Z) v_j(Z).$$

Each constituent Z_χ of $I(1)$ has unit multiplicity and known dimension D/f_χ. Its actual character label imposes

$$\frac{\chi_i(v_j)}{f_i} = \frac{\chi_j(v_i)}{f_j}.$$

After the two forced rows, the six exact bounded domains have sizes $86, 74, 74, 22, 22, 22$. Their prescribed sum and the entrywise and positive semidefinite residual Gram conditions leave exactly six frames. For every remaining row, integral closure of the modular formal-codegree relation forces D/z integral; equivalently,

$$A^2 - 10B^2 \mid 120A - 360B, \qquad A^2 - 10B^2 \mid 36A - 120B.$$

Together with (23) and the residual Gram bound for the whole partner orbit, the six frames have respectively $73, 85, 73, 28, 36, 28$ allowable row orbits.

Combine their Gram columns with the first and second induction trace identities

$$\operatorname{Tr}(\theta I(X_i)) = 0, \qquad \operatorname{Tr}(\theta^2 I(X_i)) = D \quad (i > 0).$$

After subtracting the known twist-one terms and multiplying the trace coordinates by eight, these are 56 integer equations. The moment domains above produce respectively $360, 398, 360, 138, 168, 138$ columns. An actual center would express the target as a nonnegative real combination of them.

Two fixed integer covectors contradict all six systems. Their pairings with every allowable column are nonnegative; their target pairings are -225800939520 for the first three frames and -960 for the last three. The covectors are given explicitly in `r671_center_trace_certificate.json`. The standalone verifier reconstructs every domain, tests every rational positive semidefinite condition, and checks every integer pairing directly from the source tensor. It uses no optimizer or floating-point arithmetic. Thus every possible center is excluded, and the initial Galois reduction proves the general complex claim. □

18. Galois-invariant pointed associators

The next obstruction is intrinsic to a pointed fusion subcategory; it therefore does not require a pivotal structure on an ambient category.

Lemma 18.1. *Let D be a fusion ring with precisely two FP dimensions, 1 and $d > 1$, and let g be a specified order-two invertible. If every pseudo-unitary categorification of D has the same associator class on $\langle g \rangle$, then every complex categorification has that class. If the class is nontrivial, no fusion category containing such a subcategory can have a simple M with $gM \cong M$.*

Proof. By the published [25, Lemma 3.1], every categorification of D is Galois conjugate to a pseudo-unitary one. The two elements of $H^3(C_2, \mathbb{C}^\times)$ have normalized representatives $\omega(g,g,g) = +1$ and -1, each fixed by every field automorphism. Thus their classes are preserved separately by Galois conjugation. If $gM \cong M$, the orbit $\{M\}$ is a rank-one module over $\mathrm{Vec}^{\omega}_{C_2}$. After choosing $gM \to M$, its scalar module pentagon says $d\psi = \omega$ for a two-cochain ψ. Hence $[\omega] = 0$, a contradiction. This is also the pointed module-category criterion of [22]. □

Write the ordinary and pseudo-$PSU(2)_6$ rings in the basis $(1, g, x, y)$, with $g^2 = 1$, $gx = xg = y$ and $gy = yg = x$. Both have FP dimensions $(1, 1, 1+\sqrt{2}, 1+\sqrt{2})$. Their distinguishing products and dualities are

	$x^2 = y^2$	$xy = yx$	duality
ordinary	$1 + x + y$	$g + x + y$	$x^* = x,\ y^* = y$
pseudo	$g + x + y$	$1 + x + y$	$x^* = y.$

Proposition 18.2. *The rings R8_2_312, R8_2_315 and R8_2_317 have no fusion-category categorification over $\mathbb{C}$.*

Proof. In a pseudo-unitary pseudo-$PSU(2)_6$ category, Larson's [56, Proposition 5.5 and proof of Theorem 5.8, Case 1(c)(1)] forces $\nu_2(g) = -1$. In the $c = 1$ case at issue, the positive-sign branch would require at least eight roots of unity when at most seven are available. The canonical spherical structure has dimension one on pointed simples, so $\nu_2(g) = \omega(g,g,g)$ in $\mathrm{Vec}^{\omega}_{C_2}$ [59]. Thus the pseudo ring forces the nontrivial pointed class. Lemma 18.1 makes this a constraint on every complex categorification.

For completeness, the ordinary ring forces the trivial class. Equip a hypothetical pseudo-unitary categorification with its canonical spherical structure. The odd-multiplicity parity theorem [62, Theorem 3.9] gives $\nu_2(x) = \nu_2(y) = 1$. The remaining alternative $\nu_2(g) = -1$ is excluded by the

following exact indicator certificate. Let A be the restriction matrix from the center and $C = AA^{\mathsf{T}}$ its induction Gram matrix. The complete low-column lists are precisely

$$L_1 = [(1,0,0,0),(1,0,0,2),(1,0,1,0),(1,0,1,0)],$$
$$L_2 = [(1,0,0,0),(1,0,1,1),(1,0,0,1),(1,0,1,0)],$$
$$L_3 = [(1,0,0,0),(1,0,2,0),(1,0,0,1),(1,0,0,1)].$$

They follow from the complete formal-codegree list; these columns have trivial twist, and every other column has unit coordinate zero and belongs to the residual Gram matrix. For each L_i, the necessary center data yield an integer system $B_i n = b_i$, $n \geq 0$. The stored integer vectors v_i certify

$$v_i^{\mathsf{T}} B_i \geq 0, \qquad (v_1^{\mathsf{T}} b_1, v_2^{\mathsf{T}} b_2, v_3^{\mathsf{T}} b_3) = (-48, -12, -48).$$

Each system has 38 equations and respectively 102, 152, 102 variables, including nonnegative slack variables. All three are therefore infeasible by direct integer arithmetic.

We specify the hypotheses of this finite certificate. Besides reciprocity, it imposes the center indicator identities

$$\sum_{Z \in \operatorname{Irr} Z(\mathcal{D})} [F(Z) : X] \dim(Z) \theta_Z^j = \dim(\mathcal{D}) \nu_j(X), \qquad 1 \leq j \leq 4,$$

and the trace bounds for an order-j pivotal rotation on $\operatorname{Hom}(1, X^j)$ [59]. The Cauchy theorem [61], applied to $\dim(\mathcal{D}) = 8 + 4\sqrt{2}$ of norm 32, makes every twist order a power of two. The ten joint relative-trace vectors through height four to $\mathbb{Q}(\sqrt{2})$ cover all such orders: orders larger than the divisor box 32 have the zero vector, which is included. This is a finite trace classification, not a bound on the exponent. The remaining columns satisfy the d-number and dimension-quotient integrality conditions [24, 22]. Finally Galois symmetry of the normalized center S matrix [60] pairs simple dimensions by

$$z \longmapsto \left| \frac{(8 + 4\sqrt{2}) z'}{8 - 4\sqrt{2}} \right|,$$

with equal counts in each pair. These conditions give the displayed systems. The source-bound checker reproduces every column, trace vector, matrix and separating product; it uses no optimization solver. Hence $\nu_2(g) = 1$ in every pseudo-unitary ordinary categorification, and Lemma 18.1 makes its pointed class trivial in every complex categorification.

The exact census witnesses, in the displayed rank-four basis, are

ring	ordinary subring	pseudo subring	g
R8_2_312	none needed	$(0,7,1,2)$	7
R8_2_315	$(0,7,3,4)$	$(0,7,1,2)$	7
R8_2_317	$(0,7,5,6)$	$(0,7,1,2)$	7.

For R8_2_312, each of the outside simples $3,4,5,6$ is fixed by g, contradicting the nontrivial pseudo-ring class by Lemma 18.1. For R8_2_315 and R8_2_317, the two subcategories restrict the same ambient associator to the same pointed subcategory, but force opposite classes. This is impossible. The argument conjugates the rank-four subcategories separately; it assumes no simultaneous pseudo-unitarization of the ambient category. □

19. Automorphisms of the unitary 2D2 model

The general complex exclusion is Theorem 14.1. The following model-specific argument describes the obstruction through the tensor automorphism group of the known unitary neutral category.

Proposition 19.1. *The ring R8_2_24 has no unitary categorification. More generally, it admits no categorification whose neutral 2D2 category is Galois conjugate to the known unitary one.*

Proof. Let $\mathcal{Q}$ be the known unitary $2D2$ category. The construction in Proposition 8.1 gives a tensor autoequivalence $T = \mathrm{Ad}(b)|_{\mathcal{Q}}$ swapping r, gr, with $T^2 = \mathrm{Ad}(g)$. Grossman's [55, Lemma 4.1] shows that $\mathrm{Ad}(g)$ exchanges two inequivalent algebra structures on $1 + r$, so it is nontrivial. Thus T has order four. The factor automorphism fixing M and sending λ to $-\lambda$ induces an outer, label-fixing involution S by [55, Lemma 4.2]. It commutes with b on both M and λ, hence S commutes with T and is independent of T^2. Grossman's order-eight computation [55, Theorem 4.19 and its proof] therefore gives

$$\mathrm{Aut}_{\otimes}(\mathcal{Q}) = \langle T, S\rangle \cong C_4 \times C_2.$$

Its label-fixing subgroup is $\langle T^2, S\rangle \cong C_2^2$; every label-swapping element has square $T^2 = \mathrm{Ad}(g)$.

The neutral category of a unitary categorification is equivalent to $\mathcal{Q}$ by [30, Theorem 3.31]. In $R8_2_24$, an odd invertible u has $u^2 = g$ and fixes every neutral label by conjugation. Then $\mathrm{Ad}(u)$ would be a label-fixing autoequivalence whose square is the nontrivial $\mathrm{Ad}(g)$, impossible. Applying a Galois automorphism to the entire category proves the stated Galois-class assertion as well. □

20. Compression over all constituents of the induced unit

The modular-representation argument in [63, Lemma 4.1 and the proof of Theorem 5.1] can retain every simple constituent of $I(\mathbf{1})$, rather than only the Galois orbit of the dimension character. This gives two further exclusions in the pivotal and spherical settings.

Proposition 20.1. *Let $\mathcal{C}$ be a spherical fusion category, with dimension field $K = \mathbb{Q}(\dim X : X \in \mathrm{Irr}(\mathcal{C}))$ and $G = \mathrm{Gal}(K/\mathbb{Q})$. Write f_E for the formal codegrees of its fusion algebra. Then $f_E \in K$ and, for every character χ of G with conductor q,*

$$\sum_E \left| \sum_{\sigma \in G} \chi(\sigma)\sigma(f_E^{-1}) \right|^2 \le \frac{|G|}{q}.$$

Proof. Put $D = \dim \mathcal{C}$ and $s = S/D$ for the normalized matrix of $\mathcal{Z}(\mathcal{C})$. Let A_E be the constituent of $I(\mathbf{1})$ associated to E. It has dimension D/f_E and twist one, so $f_E \in K$. If A_σ corresponds to $\sigma(\dim)$, then $s_{A_\sigma, Y} = \sigma(\dim Y/D)$. Set $w_\chi = \sum_\sigma \chi(\sigma) e_{A_\sigma}$ and let P project onto the twist-one subspace. The cited local compression estimate gives $\|P s w_\chi\|^2 \le |G|/q$. Its A_E coordinate is the inner sum in the assertion. Summing over these distinct simple coordinates proves the inequality; their multiplicities in $I(\mathbf{1})$ do not repeat coordinates. □

Corollary 20.2. *Let $B = \sum_i N_i N_i^{\mathsf{T}}$ be the regular Casimir of a fusion ring. Suppose the irreducible factors of $\det(xI - B)$ belonging to a fixed real quadratic field of discriminant q are $(x^2 - t_j x + n_j)^{m_j}$. A spherical categorification requires*

$$q \sum_j m_j \frac{t_j^2 - 4n_j}{n_j^2} \le 1.$$

Proof. The Casimir eigenvalue attached to E is $\lambda_E = (\dim E) f_E$, with regular multiplicity $(\dim E)^2$. The quadratic field lies in K. If its index in K is e', pull back its nontrivial character to G. Each conjugate pair of eigenvalues contributes $2e'^2 m_j (t_j^2 - 4n_j)/n_j^2$ to the preceding bound, whose right side is $2e'/q$. Cancelling and using $e' \ge 1$ proves the claim. □

Proposition 20.3. *The rings $R8_2_318$ and $R8_2_735$ admit no pivotal or spherical categorification over $\mathbb{C}$.*

Proof. For both source tensors, direct multiplication gives

$$\det(xI - B) = (x - 16)^3 (x - 8)(x^2 - 192x + 1024)(x^2 - 16x + 32).$$

Both quadratic factors belong to $\mathbb{Q}(\sqrt{2})$, of discriminant 8. Hence the necessary bound would read

$$8\left(\frac{192^2-4\cdot 1024}{1024^2}+\frac{16^2-4\cdot 32}{32^2}\right)=\frac{5}{4}\leq 1,$$

a contradiction.

For completeness, the eight characters of each commutative ring have formal codegrees

$$16,16,16,8,8+4\sqrt{2},8-4\sqrt{2},96+64\sqrt{2},96-64\sqrt{2}.$$

Every character in the first six positions vanishes on at least one basis element. The remaining two are the Frobenius–Perron character and its quadratic conjugate, and are duality invariant. These statements follow by substitution in the source tensor; the complete exact character tables are supplied in the associated `rank8_full_I1_certificate.json`. Simple pivotal dimensions cannot vanish. Thus a pivotal structure would have one of the last two dimension characters and would be spherical, which has just been excluded. □

These compression arguments alone give scoped exclusions. The independent module obstruction in Section 6 now excludes arbitrary complex categorifications of both rings.

An independent general proof for the four quadratic cases. The same estimate gives a second proof of Proposition 5.1, without a pseudo-unitary reduction. Each of its four rings has Casimir polynomial

$$(x-8)^4(x^2-160x+640)(x^2-32x+128).$$

The sphericalization's dimension field therefore contains both $\mathbb{Q}(\sqrt{10})$ and $\mathbb{Q}(\sqrt{2})$. Its index e over $\mathbb{Q}(\sqrt{10})$ is at least two. The first quadratic factor has quantity $Q=9/4$ in Corollary 20.2. In the sphericalization's positive-sign sector, formal codegrees are doubled, so compression requires $eQ/4\leq 1$. But $eQ/4\geq 9/8>1$, a contradiction.

21. The conductor-32 obstruction

Theorem 21.1. *The ring R8_2_857 has no categorification over* $\mathbb{C}$.

Proof. Reorder the source basis as $(0,1,7,2,5,4,3,6)$ and denote it by $X_0=\mathbf{1},X_1,\dots,X_7$. All objects are self-dual. Multiplication by X_1 is the symmetric tridiagonal matrix J_2 whose off-diagonal entries are one and whose diagonal is $(0,1,1,1,1,1,1,2)$. If M_j denotes multiplication by X_j, then

$$M_0=I,\qquad M_1=J_2,\qquad M_{j+1}=(J_2-I)M_j-M_{j-1}\quad(1\leq j\leq 6).$$

These identities are direct consequences of the source tensor.

Put $\zeta=e^{2\pi i/32}$ and $L=\mathbb{Q}(\zeta+\zeta^{-1})$. The positive dimension character is $d_j=\sum_{r=-j}^{j}\zeta^r=\sin((2j+1)\pi/32)/\sin(\pi/32)$. Its eight Galois conjugates are distinct and give all ring characters. Their formal codegrees are the conjugates of

$$D=\sum_{j=0}^{7}d_j^2=4\csc^2(\pi/32),\qquad \frac{1}{D}=\frac{2-\zeta-\zeta^{-1}}{16}.$$

For any primitive even Dirichlet character χ of conductor 32, the reciprocal Fourier sum over $\mathrm{Gal}(L/\mathbb{Q})$ is

$$\widehat{g}(\chi)=-\frac{\tau(\chi)}{16},\qquad |\widehat{g}(\chi)|^2=\frac{32}{16^2}=\frac{1}{8}.$$

Here $\tau(\chi)$ is the primitive Gauss sum: summing $\zeta^a+\zeta^{-a}$ over representatives modulo sign gives precisely its full sum. The eight constituents of $I(\mathbf{1})$ thus violate Proposition 20.1: if the spherical dimension field has index e over L, their squared Fourier norm is e^2, whereas the bound is $e/4$. Hence the ring has no spherical categorification.

Now suppose an arbitrary categorification exists, and take its canonical sphericalization with sign object ε of dimension -1. In the $\varepsilon=1$ sector the eight formal codegrees are $2\sigma(D)$. They belong

to the spherical dimension field K, so $L \subseteq K$. Put $e = [K : L]$. The same eight coordinates in the full induced-unit compression have squared norm $e^2/4$, with upper bound $e/4$. Thus $e = 1$ and $[K : \mathbb{Q}] = 8$.

Both lifts of every simple are self-dual, since duality preserves its nonzero spherical dimension while exchanging the lifts negates it. Choose the positive-second-indicator lifts and pass to the signed quotient $\varepsilon = -1$. Its constants satisfy

$$|T_{ij}^k| \le N_{ij}^k, \qquad T_{ij}^k \equiv N_{ij}^k \pmod 2, \qquad T_{ii}^j \equiv N_{ii}^j \pmod 4.$$

For the last congruence, a negative-indicator lift can occur in $X_i^* \otimes X_i$ only with even multiplicity by [62, Theorem 3.9]. The signed difference is therefore the original multiplicity minus a multiple of four. Self-duality gives full symmetry of T. Consequently $T_{1j}^j = T_{jj}^1 = 1$ for $1 \le j \le 6$.

The off-diagonal entries of the signed multiplication matrix of X_1 are ± 1. Successively switching the lifts of $X_2, \dots, X_7$ makes them all positive, while leaving the unit, X_1, and all diagonal entries unchanged. Its matrix is now J_t, with diagonal $(0, 1, 1, 1, 1, 1, 1, t)$ and $t \in \{-2, 0, 2\}$. The first seven rows give the same polynomial recurrence for all signed multiplication matrices. Thus $\dim X_1$ generates K.

If $t = -2$, the bottom-right entry of $M_2 = J_t^2 - J_t - I$ is 6, exceeding the original multiplicity $N_{27}^7 = 2$. If $t = 0$, then

$$\det(xI - J_0) = (x - 1)(x + 1)(x^2 - 2x - 1)(x^4 - 4x^3 + 2x^2 + 4x - 1).$$

The degree of $\dim X_1$ is at most four, contrary to $[K : \mathbb{Q}] = 8$. Finally, if $t = 2$, the recurrence makes every signed multiplication matrix equal to the original nonnegative one. The negative lift multiplicities all vanish, so the eight chosen lifts form a spherical fusion subcategory equivalent to the original categorification. This contradicts the spherical exclusion already proved. □

The source-bound certificate verifies all 512 dimension-character products in $\mathbb{Q}(\zeta_{32})$, all eight exact Fourier norms, and the three displayed matrix branches. The argument is independent of any construction of the corresponding folded quantum-group datum.

21.1. **A central-lift obstruction for the $2D2$ rules.**

Proposition 21.2 (No central invertible lift for the $2D2$ rules)**.** *Let $\mathcal{Q}$ be a pseudo-unitary fusion category with simple objects $1, f, r, s$ and fusion rules*

$$f^2 = 1, \qquad fr = rf = s, \qquad r^2 = s^2 = 1 + 2r + 2s, \qquad rs = sr = f + 2r + 2s.$$

There is no invertible object $z \in Z(\mathcal{Q})$ whose underlying object is f.

Proof. Put $d = 2 + \sqrt{5}$, so $d^2 = 1 + 4d$. The displayed rules imply $\mathcal{Q}_{\mathrm{ad}} = \mathcal{Q}$. Consequently the only central lift of the tensor unit is the tensor unit: such lifts are tensor automorphisms of the identity, equivalently characters of the universal grading group. Thus a hypothetical z satisfies $z^2 \cong 1$. In its pointed braided subcategory the positive ribbon twist $q = \theta_z$ belongs to $\{1, -1, i, -i\}$.

If $q = 1$, then $1 \oplus z$ is the regular algebra of a central Tannakian $\mathrm{Rep}(C_2)$. Its module category has two simple free orbits, represented by $1, r$, and their rule is $\bar{r}^2 = 1 + 4\bar{r}$. This contradicts the rank-two fusion-category classification [23].

Suppose $q = \pm i$. Write I for induction to the center. The formal-codegree induction formula, as recalled in [30, Appendix A.2], gives the distinct constituents of $I(1)$ dimensions

$$1, \quad d^2, \quad 1 + 2d, \quad 1 + 2d.$$

Every constituent has twist 1. Each has tensor-unit multiplicity one after forgetting, and none contains f, because $FI(1)$ has no f-summand. The projection formula gives $I(f) \cong z \otimes I(1)$, disjoint from $I(1)$. Since $z^2 = 1$, the monodromy of z with any simple central object is ± 1. Hence every constituent of $I(f)$ has twist $\pm i$.

In $I(r)\oplus I(s)$ the constituents belonging to $I(1)$ have total dimension

$$K = 4d^2 + 4(1+2d) = 8 + 24d.$$

Indeed their total multiplicities are, respectively, $0, 4, 2, 2$. The constituents belonging to $I(f)$ have the same total dimension. Since

$$\dim\bigl(I(r)\oplus I(s)\bigr) = 2d(4+8d) = 16 + 72d,$$

the other constituents have total dimension $24d$. The first-indicator identity gives zero twist trace on $I(r)\oplus I(s)$. Its $I(1)$ part has real trace $8+24d$, its $I(f)$ part has purely imaginary trace, and the absolute real trace of the remaining part is at most $24d$. This is impossible.

Finally suppose $q=-1$. Let $\mathcal{I}$ be an Ising category with fermion ψ and noninvertible σ. The object $z\boxtimes\psi$ is a central boson in $\mathcal{Q}\boxtimes\mathcal{I}$. Condensing its regular algebra gives a pseudo-unitary fusion category $\mathcal{D}$ of rank six. All twelve simples before condensation have free orbits. Representatives give objects $1, f, u, r, s, v$ of dimensions

$$1, 1, \sqrt{2}, d, d, \sqrt{2}d,$$

with the above rules on $1, f, r, s$ and additional rules

$$fu = u, \quad fv = v, \quad u^2 = 1 + f, \quad ur = us = v,$$

$$uv = r + s, \quad rv = sv = u + 4v, \quad v^2 = 1 + f + 4r + 4s.$$

We prove directly that these rules have no pseudo-unitary categorification.

Put $D = \dim(\mathcal{D}) = 8+16d$ and $T = Dd = 16+72d$. The six formal codegrees show that $I(1)$ has two constituents of each dimension $1, d^2, 2+4d$, all of twist 1, each with tensor-unit multiplicity one. Positivity and linear independence over $\mathbb{Q}(\sqrt{5})$ show that their restrictions have no u or v summands. The two constituents of dimension d^2 restrict to

$$1 + a_j r + (4-a_j)s \qquad (0 \le a_j \le 4),$$

and the two of dimension $2+4d$ restrict to

$$1 + f + b_j r + (4-b_j)s \qquad (0 \le b_j \le 4).$$

The r-coefficient of $FI(1)$ is 8, so if $a = a_1 + a_2$ then $b_1 + b_2 = 8 - a$. The total dimension K of the twist-one summands of $I(r)$ supplied by $I(1)$ is therefore

$$K = ad^2 + (8-a)(2+4d) = 16 - a + 32d \ge 8 + 32d.$$

Let $R = T - K$ be the dimension of the other summands. The first two indicator identities [59], with $\nu_2(r) \le 1$, imply

$$\sum_j w_j \cos t_j = -K, \qquad \sum_j w_j \cos(2t_j) = D\nu_2(r) - K \le D - K, \qquad \sum_j w_j = R,$$

where $w_j > 0$ and e^{it_j} are their dimensions and twists. If $R = 0$ the first equation is already impossible; otherwise Cauchy–Schwarz gives

$$2K^2 \le R(R + D - K), \qquad 0 \le T(T+D) - K(3T+D).$$

But the last expression is at most

$$T(T+D) - (8+32d)(3T+D) = -64 - 512d - 1088d^2 < 0.$$

This final contradiction excludes $q=-1$. □

Remark 21.3. The published Galois pseudo-unitarity reduction [25, Lemma 3.1] for categories having two Frobenius–Perron dimensions upgrades this statement to arbitrary complex categorifications of the displayed ring: a central lift survives Galois conjugation. This uses the corrected published formulation “Galois conjugate to a pseudo-unitary category”, not a claim that every categorification is itself pseudo-unitary.

Remark 21.4 (Scope of the inner-conjugation construction). Let $\mathcal{P} = \mathcal{Q}_{\mathrm{pt}}$ and $\mathcal{D} = \mathcal{Z}_{\mathcal{P}}(\mathcal{Q})$. By [47, Remark 2.4(i)], this relative center is Morita equivalent to $\mathcal{P} \boxtimes \mathcal{Q}^{\mathrm{rev}}$. Its invertibles are precisely those of $\mathcal{Z}(\mathcal{P})$, whose group is $C_2 \times C_2$ for either associator on $\mathcal{P} \simeq \mathrm{Vec}_{C_2}^{\omega}$. Thus this natural inner-conjugation construction cannot realize the C_4-based rings $R8_1_9$ or $R8_1_18$; no indicator assumption is needed.

The displayed rank-four rules have trivial universal grading, so the theorem implies $\mathrm{Inv}\,\mathcal{Z}(\mathcal{Q}) = 1$. Morita invariance then gives $|\mathrm{Inv}\,\mathcal{Z}(\mathcal{D})| = 4$. If $\mathcal{D}$ had the rules $R8_1_49$, its universal grading would also be trivial. The forgetful map on invertibles would therefore be injective, since its kernel is the group of tensor automorphisms of the identity, and hence bijective. Every invertible would have a central lift, contradicting the noncommuting invertible action in that ring. This rules out the proposed construction only. General exclusions are proved independently in Theorems 13.7 and 13.9 and Proposition 8.3.

22. Reproduction and mathematical limits

Run `python3 tools/verify_current_publication.py` from the repository root, and repeat with `-O`. The verifier checks the published documents and their website copies, the source identities, the current decision ledger and the accepted finite certificate applications, and the complete current unresolved inventory. Run `python3 tools/verify_rigidity_20261017.py` for the inner-square and associated theoretical evidence, and repeat that command with `-O`. The current-publication verifier's `-historical` option also replays the full chain of source-bound certificate applications in isolated inputs. The exact replay instructions and the scope of every recorded run appear in `docs/VERIFY.md`.

The historical rank-eight multiplicity-two full-certificate audit covers all 973 input tensors, including the 75 signed proof trees and four rank-seven subring proof transfers used by the corresponding base certificates. The theoretical applications and positive tensor comparisons are checked separately. All finite checks use integer or algebraic arithmetic. They validate their specified equations and source bindings; they do not independently prove categorical transport identities, cited theorems or census completeness. A timeout or an unfinished branch never constitutes an exclusion.

The cyclic induction and center STS arguments in Theorems 13.7 and 13.9, together with Proposition 8.3, complete the multiplicity-one classification. At multiplicity two the conclusions are 848 general exclusions and 29 unitary constructions. Theorem 14.1 excludes $R8_2_24$ by combining the saturated indicator signs, the actual inner square supplied by its invertible object, the common pivotal chart and tensor-functor rigidity. The source and weight checks are recorded in `evidence/rigidity_20261017/rank8/`.

The finite applications are conditional on the stated mathematical arguments, cited theorems and source census completeness. The multiplicity-three input is checked as described in Section 3; exhaustive enumeration is not independently rerun. This work does not classify categories up to tensor equivalence or claim that no further criterion is needed. Noncommutative Extended Haagerup models at rank eight have multiplicities four and seven [43]; they are genuine models outside the present census bound.

AI disclosure and verification status. GPT-6 Astra and GPT-6.1 Sol assisted with literature exploration, mathematical research, code development and manuscript preparation. The results, proofs and accompanying data are AI-generated and have not been independently verified by a human expert or formalized in Lean. They should be used with caution. Computational replay provides evidence for its explicit finite checks and does not replace mathematical review.

References

[1] G. Vercleyen and J. Slingerland, *On Low-Rank Multiplicity-Free Complex Fusion Categories*, *Commun. Math. Phys.* **407**, 231 (2026), DOI:10.1007/s00220-026-05705-2; https://anyonwiki.github.io/pages/Lists/losmffr.html.

[2] S. X. Cui, C. Galindo, J. Y. Plavnik and Z. Wang, *On Gauging Symmetry of Modular Categories*, *Commun. Math. Phys.* **348** (2016), 1043–1064; arXiv:1510.03475.

[3] P. Etingof, D. Nikshych and V. Ostrik, *Weakly group-theoretical and solvable fusion categories*, arXiv:0809.3031v2, Theorem 1.6 and Proposition 4.4.

[4] V. Drinfeld, S. Gelaki, D. Nikshych and V. Ostrik, *On braided fusion categories I*, arXiv:0906.0620v3, Theorem 4.18 and Proposition 4.19.

[5] V. Ostrik, *Module categories over the Drinfeld double of a finite group*, Int. Math. Res. Not. (2003), 1507–1520; corrected version arXiv:math/0202130v2.

[6] S. Burciu and S. Natale, *Fusion rules of equivariantizations of fusion categories*, J. Math. Phys. **54** (2013), 013511; corrected version arXiv:1206.6625v2.

[7] P. Etingof, D. Nikshych and V. Ostrik, *Fusion categories and homotopy theory*, Quantum Topol. **1** (2010), 209–273; arXiv:0909.3140.

[8] M. Izumi, *The classification of* 3^n *subfactors and related fusion categories*, Quantum Topol. **9** (2018), 473–562, DOI:10.4171/QT/113.

[9] P. Grossman, M. Izumi and N. Snyder, *The Asaeda–Haagerup fusion categories*, J. Reine Angew. Math. **743** (2018), 261–305; arXiv:1501.07324.

[10] P. Etingof and S. Gelaki, *Semisimple Hopf algebras of dimension pq are trivial*, J. Algebra **210** (1998), 664–669; arXiv:math/9801129.

[11] S. Palcoux, *Fusion Ring Census*, exact tensor dataset and verification records, `https://github.com/sebastienpalcoux/fusion-ring-census`, commit `b86eb6dc6eeeafe5fc5f6906d7b074e0322e9726` (2026).

[12] *A short classification of complex rank-two Grothendieck rings*, manuscript (2026).

[13] *Galois-averaged indicators and sphericality in rank three*, manuscript (2026).

[14] *Grothendieck rings of general fusion categories of rank four*, manuscript and exact-certificate supplement (2026).

[15] *Rank-four Grothendieck rings through multiplicity 128*, finite-classification manuscript and exact-certificate supplement (2026).

[16] *Rank-five Grothendieck rings through multiplicity twenty*, manuscript and exact-certificate supplement (2026).

[17] *Rank-six fusion rings: categorification criteria and models through multiplicity nine*, manuscript and exact-certificate supplement (2026).

[18] *Rank-seven Grothendieck rings: a complete classification through multiplicity five*, manuscript and exact-certificate supplement (2026).

[19] M. A. Alekseyev, W. Bruns, J. Dong and S. Palcoux, *Classifying integral Grothendieck rings up to rank 5 and beyond*, accepted in *Letters in Mathematical Physics*; arXiv:2507.07023 (2025).

[20] M. A. Alekseyev, W. Bruns, J. Dong and S. Palcoux, *Classifying integral Grothendieck rings up to rank 5 and beyond*, archived full-data preprint, Appendix C.3.2 and Lemmas 4.2–4.3, `https://bimsa.net/doc/publication/14753.pdf` (2025).

[21] P. Etingof, D. Nikshych and V. Ostrik, *On fusion categories*, Ann. of Math. **162** (2005), 581–642.

[22] P. Etingof, S. Gelaki, D. Nikshych and V. Ostrik, *Tensor Categories*, AMS Mathematical Surveys and Monographs 205 (2015).

[23] V. Ostrik, *Fusion categories of rank 2*, Math. Res. Lett. **10** (2003), 177–183.

[24] V. Ostrik, *On formal codegrees of fusion categories*, Math. Res. Lett. **16** (2009), 895–901.

[25] A. Schopieray, *Categorification of integral group rings extended by one dimension*, J. London Math. Soc. **108** (2023), 1617–1641, Lemma 3.1 and Proposition 3.2.

[26] A. Schopieray, *Norm, trace, and formal codegrees of fusion categories*, J. Algebra **568** (2021), 362–385, Theorem 4.4 and Corollary 4.3.

[27] A. Schopieray, *Greedy fusion*, arXiv:2607.13126v1 (2026), Proposition 1 and Corollary 2.

[28] Z. Liu, S. Palcoux and Y. Ren, *Classification of Grothendieck rings of complex fusion categories of multiplicity one up to rank six*, Lett. Math. Phys. **112** (2022), article 54, Theorem 2.14.

[29] A. Kirillov Jr. and V. Ostrik, *On a q-analogue of the McKay correspondence and the ADE classification of* $\widehat{\mathfrak{sl}}_2$ *conformal field theories*, Adv. Math. **171** (2002), 183–227.

[30] C. Edie-Michell, M. Izumi and D. Penneys, *Classification of* $\mathbb{Z}/2\mathbb{Z}$*-quadratic unitary fusion categories*, Quantum Topol. **16** (2025), 493–568; DOI:10.4171/QT/201.

[31] C. Jones, S. Morrison, D. Penneys and J. Plavnik, *Extension theory for braided-enriched fusion categories*, arXiv:1910.03178, Section 6.2.

[32] C. Galindo, *On braided and ribbon unitary fusion categories*, arXiv:1209.2022v3, corrected Remark 3.1 (December 2024).

[33] C. Edie-Michell, *The Brauer–Picard groups of fusion categories coming from the ADE subfactors*, Int. J. Math. **29** (2018), 1850036; arXiv:1709.04721.

[34] D. Reutter, *Uniqueness of unitary structure for unitarizable fusion categories*, arXiv:1906.09710, Theorems 1–2.

[35] Z. Yu and Q. Zhang, *Realizing modular data from centers of near-group categories*, arXiv:2412.20763v2; DOI:10.1007/s10468-025-10370-5.
[36] M. Izumi, *A Cuntz algebra approach to the classification of near-group categories*, arXiv:1512.04288v2, Theorem 12.5.
[37] P. G. Budinski, *Exotic Fusion Categories and Their Modular Data*, master's thesis, University of Alberta (2021); DOI:10.7939/r3-7p7k-fg18.
[38] J. C. Y. Teo, T. L. Hughes and E. Fradkin, *Theory of twist liquids: gauging an anyonic symmetry*, Ann. Phys. **360** (2015), 349–445; arXiv:1503.06812.
[39] C. Galindo, S.-M. Hong and E. C. Rowell, *Generalized and quasi-localizations of braid group representations*, Int. Math. Res. Not. (2013), 693–731; arXiv:1105.5048, Theorem 5.20.
[40] M. Müger, *Galois theory for braided tensor categories and the modular closure*, Adv. Math. **150** (2000), 151–201; arXiv:math/9812040.
[41] D. Tambara and S. Yamagami, *Tensor categories with fusion rules of self-duality for finite abelian groups*, J. Algebra **209** (1998), 692–707.
[42] J.-P. Serre, *Linear Representations of Finite Groups*, Springer Graduate Texts in Mathematics 42 (1977), Chapter 5.
[43] P. Grossman, S. Morrison, D. Penneys, E. Peters and N. Snyder, *The Extended Haagerup fusion categories*, Ann. Sci. Ecole Norm. Sup. **56** (2023), 589–664; arXiv:1810.06076, Theorem 1.1 and Sections 2.2–2.4.
[44] Y. Kashina, *Classification of semisimple Hopf algebras of dimension 16*, J. Algebra **232** (2000), 617–663; arXiv:math/0004114.
[45] D. Jordan and E. Larson, *On the classification of certain fusion categories*, J. Noncommut. Geom. **3** (2009), 481–499; arXiv:0812.1603.
[46] S. Gelaki and D. Naidu, *Some properties of group-theoretical categories*, J. Algebra **322** (2009), 2631–2641; arXiv:0709.4326.
[47] S. Gelaki, D. Naidu and D. Nikshych, *Centers of graded fusion categories*, Algebra Number Theory **3** (2009), 959–990; arXiv:0905.3117.
[48] S. Morrison, E. Peters and N. Snyder, *Skein theory for the D_{2n} planar algebras*, J. Pure Appl. Algebra **214** (2010), 117–139; arXiv:0808.0764.
[49] S. Morrison, E. Peters and N. Snyder, *Knot polynomial identities and quantum group coincidences*, Quantum Topol. **2** (2011), 101–156; arXiv:1003.0022.
[50] C. Edie-Michell, *Auto-equivalences of the modular tensor categories of type A, B, C and G*, arXiv:2002.03220v2.
[51] M. Bischoff, *The rank of G-crossed braided extensions of modular tensor categories*, Contemp. Math. **747** (2020), 115–120; arXiv:1807.06131v2.
[52] A. Davydov and I. Runkel, *$\mathbb{Z}/2\mathbb{Z}$-extensions of Hopf algebra module categories by their base categories*, Adv. Math. **247** (2013), 192–265; arXiv:1207.3611v2.
[53] D. E. Evans and T. Gannon, *The exoticness and realisability of twisted Haagerup–Izumi modular data*, Comm. Math. Phys. **307** (2011), 463–512; arXiv:1006.1326.
[54] A. Schopieray, *Irrational braided generalized near-groups*, Canad. Math. Bull. (2023); DOI:10.4153/S0008439523000334.
[55] P. Grossman, *Fusion categories associated to subfactors with index $3+\sqrt{5}$*, arXiv:1612.07185.
[56] H. K. Larson, *Pseudo-unitary non-self-dual fusion categories of rank 4*, arXiv:1401.1879v2.
[57] D. Nikshych and B. Riepel, *Categorical Lagrangian Grassmannians and Brauer–Picard groups of pointed fusion categories*, arXiv:1309.5026v2.
[58] C. Galindo, *Clifford theory for graded fusion categories*, arXiv:1010.5283v3, Theorem 1.3.
[59] S.-H. Ng and P. Schauenburg, *Frobenius–Schur indicators and exponents of spherical categories*, Adv. Math. **211** (2007), 34–71; arXiv:math/0601012v3.
[60] S.-H. Ng and P. Schauenburg, *Congruence subgroups and generalized Frobenius–Schur indicators*, Comm. Math. Phys. **300** (2010), 1–46; arXiv:0806.2493v3.
[61] P. Bruillard, S.-H. Ng, E. C. Rowell and Z. Wang, *Rank-finiteness for modular categories*, arXiv:1310.7050v3, the Cauchy theorem for spherical fusion categories.
[62] Z. Liu, S. Palcoux, Y. Ren and G. Vercleyen, *Triangular prism equations and categorification*, arXiv:2203.06522v5, Theorem 3.9.
[63] A. Schopieray, *The dimension block of a spherical fusion category*, arXiv:2609.35287v1 (28 September 2026), Lemma 4.1 and Theorem 5.1.
[64] P. Etingof, S. Gelaki and V. Ostrik, *Classification of fusion categories of dimension pq*, Int. Math. Res. Not. **2004** (2004), 3041–3056; arXiv:math/0304194v2, Lemma 6.2 and Theorems 6.3 and 7.1.
[65] J. Dong and H. Sun, *Structure, examples and classification for generalized near-group fusion categories*, J. Algebra **568** (2021), 386–407; arXiv:2004.08735v1, Propositions 3.3 and 3.6.
[66] P. Grossman, M. Izumi and N. Snyder, *Graded extensions of generalized Haagerup categories*, arXiv:2201.11901v1, Section 3 and Theorem 4.1.

[67] P. Grossman and M. Izumi, *Drinfeld centers of fusion categories arising from generalized Haagerup subfactors*, Quantum Topol. **13** (2022), 593–668; DOI:10.4171/QT/167, Lemma 3.5, Theorem 3.9 and Example 4.5.
[68] G. Schaumann, *Traces on module categories over fusion categories*, arXiv:1206.5716v3, Theorem 4.5 and Propositions 5.8, 5.10.
[69] S. Morrison and D. Penneys, *2-supertransitive subfactors at index* $3+\sqrt{5}$, arXiv:1406.3401v1, Lemmas 3.2, 3.4 and Theorems 7.8, 7.9.
[70] D. E. Evans and T. Gannon, *Near-group fusion categories and their doubles*, Adv. Math. **255** (2014), 586–640; DOI:10.1016/j.aim.2013.12.014; arXiv:1208.1500, Proposition 6, Table 2, Proposition 7(b), and the finite verification following Conjecture 2.
[71] P. Grossman and M. Izumi, *Infinite families of potential modular data related to quadratic categories*, arXiv:1906.07397v2, Section 3.2.3.1.
[72] P. Bonderson, E. C. Rowell and Z. Wang, *On realizing modular data*, J. Math. Sci. Univ. Tokyo **27** (2020), 65–79; arXiv:1910.07061, Section 2.4.
[73] Z. Feng, S. Ming and E. C. Rowell, *Reconstructing braided subcategories of* $SU(N)_k$, J. Algebra **635** (2023), 436–458; arXiv:2202.00121v2, Theorem 5.1.
[74] V. Ostrik, *Pivotal fusion categories of rank 3*, arXiv:1309.4822.
[75] V. Ostrik, *Module categories, weak Hopf algebras and modular invariants*, Transform. Groups **8** (2003), 177–206; arXiv:math/0111139, Theorem 2 and Remark 12.
[76] C. Edie-Michell, *A complete classification of unitary fusion categories tensor generated by an object of dimension* $(1+\sqrt{5})/2$, arXiv:1904.08909v2, Definition 2.5.
[77] J. Dong, *Classification of certain weakly integral fusion categories*, arXiv:2103.04553v3, Theorem 1.2 and Corollary 4.16.
[78] F. Calegari, S. Morrison and N. Snyder, *Cyclotomic integers, fusion categories, and subfactors*, Comm. Math. Phys. **303** (2011), 845–896, Appendix A by V. Ostrik, Propositions A.4.1 and A.6.1 and Section A.5.
[79] D. Naidu, *Categorical Morita equivalence for group-theoretical categories*, Comm. Algebra **35** (2007), 3544–3565; arXiv:math/0605530, Theorem 3.4.
[80] A. Davydov, M. Müger, D. Nikshych and V. Ostrik, *The Witt group of non-degenerate braided fusion categories*, J. Reine Angew. Math. **677** (2013), 135–177; arXiv:1009.2117v2, Section 3.3.
[81] B. Gui, *Q-systems and extensions of completely unitary vertex operator algebras*, arXiv:1908.03484v3, Theorems 2.21 and 2.28, Corollary 3.32.
[82] S. Morrison and D. Penneys, *Constructing spoke subfactors using the jellyfish algorithm*, arXiv:1208.3637v2, Section 5.1 and Theorem 5.5.
[83] P. Grossman, D. Jordan and N. Snyder, *Cyclic extensions of fusion categories via the Brauer–Picard groupoid*, arXiv:1211.6414v2, Theorem 4.9 and Remarks 4.10–4.11.
[84] H. Cartan and S. Eilenberg, *Homological Algebra*, Princeton University Press (1956), Chapter XII, Section 10.
[85] P. Grossman and N. Snyder, *The Brauer–Picard group of the Asaeda–Haagerup fusion categories*, Trans. Amer. Math. Soc. **368** (2016), 2289–2331; arXiv:1202.4396v2.
[86] T.-C. Huang, *Cyclic Haagerup–Izumi fusion categories at every odd order*, arXiv:2609.15986v1 (2026).
[87] T. Gannon, A. Schopieray and H. Yadav, *On Haagerup–Izumi fusion categories*, arXiv:2609.25185v1 (2026).
[88] P. Grossman and N. Snyder, *Quantum subgroups of the Haagerup fusion categories*, Comm. Math. Phys. **311** (2012), 617–643; arXiv:1102.2631v2, Theorem 1.5.