

The category of chord diagrams and its generic limit

33rd Summer Conference on Topology and its Applications

Sebastian Zając
(joint work with Wiesław Kubis)

Cardinal Stefan Wyszyński University in Warsaw

20.07.2018

1 Motivation

2 The category of chord diagrams


3 Fraïsse limit

Chord diagrams may be used to represent the inter-relationships between some objects.

What is a chord diagram ?

Let's consider a collection of $b \geq 1$ (pairwise, disjoint, oriented and labeled) intervals lying in the real line. We call them backbones. By a chord diagram on backbones we understand a collection of $n \geq 0$ semi-circles (called chords) whose endpoints lie at distinct interior points of backbones.

Graphical representation



T. Doslic "Secondary structures and some related combinatorial objects", DOI: <https://doi.org/10.5592/CO/CCD.2016.02>


Where can you find it ?

Physics



“Physics is like sex: sure, it may give some practical results, but that’s not why we do it.”

– Richard Feynman



Category theory is like sex: it may not give any practical results, but that’s not why we do it - Sebastian Zajac

Math

- Matrix Field Theory (0-dim quantum field theory),
- geometry, topology, representation theory,
- moduli spaces of Riemann surfaces.

- ① A natural number n is the set $\{0, 1, \dots, n - 1\}$.
- ② Recall that a *linear ordering* on a set S is an irreflexive transitive relation $<$ satisfying $x < y$ or $y < x$ for every $x \neq y$ in S . In that case $\langle S, < \rangle$ is called a *linearly ordered set*.
- ③ If $\langle S, <^S \rangle$ and $\langle T, <^T \rangle$ are linearly ordered sets then a mapping $f: S \rightarrow T$ is called *increasing* if $f(x) <^T f(y)$ whenever $x <^S y$.
- ④ If A, B are subsets of a linearly ordered set $\langle S, < \rangle$ then we write $A < B$ if $a < b$ holds for every $a \in A$ and $b \in B$.
- ⑤ we denote by $\mathcal{P}(S)$ - power-set of S (the set of all subsets of S).
- ⑥ Given a natural number k , we denote by $[S]^k$ the family of all k -element subsets of S .
- ⑦ A family $\mathcal{A} \subseteq \mathcal{P}(S)$ is called a *partition* of S if $S = \bigcup \mathcal{A}$ and $A_0 \cap A_1 = \emptyset$ for every $A_0 \neq A_1$ in \mathcal{A} .
- ⑧ A *graph* (more formally: a simple undirected graph) is a structure of the form $\langle S, E \rangle$, where $E \subseteq [S]^2$.

The category of chord diagrams

We fix three positive integers k , ℓ , and m and a multifunction

$$\varphi: \ell \times \ell \rightarrow m$$

The objects of $\mathfrak{C}_{k,\varphi}$ are structures of the form

$$\mathbb{S} = \langle S, <^S, \{B_i^S\}_{i < k}, \{N_i^S\}_{i < \ell}, \{E_i\}_{i < m} \rangle,$$

where:

- (D1) $\langle S, <^S \rangle$ is a finite linearly ordered set.
- (D2) $\{B_i^S\}_{i < k}$ and $\{N_i^S\}_{i < \ell}$ are partitions of S .
- (D3) $B_{i_0} < B_{i_1}$ whenever $i_0 < i_1 < k$.
- (D4) $\langle S, E_i \rangle$ is a graph for every $i < m$.
- (D5) $E_{i_0} \cap E_{i_1} = \emptyset$ whenever $i_0 \neq i_1$.
- (D6) If $x \in N_{i_0}$, $y \in N_{i_1}$ and $\langle x, y \rangle \in E_j$, then $j \in \varphi(i_0, i_1)$.

The sets B_i are called *backbones*, while the sets N_i are *types of nodes* and E_i are *types of edges*.

A $\mathfrak{C}_{k,\varphi}$ -morphism from \mathbb{S} to $\mathbb{T} = \langle T, <^T, \{B_i^T\}_{i < k}, \{N_i^T\}_{i < \ell}, \{E_i\}_{i < m} \rangle$ is a mapping $f: S \rightarrow T$ that preserves the linear orderings (that is, $x <^S y \implies f(x) <^T f(y)$) and satisfies for every $x, y \in S$:

$$(M1) \quad f(x) \in B_i^T \iff x \in B_i^S$$

$$(M2) \quad f(x) \in N_i^T \iff x \in N_i^S$$


$$(M3) \quad \langle f(x), f(y) \rangle \in E_i^T \iff \langle x, y \rangle \in E_i^S.$$

Informally, a $\mathfrak{C}_{k,\varphi}$ -arrow is a mapping that preserves the structure of \mathbb{S} , “adding” new vertices and new edges of various types.


In the language of model theory, $\mathfrak{C}_{k,\varphi}$ -arrows are called *embeddings*.

It is clear that $\mathfrak{C}_{k,\varphi}$ forms a category.


Identity



Morphisms - embeddings



Composition of morphisms



The amalgamation property

Given


$$f : S \rightarrow T, \quad g : S \rightarrow K,$$

there should exist an object W and embeddings $f' : T \rightarrow W$ and $g' : K \rightarrow W$ satisfying

$$f' \circ f = g' \circ g$$

$$\begin{array}{ccc} K & \xrightarrow{g} & W \\ g' \uparrow & & \uparrow f' \\ S & \xrightarrow{f} & T \end{array}$$

The chord category as a Fraïssé category example



Universal chord diagram

In one backbone case Fraïssé limit is the Rado graph with a linear ordering isomorphic to the rationals.

