

MATH 345 Skeleton Notes

Unit 3: Beyond linear maps: local linearization

Section 3.3: One-parameter maps and changes

© 2024–2026 Jacob Denson, Kaitlyn Phillipson, Sebastien Roch. Except where otherwise noted, this work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

Vector-valued functions of one variable

Basic definitions

A map $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^n$ takes one scalar parameter and outputs a vector. It traces a curve.

Anatomy of a linear map lines are the special case $\mathbf{r}(t) = \mathbf{x}_0 + t\mathbf{d}$. Indeed, we already encountered parameterizations of curves when discussing parametric equations for straight lines, as in Parametric equation for a line.

Activity

The line L_1 passes through point $(3, -2, 5)$ and is parallel to the vector $\mathbf{v} = (2, -3, 4)$.

Find a parameterization of the line L_1 .

Activity

The line L_1 passes through point $(3, -2, 5)$ and is parallel to the vector $\mathbf{v} = (2, -3, 4)$.

Another line L_2 passes through the point $(1, 3, -1)$ and is perpendicular to L_1 . If L_2 also lies in the plane $z = -1$, find a parameterization of L_2 .

The parameterizations of straight lines given above are given by vector-valued functions whose components are linear functions. But we can also parameterize curves using nonlinear functions.

Example As we vary t over its allowed values, the vector-valued function $\mathbf{r}(t)$ traces a curve in $\mathbb{R}^n$. We will not graph functions by hand in this course. For example, consider

$$\mathbf{r}(t) = 4 \cos t \mathbf{i} + 4 \sin t \mathbf{j} + t \mathbf{k}, \quad 0 \leq t \leq 4\pi.$$

Then the curve it traces is illustrated below.

Activity

Recall the vectors $\mathbf{i}$ and $\mathbf{j}$ in $\mathbb{R}^2$ from Standard basis. Consider the vector-valued function $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^3$ defined by the equation

$$\mathbf{r}(t) = (t^2 - 3t)\mathbf{i} + (4t + 1)\mathbf{j},$$

i.e., the equation

$$\mathbf{r}(t) = \begin{bmatrix} t^2 - 3t \\ 4t + 1 \end{bmatrix}.$$

Evaluate $\mathbf{r}(0)$.

Activity

Recall the vectors $\mathbf{i}$ and $\mathbf{j}$ in $\mathbb{R}^2$ from Standard basis. Consider the vector-valued function $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^3$ defined by the equation

$$\mathbf{r}(t) = (t^2 - 3t)\mathbf{i} + (4t + 1)\mathbf{j},$$

i.e., the equation

$$\mathbf{r}(t) = \begin{bmatrix} t^2 - 3t \\ 4t + 1 \end{bmatrix}.$$

Evaluate $\mathbf{r}(1)$.

Activity

Recall the vectors $\mathbf{i}$ and $\mathbf{j}$ in $\mathbb{R}^2$ from Standard basis. Consider the vector-valued function $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^3$ defined by the equation

$$\mathbf{r}(t) = (t^2 - 3t)\mathbf{i} + (4t + 1)\mathbf{j},$$

i.e., the equation

$$\mathbf{r}(t) = \begin{bmatrix} t^2 - 3t \\ 4t + 1 \end{bmatrix}.$$

Evaluate $\mathbf{r}(-4)$.

We use parameterizations mainly to describe position, velocity, and tangent direction along a curve.

Limits and continuity

Definition: Limit of a Vector-Valued Function A vector-valued function $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^n$ approaches the $\mathbf{v} \in \mathbb{R}^n$ as t approaches a , written

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{v},$$

provided

$$\lim_{t \rightarrow a} \|\mathbf{r}(t) - \mathbf{v}\| = 0.$$

This is the limit of a scalar-valued function, and can thus be evaluated using the methods of single-variable calculus.

In practice, calculating and simplifying the expression $\|\mathbf{r}(t) - \mathbf{v}\|$ in order to evaluate the limit

$$\lim_{t \rightarrow a} \|\mathbf{r}(t) - \mathbf{v}\| = 0$$

can be quite cumbersome. Instead, the following theorem allows us to evaluate limits much more easily.

Theorem: Limit of a Vector-Valued Function Let $\mathbf{r}(t) : \mathbb{R} \rightarrow \mathbb{R}^n$ be a function with component functions $r_1, \dots, r_n : \mathbb{R} \rightarrow \mathbb{R}$. Then if

$$\lim_{t \rightarrow a} r_1(t) = v_1 \quad \cdots \quad \lim_{t \rightarrow a} r_n(t) = v_n,$$

and we define the vector $\mathbf{v} = (v_1, \dots, v_n)$, then

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{v}.$$

Equivalently,

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \begin{bmatrix} \lim_{t \rightarrow a} r_1(t) \\ \vdots \\ \lim_{t \rightarrow a} r_n(t) \end{bmatrix}.$$

Activity

Calculate $\lim_{t \rightarrow 3} \mathbf{r}(t)$ for the given vector-valued functions $\mathbf{r}(t)$.
The function $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^2$ defined by

$$\mathbf{r}(t) = (t^2 - 3t + 4)\mathbf{i} + (4t + 3)\mathbf{j}.$$

Activity

Calculate $\lim_{t \rightarrow 3} \mathbf{r}(t)$ for the given vector-valued functions $\mathbf{r}(t)$.
The function $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^3$ defined by

$$\mathbf{r}(t) = \left(\frac{2t - 4}{t + 1} \right) \mathbf{i} + \left(\frac{t^2 + 1}{2} \right) \mathbf{j} + (4t - 3)\mathbf{k}.$$

Definition: Continuity of Vector-Valued Functions A vector-valued function $\mathbf{r}(t)$ is at point $t = a$ if

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{r}(a)$$

That is, a vector-valued function is continuous at point $t = a$ when the limit of the function as t approaches a equals the value of the function at a .

Theorem A vector-valued function $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^n$ with component functions $r_1, \dots, r_n$ is continuous at $t = a$ if and only if each of the component functions $r_1, \dots, r_n$ is continuous at $t = a$.

Derivatives

Now we define the derivative of a vector-valued function from $\mathbb{R}$ to $\mathbb{R}^n$.

Definition: Derivative of a Vector-Valued Function The derivative of a function $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^n$ is

$$\frac{d\mathbf{r}}{dt}(t) = \mathbf{r}'(t) = \lim_{\Delta t \rightarrow 0} \frac{\mathbf{r}(t + \Delta t) - \mathbf{r}(t)}{\Delta t},$$

provided the limit exists. If $\mathbf{r}'(t)$ exists, then we say $\mathbf{r}$ is

differentiable

at t . If $\mathbf{r}'(t)$ exists for all t in an open interval

(a, b) , then we say $\mathbf{r}$ is

differentiable on (a, b) . The vector $\mathbf{r}'(t)$ is called the

tangent vector

of the curve described by the function $\mathbf{r}$ at

Remark The derivative $\mathbf{r}' : \mathbb{R} \rightarrow \mathbb{R}^n$ of a differentiable function $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^n$ is *also* a vector-valued function.

As in single variable calculus, we will often have much more efficient methods of calculating the derivative of a function $\mathbf{r}$. But you should be able to work directly from the definition in simple examples.

Activity

Use the definition to calculate the derivative of the function

$$\mathbf{r}(t) = (2t^2 + 3)\mathbf{i} + (5t - 6)\mathbf{j}.$$

Since Limit of a Vector-Valued Function tells us taking limits of functions from $\mathbb{R}$ to $\mathbb{R}^n$ is equivalent to taking the limits of each component function, taking the derivative of such a function is equivalent to taking the derivative of each component function.

Theorem If $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^n$ has components $r_1, \dots, r_n : \mathbb{R} \rightarrow \mathbb{R}$, then $\mathbf{r}$ is differentiable at t if and only if each component function $r_1, \dots, r_n$ is differentiable at t , and

$$\mathbf{r}'(t) = \begin{bmatrix} r'_1(t) \\ \vdots \\ r'_n(t) \end{bmatrix}.$$

Because of the theorem, we can apply all the techniques from single-variable calculus to differentiate functions from $\mathbb{R}$ to $\mathbb{R}^n$.

Activity

Find $\mathbf{r}'(t)$, where

$$\mathbf{r}(t) = e^{2t} \mathbf{i} + \ln(1 + t) \mathbf{j} - (\cos t) \mathbf{k}$$

If $\mathbf{r} : \mathbb{R} \rightarrow \mathbb{R}^n$, then $\mathbf{r}' : \mathbb{R} \rightarrow \mathbb{R}^n$ is also a vector-valued function of the same type, so we can consider the second derivatives $\mathbf{r}''$, i.e., the derivative of the function $\mathbf{r}'$.

Activity

Find $\mathbf{r}''(t)$, where $\mathbf{r}$ is as in the activity.

Differentiation rules

Theorem Suppose $\mathbf{u}, \mathbf{v} : \mathbb{R} \rightarrow \mathbb{R}^n$ are differentiable, c is a scalar, and $a : \mathbb{R} \rightarrow \mathbb{R}$ is a scalar-valued function. Then

Sum Rule

$$(\mathbf{u} + \mathbf{v})'(t) = \mathbf{u}'(t) + \mathbf{v}'(t)$$

Product Rule

$$(c\mathbf{u})'(t) = c\mathbf{u}'(t)$$

Product Rule

$$(a\mathbf{u})'(t) = a'(t)\mathbf{u}(t) + a(t)\mathbf{u}'(t)$$

Dot Product rule

$$(\mathbf{u} \cdot \mathbf{v})'(t) = \mathbf{u}'(t) \cdot \mathbf{v}(t) + \mathbf{u}(t) \cdot \mathbf{v}'(t)$$

Proof We prove the dot product rule, in the two dimensional case. Let

$$\mathbf{u}(t) = u_1(t)\mathbf{i} + u_2(t)\mathbf{j}$$

and let

$$\mathbf{v}(t) = v_1(t)\mathbf{i} + v_2(t)\mathbf{j}.$$

Then

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2.$$

Applying the sum and product rules for functions from $\mathbb{R}$ to $\mathbb{R}$, we find that

$$\begin{aligned} (\mathbf{u} \cdot \mathbf{v})' &= (u_1 v_1)' + (u_2 v_2)' \\ &= (u_1' v_1 + u_1 v_1') + (u_2' v_2 + u_2 v_2') \\ &= (u_1' v_1 + u_2' v_2) + (u_1 v_1' + u_2 v_2') \\ &= \mathbf{u}' \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{v}'. \end{aligned}$$

Activity

Find $(\mathbf{u} \cdot \mathbf{v})'$, where

$$\mathbf{u}(t) = \cos t \mathbf{i} + \sin t \mathbf{j} + t \mathbf{k}$$

and

$$\mathbf{v}(t) = t \mathbf{i} + \ln t \mathbf{j} + \mathbf{k}.$$

Remark Note that in the activity, since $\mathbf{v}$ is only well-defined for $t > 0$, the function $\mathbf{u} \cdot \mathbf{v}$ is also only well-defined for $t > 0$, as is its derivative.