

MATH 345 Skeleton Notes

Unit 4: Orthogonality and least squares

Section 4.4: \$QR\$ factorization and least squares

© 2024–2026 Jacob Denson, Kaitlyn Phillipson, Sebastien Roch. Except where otherwise noted, this work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

QR factorization

Recall that a list of vectors is orthonormal when the vectors have length 1 and are pairwise orthogonal. Also recall the notion of an upper triangular matrix from Triangular and Diagonal Matrices.

Definition Let A be an $m \times n$ matrix with independent columns. A

of A is an $m \times n$ matrix Q , with orthonormal columns, and an invertible upper triangular $n \times n$ matrix R , with positive entries on its diagonal, so that $A = QR$.

Warning: Rectangular Q In a QR factorization of an $m \times n$ matrix with $m > n$, the matrix Q has orthonormal columns, so $Q^T Q = I_n$. But Q is not square, and QQ^T is not usually I_m .

Normal equations are conceptually important, but QR is often a better computational method. If $A = QR$ with Q having orthonormal columns, then

$$\|Ax - b\| = \|QRx - b\|.$$

Orthogonality reduces the least-squares problem to

$$R\hat{x} = Q^T b,$$

which is triangular and easier to solve accurately.

Activity: Reading QR least-squares code

The following code uses QR to solve the same least-squares problem from the Unit 2 target activity.

```
A = np.array([
    [1.0, 0.0],
    [0.0, 1.0],
    [1.0, 1.0],
])
b = np.array([1.0, 2.0, 4.0])

Q, R = np.linalg.qr(A)
x_qr = np.linalg.solve(R, Q.T @ b)
x_lstsq = np.linalg.lstsq(A, b, rcond=None)[0]

Q.T @ Q, x_qr, x_lstsq
```

1. What should $Q.T @ Q$ be close to?
2. Why does the code solve a system with R instead of a system with A ?
3. Why should x_qr and x_lstsq agree?
4. Does this mean $Ax = b$ has an exact solution?

Activity

Let

$$Q = \begin{bmatrix} \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & -\frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{6}} \end{bmatrix}.$$

Check that the columns are orthonormal and compute $Q^T Q$.

Activity

Solve the linear system $Rx = \mathbf{b}$ where

$$R = \begin{bmatrix} 4 & 2 & 7 \\ 0 & 3 & 5 \\ 0 & 0 & 2 \end{bmatrix} \quad \text{and} \quad \mathbf{b} = \begin{bmatrix} 30 \\ 21 \\ 6 \end{bmatrix}$$

The QR factorization is a matrix version of the *Gram-Schmidt orthogonalization process*, which has many applications in many numerical algorithms.

Theorem Every $m \times n$ matrix A with linearly independent columns has a QR factorization $A = QR$.

Proof Let $A = [\mathbf{c}_1 \mathbf{c}_2 \cdots \mathbf{c}_n]$ be the matrix with columns $\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_n$. Apply Gram-Schmidt process to obtain orthogonal vectors $\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_n$ where: $\mathbf{f}_1 = \mathbf{c}_1$ and

$$\mathbf{f}_k = \mathbf{c}_k - \frac{\mathbf{c}_k \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1 - \frac{\mathbf{c}_k \cdot \mathbf{f}_2}{\|\mathbf{f}_2\|^2} \mathbf{f}_2 - \cdots - \frac{\mathbf{c}_k \cdot \mathbf{f}_{k-1}}{\|\mathbf{f}_{k-1}\|^2} \mathbf{f}_{k-1}$$

for $k = 2, 3, \dots, n$.

Let $\mathbf{q}_k = \frac{\mathbf{f}_k}{\|\mathbf{f}_k\|}$ for each k . Then $\mathbf{q}_1, \mathbf{q}_2, \dots, \mathbf{q}_n$ are orthonormal vectors.

We can rewrite each column $\mathbf{c}_k$ as a linear combination of the $\mathbf{q}_i$: $\mathbf{c}_1 = \|\mathbf{f}_1\| \mathbf{q}_1$,

$$\mathbf{c}_2 = (\mathbf{c}_2 \cdot \mathbf{q}_1) \mathbf{q}_1 + \|\mathbf{f}_2\| \mathbf{q}_2$$

$$\mathbf{c}_3 = (\mathbf{c}_3 \cdot \mathbf{q}_1) \mathbf{q}_1 + (\mathbf{c}_3 \cdot \mathbf{q}_2) \mathbf{q}_2 + \|\mathbf{f}_3\| \mathbf{q}_3$$

and so on. This gives us the matrix factorization:

$$\begin{aligned} A &= [\mathbf{c}_1 \mathbf{c}_2 \cdots \mathbf{c}_n] \\ &= [\mathbf{q}_1 \mathbf{q}_2 \cdots \mathbf{q}_n] \begin{bmatrix} \|\mathbf{f}_1\| & \mathbf{c}_2 \cdot \mathbf{q}_1 & \mathbf{c}_3 \cdot \mathbf{q}_1 & \cdots & \mathbf{c}_n \cdot \mathbf{q}_1 \\ 0 & \|\mathbf{f}_2\| & \mathbf{c}_3 \cdot \mathbf{q}_2 & \cdots & \mathbf{c}_n \cdot \mathbf{q}_2 \\ 0 & 0 & \|\mathbf{f}_3\| & \cdots & \mathbf{c}_n \cdot \mathbf{q}_3 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \|\mathbf{f}_n\| \end{bmatrix} \end{aligned}$$

So $A = QR$ where $Q = [\mathbf{q}_1 \mathbf{q}_2 \cdots \mathbf{q}_n]$ has orthonormal columns and R is upper triangular with positive diagonal entries.

Activity

Consider the matrix

$$A = \begin{bmatrix} 3 & 1 \\ 4 & 2 \end{bmatrix}.$$

Find its QR factorization.

Activity

Find the QR factorization of the matrix

$$A = \begin{bmatrix} 6 & 2 \\ 0 & 3 \end{bmatrix}.$$

Theorem Every square, invertible matrix A can be written as $A = QR$, where Q is an orthogonal matrix, and R is an upper triangular matrix with positive diagonal entries.