

MATH 345 Skeleton Notes

Unit 4: Orthogonality and least squares

Section 4.2: Orthogonal projection

© 2024–2026 Jacob Denson, Kaitlyn Phillipson, Sebastien Roch. Except where otherwise noted, this work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

Orthogonal complements and projections

Definition: The orthogonal complement of a subspace If U is a subspace of $\mathbb{R}^n$, the

of U is the subset of $\mathbb{R}^n$ given by

$$U^\perp = \{\mathbf{x} \text{ in } \mathbb{R}^n \mid \mathbf{x} \cdot \mathbf{y} = 0 \text{ for all } \mathbf{y} \text{ in } U\}$$

Theorem The following properties of the orthogonal complement hold:

1. For any subspace U of $\mathbb{R}^n$, $U^\perp$ is a subspace of $\mathbb{R}^n$
2. $\{0\}^\perp = \mathbb{R}^n$ and $(\mathbb{R}^n)^\perp = \{0\}$
3. If $U = \text{span}\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$, then

$$U^\perp = \{\mathbf{x} \text{ in } \mathbb{R}^n \mid \mathbf{x} \cdot \mathbf{x}_i = 0 \text{ for } i = 1, 2, \dots, k\}$$

Activity

Find $U^\perp$ if $U = \text{span}\{(1, 1, 0), (0, 1, 1)\}$ in $\mathbb{R}^3$.

Lemma: Expanding an orthogonal set Let $\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_m\}$ be an orthogonal set in $\mathbb{R}^n$. Given $\mathbf{x}$ in $\mathbb{R}^n$, write

$$\mathbf{f}_{m+1} = \mathbf{x} - \frac{\mathbf{x} \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1 - \frac{\mathbf{x} \cdot \mathbf{f}_2}{\|\mathbf{f}_2\|^2} \mathbf{f}_2 - \dots - \frac{\mathbf{x} \cdot \mathbf{f}_m}{\|\mathbf{f}_m\|^2} \mathbf{f}_m$$

Then:

- $\mathbf{f}_{m+1} \cdot \mathbf{f}_k = 0$ for $k = 1, 2, \dots, m$
- If $\mathbf{x} \notin \text{span}\{\mathbf{f}_1, \dots, \mathbf{f}_m\}$, then $\mathbf{f}_{m+1} \neq \mathbf{0}$ and $\{\mathbf{f}_1, \dots, \mathbf{f}_m, \mathbf{f}_{m+1}\}$ is an orthogonal set.

Proof For convenience, let $t_i = \frac{\mathbf{x} \cdot \mathbf{f}_i}{\|\mathbf{f}_i\|^2}$ for each i . Given $1 \leq k \leq m$, we calculate that

$$\begin{aligned} \mathbf{f}_{m+1} \cdot \mathbf{f}_k &= (\mathbf{x} - t_1 \mathbf{f}_1 - t_2 \mathbf{f}_2 - \dots - t_m \mathbf{f}_m) \cdot \mathbf{f}_k \\ &= \mathbf{x} \cdot \mathbf{f}_k - t_1 (\mathbf{f}_1 \cdot \mathbf{f}_k) - \dots - t_k (\mathbf{f}_k \cdot \mathbf{f}_k) - \dots - t_m (\mathbf{f}_m \cdot \mathbf{f}_k) \end{aligned}$$

Since $\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_m\}$ is an orthogonal set, $\mathbf{f}_i \cdot \mathbf{f}_k = 0$ for all $i \neq k$. So

$$\begin{aligned} \mathbf{f}_{m+1} \cdot \mathbf{f}_k &= \mathbf{x} \cdot \mathbf{f}_k - t_k (\mathbf{f}_k \cdot \mathbf{f}_k) \\ &= \mathbf{x} \cdot \mathbf{f}_k - \frac{\mathbf{x} \cdot \mathbf{f}_k}{\|\mathbf{f}_k\|^2} \cdot \|\mathbf{f}_k\|^2 \\ &= \mathbf{x} \cdot \mathbf{f}_k - \mathbf{x} \cdot \mathbf{f}_k = 0 \end{aligned}$$

This proves the first property of the lemma. To prove the second, we note that if $\mathbf{f}_{m+1}$ was zero, then $\mathbf{x}$ *would* be in the span of the vectors

$\{f_1, \dots, f_m\}$. So if x is not in the span, then f_{m+1} can not be equal to zero. And we have proved f_{m+1} is orthogonal to the vectors $f_1, \dots, f_m$, so that $\{f_1, \dots, f_{m+1}\}$ is an orthogonal set.

Activity

Use Expanding an orthogonal set to extend the orthogonal set $\{f_1, f_2\}$ in $\mathbb{R}^3$ to an orthogonal basis $\{f_1, f_2, f_3\}$, where

$$f_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \quad \text{and} \quad f_2 = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix},$$

using the vector $x = (1, 1, 2)$ as in that lemma.

Definition: The orthogonal projection onto a subspace Let U be a subspace of $\mathbb{R}^n$. Then the

of the vector $x \in \mathbb{R}^n$ onto U is the vector $y = \text{proj}_U(x) \in U$ which is *closest* to x , in the sense that for any $z \in U$ with $z \neq y$, $\|x - y\| < \|x - z\|$.

Remark The following theorem gives us a practical way to calculate orthogonal projections.

Theorem: Projection theorem Let U be a subspace of $\mathbb{R}^n$ with orthogonal basis $\{f_1, \dots, f_m\}$. For any vector $x \in \mathbb{R}^n$, define

$$\mathbf{y} = \frac{\mathbf{x} \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1 + \frac{\mathbf{x} \cdot \mathbf{f}_2}{\|\mathbf{f}_2\|^2} \mathbf{f}_2 + \cdots + \frac{\mathbf{x} \cdot \mathbf{f}_m}{\|\mathbf{f}_m\|^2} \mathbf{f}_m.$$

Then $\mathbf{y} = \text{proj}_U(\mathbf{x})$, and $\mathbf{x} - \text{proj}_U(\mathbf{x}) \in U^\perp$. In particular, if $\mathbf{x}$ is already in U , then $\mathbf{x}$ equals the displayed sum.

Proof If $\{\mathbf{f}_1, \dots, \mathbf{f}_m\}$ is an orthogonal basis for U , and we define

$$\mathbf{y} = \frac{\mathbf{x} \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1 + \frac{\mathbf{x} \cdot \mathbf{f}_2}{\|\mathbf{f}_2\|^2} \mathbf{f}_2 + \cdots + \frac{\mathbf{x} \cdot \mathbf{f}_m}{\|\mathbf{f}_m\|^2} \mathbf{f}_m,$$

then $\mathbf{y}$ is evidently a linear combination of the basis $\{\mathbf{f}_1, \dots, \mathbf{f}_m\}$, so $\mathbf{y} \in U$. And Expanding an orthogonal set states that $\mathbf{x} - \mathbf{y} \in U^\perp$. Thus if $\mathbf{z} \in U$ and $\mathbf{z} \neq \mathbf{y}$, the Pythagorean theorem (Pythagoras' theorem) tells us that

$$\begin{aligned} \|\mathbf{x} - \mathbf{z}\|^2 &= \|(\mathbf{x} - \mathbf{y}) + (\mathbf{y} - \mathbf{z})\|^2 \\ &= \|\mathbf{x} - \mathbf{y}\|^2 + \|\mathbf{y} - \mathbf{z}\|^2. \end{aligned}$$

Since $\mathbf{y} \neq \mathbf{z}$, we find that

$$\|\mathbf{x} - \mathbf{z}\|^2 > \|\mathbf{x} - \mathbf{y}\|^2,$$

and taking square roots gives that

$$\|\mathbf{x} - \mathbf{z}\| > \|\mathbf{x} - \mathbf{y}\|.$$

Thus the vector $\mathbf{y}$ we have defined is equal to $\text{proj}_U(\mathbf{x})$. Since we have already verified that $\mathbf{x} - \mathbf{y} = \mathbf{x} - \text{proj}_U(\mathbf{x})$ is in $U^\perp$, the theorem follows.

Activity

Find $\text{proj}_U \mathbf{x}$ where $U = \text{span}\{(1, 1, 0), (1, -1, 0)\}$ and $\mathbf{x} = (2, 0, 1)$ in $\mathbb{R}^3$.

Activity: Projection onto a line

Let

$$\mathbf{u} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} 2 \\ 5 \end{bmatrix}, \quad L = \text{span}\{\mathbf{u}\}.$$

1. Compute $\text{proj}_L(\mathbf{x})$.
2. Compute the residual $\mathbf{r} = \mathbf{x} - \text{proj}_L(\mathbf{x})$.
3. Check that $\mathbf{r} \cdot \mathbf{u} = 0$.

Activity: Projection onto a line in code

The following code computes the same projection and residual as the previous activity.

```
u = np.array([3.0, 4.0])
x = np.array([2.0, 5.0])

y = (x @ u) / (u @ u) * u
r = x - y
```

$y, r, r @ u$

1. Which variable stores the projection?
2. Which variable stores the residual?
3. Why should $r @ u$ be close to zero?
4. What formula is represented by $(x @ u) / (u @ u) * u$?

The Unit 1 attention activities, including Many-token attention output shape check, used dot products to make scores and then used weights to mix value vectors. In the language of this unit, the output lies in the span of the value vectors. Thus attention uses the same two operations we have been studying: dot products and linear combinations.

Note: Attention is not orthogonal projection Attention and projection both use dot products, but they solve different problems. In a simple attention step, dot products produce scores, the scores are normalized into weights, and the output is a weighted average of value vectors.

In orthogonal projection, the goal is different: find the closest vector in a subspace. The residual is perpendicular to the subspace.

So both ideas use dot products, but they answer different questions.

Attention asks which stored vectors should be mixed. Projection asks which vector in a subspace is closest. This distinction matters: not every dot-product method is a projection method.

Activity: Weighted average or projection?

For each description, decide whether it is closer to attention or orthogonal projection:

1. The output is a weighted average of stored value vectors.
2. The output is the closest point in a subspace.
3. The residual is orthogonal to every column of A .
4. The weights are obtained by normalizing relevance scores.

Gram-Schmidt orthogonalization

Repeatedly applying Expanding an orthogonal set gives the following result:

Theorem: The Gram-Schmidt orthogonalization algorithm Let U be a subspace of $\mathbb{R}^n$. Then U has an orthogonal basis. Given a basis $\{\mathbf{x}_1, \dots, \mathbf{x}_m\}$ of U , we can iteratively define an orthogonal basis $\{\mathbf{f}_1, \dots, \mathbf{f}_m\}$ of U by defining

$$\begin{aligned}\mathbf{f}_1 &= \mathbf{x}_1 \\ \mathbf{f}_2 &= \mathbf{x}_2 - \frac{\mathbf{x}_2 \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1 \\ \mathbf{f}_3 &= \mathbf{x}_3 - \frac{\mathbf{x}_3 \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1 - \frac{\mathbf{x}_3 \cdot \mathbf{f}_2}{\|\mathbf{f}_2\|^2} \mathbf{f}_2\end{aligned}$$

and so on, i.e. for $2 \leq k \leq m$ defining

$$\mathbf{f}_k = \mathbf{x}_k - \frac{\mathbf{x}_k \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \mathbf{f}_1 - \frac{\mathbf{x}_k \cdot \mathbf{f}_2}{\|\mathbf{f}_2\|^2} \mathbf{f}_2 - \dots - \frac{\mathbf{x}_k \cdot \mathbf{f}_{k-1}}{\|\mathbf{f}_{k-1}\|^2} \mathbf{f}_{k-1}.$$

Then

- $\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_m\}$ is an orthogonal basis of U .
- $\text{span}\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_k\} = \text{span}\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k\}$ for each $k = 1, 2, \dots, m$.

Activity

Find an orthogonal basis of $\text{row}(A)$ (recall Column and Row Space), where

$$A = \begin{bmatrix} 1 & 1 & -1 & -1 \\ 3 & 2 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{bmatrix}.$$

Closest-point activity

Activity

Find the point in the plane defined by the equation $2x + y - z = 0$, which is closest to the point $(1, 2, -1)$.

