

# MATH 345 Skeleton Notes

## Unit 4: Orthogonality and least squares

### Section 4.1: Orthogonality and orthogonal bases

---

© 2024–2026 Jacob Denson, Kaitlyn Phillipson, Sebastien Roch. Except where otherwise noted, this work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

#### Basic definitions

**Definition** Recall the definition of the dot product of two vectors, introduced in Dot product, and the length of a vector as introduced

in Length of a vector. The  between two points  $\mathbf{x} = (x_1, \dots, x_n)$  and  $\mathbf{y} = (y_1, \dots, y_n)$  is the length of the vector  $\mathbf{x} - \mathbf{y}$ , i.e.

$$d(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\| = \sqrt{(x_1 - y_1)^2 + \dots + (x_n - y_n)^2}.$$

If  $\mathbf{x}$  is any non-zero vector, then the vector  $\mathbf{u} = \mathbf{x}/\|\mathbf{x}\|$  is the

(a unit vector is a vec-

tor of length one). Two vectors  $\mathbf{x}$  and  $\mathbf{y}$  are  or

if  $\mathbf{x} \cdot \mathbf{y} = 0$ .

**Theorem: The Cauchy-Schwarz inequality** If  $\mathbf{x}, \mathbf{y}$  are vectors in  $\mathbb{R}^n$ ,

$$|\mathbf{x} \cdot \mathbf{y}| \leq \|\mathbf{x}\| \|\mathbf{y}\|.$$

Moreover,  $|\mathbf{x} \cdot \mathbf{y}| = \|\mathbf{x}\| \|\mathbf{y}\|$  only when  $\mathbf{x}$  and  $\mathbf{y}$  are scalar multiples of each other.

### Activity: Cosine similarity revisited

Let  $\mathbf{u}$  and  $\mathbf{v}$  be nonzero vectors in  $\mathbb{R}^n$ . The cosine similarity from Unit 1 was

$$\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|}.$$

1. Use the Cauchy-Schwarz inequality to explain why this number is between  $-1$  and  $1$ .
2. What condition makes the cosine similarity equal to  $0$ ?
3. In Unit 1, this number measured cosine similarity. What does a value of  $0$  say geometrically?

**Corollary: The triangle inequality** If  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ , then

$$\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|.$$

**Remark** The triangle inequality for vectors implies the triangle inequality for the distance function, i.e. that

$$d(\mathbf{x}, \mathbf{z}) \leq d(\mathbf{x}, \mathbf{y}) + d(\mathbf{y}, \mathbf{z}).$$

Indeed, we calculate that

$$\begin{aligned}
 d(\mathbf{x}, \mathbf{z}) &= \|\mathbf{x} - \mathbf{z}\| \\
 &= \|\mathbf{x} - \mathbf{y} + \mathbf{y} - \mathbf{z}\| \\
 &\leq \|\mathbf{x} - \mathbf{y}\| + \|\mathbf{y} - \mathbf{z}\|.
 \end{aligned}$$

**Theorem** Let  $\mathbf{x}$ ,  $\mathbf{y}$ , and  $\mathbf{z}$  be vectors in  $\mathbb{R}^n$ , and let  $a$  be a scalar in  $\mathbb{R}$ . Then the following properties hold:

- $\mathbf{x} \cdot \mathbf{y} = \mathbf{y} \cdot \mathbf{x}$ .
- $\mathbf{x} \cdot (\mathbf{y} + \mathbf{z}) = \mathbf{x} \cdot \mathbf{y} + \mathbf{x} \cdot \mathbf{z}$ .
- $(a\mathbf{x}) \cdot \mathbf{y} = a(\mathbf{x} \cdot \mathbf{y}) = \mathbf{x} \cdot (a\mathbf{y})$ .
- $\|\mathbf{x}\|^2 = \mathbf{x} \cdot \mathbf{x}$ .
- $\mathbf{x} \cdot \mathbf{x} \geq 0$ , and  $\mathbf{x} \cdot \mathbf{x} = 0$  if and only if  $\mathbf{x} = \mathbf{0}$ .
- $\|a\mathbf{x}\| = |a|\|\mathbf{x}\|$ .

### Activity

Show that  $\|\mathbf{x} + \mathbf{y}\|^2 = \|\mathbf{x}\|^2 + 2(\mathbf{x} \cdot \mathbf{y}) + \|\mathbf{y}\|^2$  for any  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ .

### Activity

Suppose that  $\mathbb{R}^n = \text{span}\{\mathbf{f}_1, \dots, \mathbf{f}_k\}$  for some vectors  $\mathbf{f}_i$ . If  $\mathbf{x} \in \mathbb{R}^n$ , and  $\mathbf{x} \cdot \mathbf{f}_i = 0$  for each  $i$ , then show  $\mathbf{x} = \mathbf{0}$ .

## Orthogonal sets and the expansion theorem

**Definition: Orthogonal and orthonormal sets** A set of vectors  $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$  is called an

if the set does not contain the zero vector,

and  $\mathbf{x}_i \cdot \mathbf{x}_j = 0$  for  $i \neq j$ . A set  $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$  is called if it is orthogonal and if in addition,  $\|\mathbf{x}_i\| = 1$  for each  $i$ .

Note that  $\{\mathbf{x}\}$  is an orthogonal set if  $\mathbf{x} \neq \mathbf{0}$ . The standard basis  $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$  (see Standard basis) is an important orthonormal set in  $\mathbb{R}^n$ .

## Orthogonal sets and bases

### Activity

Consider the following vectors in  $\mathbb{R}^4$ , written as column vectors:

$$\mathbf{f}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ -1 \end{bmatrix} \quad \mathbf{f}_2 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 2 \end{bmatrix} \quad \mathbf{f}_3 = \begin{bmatrix} -1 \\ 0 \\ 1 \\ 0 \end{bmatrix} \quad \mathbf{f}_4 = \begin{bmatrix} -1 \\ 3 \\ -1 \\ 1 \end{bmatrix}.$$

Verify that these vectors form an orthogonal set in  $\mathbb{R}^4$ . Then find the orthonormal set obtained by ‘normalizing’ these vectors, i.e. multiplying each vector by an appropriate constant to obtain a unit vector.

**Theorem: Pythagoras' theorem** If  $\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_k\}$  is an orthogonal set in  $\mathbb{R}^n$ , then

$$\|\mathbf{f}_1 + \mathbf{f}_2 + \dots + \mathbf{f}_k\|^2 = \|\mathbf{f}_1\|^2 + \|\mathbf{f}_2\|^2 + \dots + \|\mathbf{f}_k\|^2.$$

**Theorem** Every orthogonal set in  $\mathbb{R}^n$  is linearly independent.

**Proof** Let  $\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_k\}$  be an orthogonal set in  $\mathbb{R}^n$ . To show the set is linearly independent (recalling Linear Independence), we must show the only solution to the linear equation

$$t_1\mathbf{f}_1 + t_2\mathbf{f}_2 + \dots + t_k\mathbf{f}_k = \mathbf{0}$$

occurs when  $t_1 = \dots = t_k = 0$ .

If we take the dot product of both sides of the equation with  $\mathbf{f}_1$ , we find that

$$\mathbf{f}_1 \cdot (t_1\mathbf{f}_1 + t_2\mathbf{f}_2 + \dots + t_k\mathbf{f}_k) = \mathbf{f}_1 \cdot \mathbf{0}$$

and therefore that

$$t_1(\mathbf{f}_1 \cdot \mathbf{f}_1) + t_2(\mathbf{f}_1 \cdot \mathbf{f}_2) + \dots + t_k(\mathbf{f}_1 \cdot \mathbf{f}_k) = 0$$

Since the set of vectors is orthogonal,  $\mathbf{f}_1 \cdot \mathbf{f}_j = 0$  for all  $j \neq 1$ , and  $\mathbf{f}_1 \cdot \mathbf{f}_1 = \|\mathbf{f}_1\|^2$ , so we conclude that

$$t_1\|\mathbf{f}_1\|^2 = 0.$$

Since  $\mathbf{f}_1 \neq \mathbf{0}$  (by definition, an orthogonal set cannot contain the zero vector),  $\|\mathbf{f}_1\|^2 > 0$ , and so it must be true that  $t_1 = 0$ . But now repeating this argument with 1 replaced by any index  $i$ , we conclude that  $t_i = 0$ . Therefore the set  $\{\mathbf{f}_1, \dots, \mathbf{f}_k\}$  is linearly independent.

**Theorem: Expansion theorem** Let  $\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_m\}$  be an orthogonal basis of a subspace  $U$  of  $\mathbb{R}^n$ . If  $\mathbf{v}$  is any vector in  $U$ , we have

$$\mathbf{v} = \left( \frac{\mathbf{v} \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2} \right) \mathbf{f}_1 + \left( \frac{\mathbf{v} \cdot \mathbf{f}_2}{\|\mathbf{f}_2\|^2} \right) \mathbf{f}_2 + \dots + \left( \frac{\mathbf{v} \cdot \mathbf{f}_m}{\|\mathbf{f}_m\|^2} \right) \mathbf{f}_m$$

**Proof** Since  $\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_m\}$  spans  $U$ , any  $\mathbf{v} \in U$  can be written uniquely as

$$\mathbf{v} = t_1 \mathbf{f}_1 + t_2 \mathbf{f}_2 + \dots + t_m \mathbf{f}_m$$

for some  $t_1, \dots, t_m$ . To find coefficient  $t_1$ , take the dot product of both sides with  $\mathbf{f}_1$ , obtaining that

$$\begin{aligned} \mathbf{v} \cdot \mathbf{f}_1 &= (t_1 \mathbf{f}_1 + t_2 \mathbf{f}_2 + \dots + t_m \mathbf{f}_m) \cdot \mathbf{f}_1 \\ &= t_1(\mathbf{f}_1 \cdot \mathbf{f}_1) + t_2(\mathbf{f}_2 \cdot \mathbf{f}_1) + \dots + t_m(\mathbf{f}_m \cdot \mathbf{f}_1). \end{aligned}$$

Since the basis is orthogonal,  $\mathbf{f}_i \cdot \mathbf{f}_j = 0$  for  $i \neq j$ , and so the equation simplifies to

$$\mathbf{v} \cdot \mathbf{f}_1 = t_1 \|\mathbf{f}_1\|^2.$$

Rearranging, we conclude that

$$t_1 = \frac{\mathbf{v} \cdot \mathbf{f}_1}{\|\mathbf{f}_1\|^2}$$

A similar argument shows that

$$t_i = \frac{\mathbf{v} \cdot \mathbf{f}_i}{\|\mathbf{f}_i\|^2}$$

for each  $i$ , concluding the proof.

**Definition: The Fourier expansion in an orthogonal basis** Given an orthogonal basis  $\{\mathbf{f}_1, \dots, \mathbf{f}_m\}$  for some subspace  $V$  of  $\mathbb{R}^n$ , the expansion of  $\mathbf{v} \in V$  as a linear combination of this basis is called the

of the vector  $\mathbf{v}$ , and the coefficients

$$t_i = \frac{\mathbf{v} \cdot \mathbf{f}_i}{\|\mathbf{f}_i\|^2}$$

are called the

of  $\mathbf{v}$ .

### Activity

Let  $\mathbf{v} = (a, b, c, d)$  be a vector in  $\mathbb{R}^4$ . Find the Fourier expansion of  $\mathbf{v}$  as a linear combination of the orthogonal basis  $\{\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3, \mathbf{f}_4\}$  given in Activity act-6-1-orthogonal-sets.

Dot products and orthogonality now become tools for finding closest vectors.