

MATH 345 Skeleton Notes

Unit 7: Abstract vector spaces and polynomial approximation

Section 7.2: Inner product spaces

© 2024–2026 Jacob Denson, Kaitlyn Phillipson, Sebastien Roch. Except where otherwise noted, this work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

Unit 4 used the dot product on $\mathbb{R}^n$ to define length, distance, orthogonality, projection, and least squares. An inner product space keeps exactly the rules needed for those ideas. This section introduces the abstract definition and keeps three examples in view: $\mathbb{R}^n$, polynomial spaces, and matrix spaces.

First, we define inner products.

Definition: Inner Products Let V be a vector space. An on V is a function $\langle \cdot, \cdot \rangle$ that assigns to each of vectors $\mathbf{v}, \mathbf{w}$ in V a $\langle \mathbf{v}, \mathbf{w} \rangle$ satisfying the following properties:

1. $\langle \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{w}, \mathbf{v} \rangle$ for all $\mathbf{v}, \mathbf{w}$ in V .
2. $\langle \mathbf{v} + \mathbf{w}, \mathbf{u} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle + \langle \mathbf{w}, \mathbf{u} \rangle$ for all $\mathbf{u}, \mathbf{v}, \mathbf{w}$ in V .
3. $\langle r\mathbf{v}, \mathbf{w} \rangle = r\langle \mathbf{v}, \mathbf{w} \rangle$ for all $\mathbf{v}$ and $\mathbf{w}$ in V and r in $\mathbb{R}$.
4. $\langle \mathbf{v}, \mathbf{v} \rangle > 0$ for all $\mathbf{v} \neq \mathbf{0}$ in V .

The vector space V together with $\langle \cdot, \cdot \rangle$ is called an .

Theorem For an inner product $\langle \cdot, \cdot \rangle$ on a vector space V , the following additional properties automatically hold:

1. $\langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle$ for any $\mathbf{u}, \mathbf{v}$ in V .
2. $\langle \mathbf{u}, r\mathbf{v} \rangle = r\langle \mathbf{u}, \mathbf{v} \rangle$ for any $\mathbf{u}, \mathbf{v}$ in V , and r in $\mathbb{R}$.
3. $\langle \mathbf{v}, \mathbf{0} \rangle = 0 = \langle \mathbf{0}, \mathbf{v} \rangle$
4. $\langle \mathbf{v}, \mathbf{v} \rangle = 0$ if and only if $\mathbf{v} = \mathbf{0}$.

The dot product on $\mathbb{R}^n$ is an inner product. We will often call the dot product the

Activity

Let

$$\mathbf{u} = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} \quad \text{and} \quad \mathbf{v} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$$

be vectors in $\mathbb{R}^2$, and define

$$\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_1 - u_2 v_1 - u_1 v_2 + 3u_2 v_2$$

Show this definition satisfies the 4th property of Theorem thm-7-4-inner-product-properties, i.e. that $\langle \mathbf{u}, \mathbf{u} \rangle > 0$ for $\mathbf{u} \neq \mathbf{0}$.

Theorem Let $E = \{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ be the standard basis for $\mathbb{R}^n$, and let $\langle \cdot, \cdot \rangle$ be any inner product on $\mathbb{R}^n$. Let $c_{ij} = \langle \mathbf{e}_i, \mathbf{e}_j \rangle$ and consider the matrix

$C = [c_{ij}]$. Then

1. C is a

2. $\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u}^T C \mathbf{v}$ for all $\mathbf{u}, \mathbf{v}$ in $\mathbb{R}^n$.

Definition Given an inner product $\langle \cdot, \cdot \rangle$ on $\mathbb{R}^n$, the symmetric matrix C analyzed in Theorem thm-7-4-inner-product-in-coordinates is called the

associated with the inner product.

Remark If a symmetric $n \times n$ matrix C satisfies $\mathbf{x}^T C \mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$, then C defines an inner product on $\mathbb{R}^n$. Moreover, these are the *only* symmetric matrices that define inner products. They are called *positive definite matrices*.

Definition: Positive Definite Matrices An $n \times n$ matrix A is

if it is symmetric, and if $\mathbf{x}^T A \mathbf{x} > 0$ for all $\mathbf{x} \neq \mathbf{0}$.

Activity

Consider the inner product on $\mathbb{R}^2$ defined by

$$\langle \mathbf{u}, \mathbf{v} \rangle = u_1 v_1 - u_2 v_1 - u_1 v_2 + 3u_2 v_2.$$

Find C such that $\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u}^T C \mathbf{v}$.

Definition: The Norm of a Vector In an inner product space, the

of a vector $\mathbf{v}$ is

$$\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}.$$

The between two vectors $\mathbf{u}$ and $\mathbf{v}$ in an inner product space is

$$d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\|.$$

Definition: Unit Vectors A vector $\mathbf{v}$ in an inner product space V is

called a if $\|\mathbf{v}\| = 1$. This is equivalent to the equa-

tion $\langle \mathbf{v}, \mathbf{v} \rangle = 1$. The set of all unit vectors in V is called the in V .

Theorem If $\mathbf{v}$ is a nonzero vector of an inner product space, then

$$\mathbf{u} = \frac{1}{\|\mathbf{v}\|} \mathbf{v}$$

is the unique unit vector that is a positive multiple of $\mathbf{v}$. We often say that $\mathbf{u}$ is the unit vector ‘in the same direction as $\mathbf{v}$ ’.

Definition: Orthogonal Vectors Two vectors $\mathbf{u}$ and $\mathbf{v}$ in an inner

product space are if

$$\langle \mathbf{u}, \mathbf{v} \rangle = 0.$$

The next examples show that inner products are not limited to ordinary coordinate vectors. The objects can be polynomials or matrices, as long as the rule for $\langle \cdot, \cdot \rangle$ satisfies the inner-product properties.

Spaces of polynomials

Activity

Let $V = \mathbf{P}_1$ be the vector space of all polynomials of degree at most 1. Define

$$\langle f, g \rangle = \int_0^1 f(x)g(x) dx.$$

Does this definition define an inner product on V ?

Activity

Consider $\mathbf{P}_1$ with the inner product

$$\langle p, q \rangle = \int_0^1 p(x)q(x) dx.$$

Task Calculate the norm of the polynomial $p(x) = x$.

Task Calculate the distance between $q_1(x) = 2x + 1$ and $q_2(x) = 3x$.

Task Find the unit vector in the same direction as $p(x) = x$.

Task Are x and $x - 2/3$ orthogonal polynomials in $\mathbf{P}_1$?

Spaces of matrices

Definition: The Trace of a Matrix The trace of a square $n \times n$ matrix A , denoted by $\text{Tr}(A)$, is the sum of its diagonal entries. For $A = [a_{ij}]$, we have

$$\text{Tr}(A) = a_{11} + a_{22} + \cdots + a_{nn}.$$

Theorem For any $n \times n$ matrices A and B ,

$$\text{Tr}(A^T) = \text{Tr}(A) \quad \text{and} \quad \text{Tr}(A + B) = \text{Tr}(A) + \text{Tr}(B).$$

Activity

Consider $\mathbf{M}_{22}$, and let A, B be matrices in $\mathbf{M}_{22}$. Define

$$\langle A, B \rangle = \text{Tr}(A^T B).$$

Task Find $\langle A, B \rangle$ for $A = \begin{bmatrix} 1 & 2 \\ 2 & -3 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 2 \\ -1 & 1 \end{bmatrix}$.

Task Show that the map $\langle, \rangle$ defined above is an inner product.

Task Find the distance between the two matrices

$$A = \begin{bmatrix} 3 & 1 \\ 0 & -1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} -1 & 0 \\ 1 & 2 \end{bmatrix}.$$

Task Find a value for x so that the two matrices

$$A = \begin{bmatrix} 1 & 0 \\ -2 & 2 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} -3 & 2 \\ x & 0 \end{bmatrix}$$

are orthogonal.