


Is timing everything? The effects of measurement timing on the performance of nonlinear longitudinal models

—
Sebastian Sciarra


Model of workplace deviance

Model of workplace deviance




Christian & Ellis (2011)

Model of workplace deviance




Christian & Ellis (2011)

Model of workplace deviance



Christian & Ellis (2011)


Model of workplace deviance



Christian & Ellis (2011)

Time is integral to understanding psychological processes


Model of workplace deviance



Christian & Ellis (2011)


Time is integral to understanding psychological processes

Model of thriving at work



Time is integral to understanding psychological processes


Model of organizational trust



Mayer et al. (1995)

Time is integral to understanding psychological processes

Model of turnover intentions




Chen et al. (2011)

Change over time is likely to be nonlinear



Change over time is likely to be nonlinear


Forgetting curve



Li (2017)

Change over time is likely to be nonlinear


Number of speech errors throughout child development



Burchinal & Appelbaum (1991)

Change over time is likely to be nonlinear

Development of habits




Fournier et al. (2017)

Experiment 1

Question 1: Does placing measurements near periods of change increase model performance?


Question 2: How to space measurements when the nature of change is unknown?

The Monte Carlo method




1. Population definition

μ (known)




The Monte Carlo method




1. Population definition

μ (known)



2. Sample generation




The Monte Carlo method


 = model
 = sample

1. Population definition


μ (known)



2. Sample generation



3. Modelling




The Monte Carlo method


 = model
 = sample

1. Population definition


μ (known)



2. Sample generation



4. Model performance




3. Modelling

The Monte Carlo method


 = model
 = sample

1. Population definition


μ (known)




2. Sample generation



μ (estimated)




4. Model performance




3. Modelling

The Monte Carlo method: Population definition




Lawrence et al. (2001)


The Monte Carlo method: Population definition



The Monte Carlo method: Population definition




The Monte Carlo method: Population definition



1. Baseline (θ) = 3.00
2. Maximal elevation (α) = 3.32
3. Days to halfway elevation (β) = 180.00

The Monte Carlo method: Population definition




1. Baseline (θ) = 3.00
2. Maximal elevation (α) = 3.32
3. Days to halfway elevation (β) = 180.00
4. Triquarter-halfway delta (γ) = 20.00

The Monte Carlo method


 = model
 = sample

1. Population definition

μ (known)



2. Sample generation




μ (estimated)

Parameter bias

\bar{x}

4. Model performance




3. Modelling



The Monte Carlo method

 = model
 = sample


1. Population definition



2. Sample generation




4. Model performance




3. Modelling


The Monte Carlo method: Population definition




The Monte Carlo method: Population definition




The Monte Carlo method: Population definition




The Monte Carlo method: Population definition




The Monte Carlo method: Population definition



The Monte Carlo method: Population definition



The Monte Carlo method: Population definition




*Random effects** (grey curves)

1. Baseline (θ) = 0.05
2. Maximal elevation (α) = 0.05
3. Days to halfway elevation (β) = 10.00
4. Triquarter-halfway delta (γ) = 4.00

*random effects are given in standard deviation units

The Monte Carlo method: Population definition



Fixed effects (black curve)

1. Baseline (θ) = 3.00
2. Maximal elevation (α) = 3.32
3. Days to halfway elevation (β) = 180.00
4. Triquarter-halfway delta (γ) = 20.00

*Random effects** (grey curves)


1. Baseline (θ) = 0.05
2. Maximal elevation (α) = 0.05
3. Days to halfway elevation (β) = 10.00
4. Triquarter-halfway delta (γ) = 4.00

*random effects are given in standard deviation units



The Monte Carlo method

 = model
 = sample


1. Population definition



2. Sample generation



4. Model performance




3. Modelling

The Monte Carlo method

 = model
 = sample



1. Population definition

$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (known)



2. Sample generation

$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (estimated)




4. Model performance

3. Modelling

Independent variable I: Nature of change

Question 2: How to space measurements when the nature of change is unknown?


Independent variable I: Nature of change



Question 2: How to space measurements when the nature of change is unknown?

Independent variable I: Nature of change


Logistic curve with $\beta_{fixed} = 180$



Question 2: How to space measurements when the nature of change is unknown?

Independent variable I: Nature of change

Logistic curve with $\beta_{\text{fixed}} = 280$



Question 2: How to space measurements when the nature of change is unknown?


The Monte Carlo method

 = model
 = sample

1. Population definition

IV 1: Nature of change
($\beta_{fixed} = 80, 180, \text{ or } 280$)


2. Sample generation



Independent variable II: Number of measurements


Independent variable II: Number of measurements

Five measurements




Independent variable II: Number of measurements

Seven measurements




Independent variable II: Number of measurements

Nine measurements




Independent variable II: Number of measurements

Eleven measurements




The Monte Carlo method




1. Population definition

IV 1: Nature of change
($\beta_{fixed} = 80, 180, \text{ or } 280$)

$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (known)




$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (estimated)




2. Sample generation

IV 2: Number of measurements (5, 7, 9, or 11)



4. Model performance




3. Modelling

Independent variable III: Measurement spacing

Question 2: How to space measurements when the nature of change is unknown?

Independent variable III: Measurement spacing


Equal spacing



Question 2: How to space measurements when the nature of change is unknown?

Independent variable III: Measurement spacing


Time-interval increasing spacing



Question 2: How to space measurements when the nature of change is unknown?

Independent variable III: Measurement spacing


Time-interval decreasing spacing



Question 2: How to space measurements when the nature of change is unknown?


Independent variable III: Measurement spacing

Middle-and-extreme spacing




Question 2: How to space measurements when the nature of change is unknown?

The Monte Carlo method




1. Population definition

IV 1: Nature of change
($\beta_{fixed} = 80, 180, \text{ or } 280$)




$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (estimated)



2. Sample generation

IV 2: Number of measurements (5, 7, 9, or 11)

IV 3: Spacing of measurements
(equal, time-in., time-dec.,
middle-and-extreme)



4. Model performance

3. Modelling

Monte Carlo method: Modelling



Monte Carlo method: Modelling

Structured latent growth curve model


Monte Carlo method: Modelling

Structured latent growth curve model = latent growth curve model

+

Taylor series
approximations


The Monte Carlo method




1. Population definition

IV 1: Nature of change
($\beta_{\text{fixed}} = 80, 180, 280$)

$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (known)




$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (estimated)




2. Sample generation

IV 2: Number of measurements (5, 7, 9, 11)

IV 3: Spacing of measurements (equal, time-inc., time dec., mid-extreme)




4. Model performance



3. Modelling

Structured latent growth model


The Monte Carlo method




1. Population definition

IV 1: Nature of change
($\beta_{\text{fixed}} = 80, 180, 280$)

$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (known)




$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (estimated)



2. Sample generation


IV 2: Number of measurements (5, 7, 9, 11)

IV 3: Spacing of measurements (equal, time-inc., time dec., mid-extreme)



$n = k = 1000$

4. Model performance



3. Modelling

Structured latent growth model

Experiment 1

Question 1: Does placing measurements near periods of change increase model performance?

Question 2: How to space measurements when the nature of change is unknown?

Experiment 1

Question 1: Does placing measurements near periods of change increase model performance?

Question 2: How to space measurements when the nature of change is unknown?

Experiment 1

Question 1: Does placing measurements near periods of change increase model performance?

IV 1: Nature of change ($\beta_{\text{fixed}} = 80, 180, 280$)

IV 2: Number of measurements (5, 7, 9, 11)

Question 2: How to space measurements when the nature of change is unknown?

IV 3: Spacing of measurements (equal, time-inc., time dec., mid-extreme)

Experiment 1

Question 1: Does placing measurements near periods of change increase model performance?

IV 1: Nature of change ($\beta_{\text{fixed}} = 80, 180, 280$)

3

x

4

= 48 cells

Question 2: How to space measurements when the nature of change is unknown?

IV 2: Number of measurements (5, 7, 9, 11)

IV 3: Spacing of measurements (equal, time-inc., time dec., mid-extreme)

x

4

Experiment 1

Question 1: Does placing measurements near periods of change increase model performance?

IV 1: Nature of change ($\beta_{\text{fixed}} = 80, 180, 280$)

3

x

= 48 cells

1 cell

Question 2: How to space measurements when the nature of change is unknown?

IV 2: Number of measurements (5, 7, 9, 11)

4

x

4

IV 3: Spacing of measurements (equal, time-inc., time dec., mid-extreme)

Experiment 1

Question 1: Does placing measurements near periods of change increase model performance?

IV 1: Nature of change ($\beta_{fixed} = 80, 180, 280$)

3

X

4

= 48 cells

1 cell —→

Nature of change ($\beta_{fixed} = 180$)

Number of measurements (NM) = 5

Spacing = time-interval increasing

Question 2: How to space measurements when the nature of change is unknown?

IV 3: Spacing of measurements (equal, time-inc., time dec., mid-extreme)

X

4

Experiment 1

Question 1: Does placing measurements near periods of change increase model performance?

IV 1: Nature of change ($\beta_{fixed} = 80, 180, 280$)

3

X

4

= 48 cells

1 cell

Nature of change ($\beta_{fixed} = 180$)

Number of measurements (NM) = 5

Spacing = time-interval increasing

Question 2: How to space measurements when the nature of change is unknown?

IV 3: Spacing of measurements (equal, time-inc., time dec., mid-extreme)

X

4

Fixed-Effect Triquarter-Halfway Delta Parameter (γ_{fixed})

Experiment 1

Question 1: Does placing measurements near periods of change increase model performance?

IV 1: Nature of change ($\beta_{fixed} = 80, 180, 280$)

3 X 4
= 48 cells

1 cell

IV 2: Number of measurements (5, 7, 9, 11)

Question 2: How to space measurements when the nature of change is unknown?


IV 3: Spacing of measurements (equal, time-inc., time dec., mid-extreme)

X 4

Nature of change ($\beta_{fixed} = 180$)
Number of measurements (NM) = 5
Spacing = time-interval increasing


$$n = k = 1000$$

Fixed-Effect Triquarter-Halfway Delta Parameter (γ_{fixed})





Monte Carlo method: Model performance

*Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$,
NM = 5, Spacing = Time-Interval Increasing)*





Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{fixed} = 180.00$, $NM = 5$, Spacing = Time-Interval Increasing)





Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)




Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)



Monte Carlo method: Model performance


Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)



Bias: difference between average estimated value and population value

Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{fixed} = 180.00$, $NM = 5$, Spacing = Time-Interval Increasing)




Bias: difference between average estimated value and population value

Estimates were biased if bias > 10% of population value

Muthén et al. (1997)

Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{fixed} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)




Bias: difference between average estimated value and population value

Estimates were biased if bias > 10% of population value

Muthén et al. (1997)

Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{fixed} = 180.00$, $NM = 5$, Spacing = Time-Interval Increasing)




Bias: difference between average estimated value and population value

Estimates were biased if bias > 10% of population value

Muthén et al. (1997)

Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{fixed} = 180.00$, $NM = 5$, Spacing = Time-Interval Increasing)



Bias: difference between average estimated value and population value


Estimates were biased if bias > 10% of population value

Precision: range of values covered by the middle 95% of estimates

Muthén et al. (1997)

Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{fixed} = 180.00$, $NM = 5$, Spacing = Time-Interval Increasing)



Bias: difference between average estimated value and population value


Estimates were biased if bias > 10% of population value

Precision: range of values covered by the middle 95% of estimates

Muthén et al. (1997)

Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)



Bias: difference between average estimated value and population value


Estimates were biased if bias > 10% of population value

Precision: range of values covered by the middle 95% of estimates

Muthén et al. (1997)

Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, $NM = 5$, Spacing = Time-Interval Increasing)



Bias: difference between average estimated value and population value

Estimates were biased if bias > 10% of population value


Precision: range of values covered by the middle 95% of estimates

Estimation was imprecise if a whisker length > 10% of population value

Muthén et al. (1997)

Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Delta Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)




Bias/Precision Plot for Fixed-Effect Triquarter-Halfway Delta Parameter



Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Delta Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)




Bias/Precision Plot for Fixed-Effect Triquarter-Halfway Delta Parameter




Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Delta Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)




Bias/Precision Plot for Fixed-Effect Triquarter-Halfway Delta Parameter




Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Delta Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)




Bias/Precision Plot for Fixed-Effect Triquarter-Halfway Delta Parameter




Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Delta Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)




Bias/Precision Plot for Fixed-Effect Triquarter-Halfway Delta Parameter




Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Delta Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)





Bias/Precision Plot for Fixed-Effect Triquarter-Halfway Delta Parameter




Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Delta Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)





Bias/Precision Plot for Fixed-Effect Triquarter-Halfway Delta Parameter




Number of Measurements	■ 5	● 7	▲ 9	◆ 11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		

Monte Carlo method: Model performance

Values Estimated for Fixed-Effect Triquarter-Halfway Delta Parameter (γ_{fixed} ; $\beta_{\text{fixed}} = 180.00$, NM = 5, Spacing = Time-Interval Increasing)



Bias/Precision Plot for Fixed-Effect Triquarter-Halfway Delta Parameter for Time-Interval Increasing Spacing




Number of Measurements	\blacksquare 5 \bullet 7 \blacktriangle 9 \blacklozenge 11
Is Unbiased?	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No
Is Precise?	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No

Parameter Values Estimated for Day- and Likert-Unit Parameters in Experiment 1

Measurement	Number of Measurements	β_{fixed} (Days to halfway elevation)			β_{random} (Days to halfway elevation)			γ_{fixed} (Triquarter-halfway delta)			γ_{random} (Triquarter-halfway delta)		
					Pop value = 10.00			Pop value = 20.00			Pop value = 4.00		
		80	180	280	80	180	280	80	180	280	80	180	280
Measurement	Number of Measurements	80	180	280	80	180	280	80	180	280	80	180	280
Equal spacing	5	79.73	179.78	279.81 [□]	10.14	10.40	10.08	19.37	19.49	19.71	7.41 [□]	14.53 [□]	8.11 [□]
	7	80.21	178.99	279.55 [□]	10.16	10.55	10.13	20.67	20.83	20.60	4.37	5.14 [□]	4.41 [□]
	9	80.00	179.94	279.99 [□]	10.29	10.37	10.34	20.77	20.76	20.67	4.24	4.14	4.30
	11	80.03	180.01	279.88 [□]	10.27	10.29	10.32	20.64	20.70	20.64	4.13	4.08	4.18
Time-interval increasing	5	79.88	180.10	274.37 [□]	10.32	9.73	13.04 [□]	20.71	20.39	18.32	4.57 [□]	4.99 [□]	6.20 [□]
	7	80.19	179.82	279.86 [□]	10.42	10.47	10.14	20.66	20.79	19.78	4.29	4.87 [□]	7.03 [□]
	9	79.59	179.06	279.70 [□]	10.07	10.22	10.20	20.33	20.66	20.72	4.17	4.25	4.32
	11	79.89	179.84	279.62 [□]	10.38	10.30	10.47	20.78	20.75	20.68	4.23	4.18	4.13
Time-interval decreasing	5	70.67	179.92	279.63 [□]	15.28 [□]	9.80	10.22	16.63	20.07	20.55	5.48 [□]	5.17 [□]	4.59 [□]
	7	78.23	178.22	279.84 [□]	10.08	10.46	10.39	19.38	20.59	20.69	6.80 [□]	5.09 [□]	4.24
	9	79.95	179.34	278.98 [□]	10.03	10.20	10.05	20.42	20.54	20.28	4.37	4.32	4.19
	11	79.42	179.70	279.52 [□]	10.38	10.13	10.06	20.75	20.45	20.31	4.17	4.16	4.17
Middle-and-extreme spacing	5	71.95	179.61	287.73 [□]	16.78 [□]	10.26	16.74 [□]	15.59	20.61	17.09	6.54 [□]	4.24	8.61 [□]
	7	80.45	180.00	279.15 [□]	13.93 [□]	10.25	13.69 [□]	20.71	20.58	20.61	5.21 [□]	4.16	4.98 [□]
	9	80.28	180.05	279.63 [□]	10.42	10.24	10.24	20.91	20.65	20.85	4.74 [□]	4.26	4.72 [□]
	11	80.19	179.96	279.86 [□]	10.27	10.28	10.15	20.71	20.70	20.71	4.14	4.08	4.16

Experiment 1 design (Monte Carlo method)





1. Population definition

IV 1: Nature of change
($\beta_{\text{fixed}} = 80, 180, 280$)

2. Sample generation

IV 2: Number of measurements (5, 7, 9, 11)

IV 3: Spacing of measurements (equal, time-inc., time dec., mid-extreme)




Question 1: Does placing measurements near periods of change increase model performance?

Question 1: Does placing measurements near periods of change increase model performance?

Answer 1: Model performance increases when measurements are placed near periods of change

Answer 1: Model performance increases when measurements are placed near periods of change




Day

Highest Model Performance?

- Yes
- No

Answer 1: Model performance increases when measurements are placed near periods of change

*Time-interval
increasing spacing*




Nature of Change (Population Value Set for β_{fixed})

Number of Measurements	■ 5	● 7	▲ 9	◆ 11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		

Answer 1: Model performance increases when measurements are placed near periods of change

*Time-interval
increasing spacing*




Nature of Change (Population Value Set for β_{fixed})

Number of Measurements	■ 5	● 7	▲ 9	◆ 11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		


Answer 1: Model performance increases when measurements are placed near periods of change

*Time-interval
increasing spacing*




Answer 1: Model performance increases when measurements are placed near periods of change


A : $\beta_{\text{fixed}} = 180$ With Equal Spacing




B : $\beta_{\text{fixed}} = 80$ With Time-Interval Increasing Spacing



C : $\beta_{\text{fixed}} = 280$ With Time-Interval Decreasing Spacing



D : $\beta_{\text{fixed}} = 180$ With Middle-and-Extreme Spacing



Highest Model Performance?

- Yes
- No

Question 1: Does placing measurements near periods of change increase model performance?

Answer 1: Model performance increases when measurements are placed near periods of change

Question 2: How to space measurements when the nature of change is unknown?

Question 1: Does placing measurements near periods of change increase model performance?

Answer 1: Model performance increases when measurements are placed near periods of change

Question 2: How to space measurements when the nature of change is unknown?

Answer 2: When the nature of change is unknown, model performance is highest with equal spacing

Answer 2: When the nature of change is unknown, model performance is highest with equal spacing

Spacing Schedule	Qualitative Description
Equal (see Figure 2.5 and Table 2.3)	
Time-interval increasing (see Figure 2.7 and Table 2.6)	
Time-interval decreasing (see Figure 2.9 and Table 2.9)	
Middle-and-extreme (see Figure 2.11 and Table 2.9)	

Answer 2: When the nature of change is unknown, model performance is highest with equal spacing

Spacing Schedule	Qualitative Description
Equal (see Figure 2.5 and Table 2.3)	Largest improvements in bias and precision with NM = 7
Time-interval increasing (see Figure 2.7 and Table 2.6)	
Time-interval decreasing (see Figure 2.9 and Table 2.9)	
Middle-and-extreme (see Figure 2.11 and Table 2.9)	

Answer 2: When the nature of change is unknown, model performance is highest with equal spacing

Spacing Schedule	Qualitative Description
Equal (see Figure 2.5 and Table 2.3)	Largest improvements in bias and precision with NM = 7
Time-interval increasing (see Figure 2.7 and Table 2.6)	Largest improvements in bias and precision with NM = 9
Time-interval decreasing (see Figure 2.9 and Table 2.9)	Largest improvements in bias and precision with NM = 9
Middle-and-extreme (see Figure 2.11 and Table 2.9)	Largest improvements in bias and precision with NM = 9

Answer 2: When the nature of change is unknown, model performance is highest with equal spacing



Error Bar Summary

Spacing Schedule	Qualitative Description
Equal (see Figure 2.5 and Table 2.3)	Largest improvements in bias and precision with NM = 7
Time-interval increasing (see Figure 2.7 and Table 2.6)	Largest improvements in bias and precision with NM = 9
Time-interval decreasing (see Figure 2.9 and Table 2.9)	Largest improvements in bias and precision with NM = 9
Middle-and-extreme (see Figure 2.11 and Table 2.9)	Largest improvements in bias and precision with NM = 9

Answer 2: When the nature of change is unknown, model performance is highest with equal spacing



		Error Bar Summary
Spacing Schedule	Qualitative Description	β_{fixed}
Equal (see Figure 2.5 and Table 2.3)	Largest improvements in bias and precision with NM = 7	5.64
Time-interval increasing (see Figure 2.7 and Table 2.6)	Largest improvements in bias and precision with NM = 9	4.97
Time-interval decreasing (see Figure 2.9 and Table 2.9)	Largest improvements in bias and precision with NM = 9	4.88
Middle-and-extreme (see Figure 2.11 and Table 2.9)	Largest improvements in bias and precision with NM = 9	6.51

Answer 2: When the nature of change is unknown, model performance is highest with equal spacing



		Error Bar Summary	
Spacing Schedule	Qualitative Description	β_{fixed}	γ_{fixed}
Equal (see Figure 2.5 and Table 2.3)	Largest improvements in bias and precision with NM = 7	5.64	4.37
Time-interval increasing (see Figure 2.7 and Table 2.6)	Largest improvements in bias and precision with NM = 9	4.97	3.45
Time-interval decreasing (see Figure 2.9 and Table 2.9)	Largest improvements in bias and precision with NM = 9	4.88	3.40
Middle-and-extreme (see Figure 2.11 and Table 2.9)	Largest improvements in bias and precision with NM = 9	6.51	5.55

Answer 2: When the nature of change is unknown, model performance is highest with equal spacing



Spacing Schedule	Qualitative Description	Error Bar Summary		
		β_{fixed}	γ_{fixed}	β_{random}
Equal (see Figure 2.5 and Table 2.3)	Largest improvements in bias and precision with NM = 7	5.64	4.37	7.74
Time-interval increasing (see Figure 2.7 and Table 2.6)	Largest improvements in bias and precision with NM = 9	4.97	3.45	6.31
Time-interval decreasing (see Figure 2.9 and Table 2.9)	Largest improvements in bias and precision with NM = 9	4.88	3.40	6.15
Middle-and-extreme (see Figure 2.11 and Table 2.9)	Largest improvements in bias and precision with NM = 9	6.51	5.55	9.02

Answer 2: When the nature of change is unknown, model performance is highest with equal spacing



Spacing Schedule	Qualitative Description	Error Bar Summary			
		β_{fixed}	γ_{fixed}	β_{random}	γ_{random}
Equal (see Figure 2.5 and Table 2.3)	Largest improvements in bias and precision with NM = 7	5.64	4.37	7.74	7.02
Time-interval increasing (see Figure 2.7 and Table 2.6)	Largest improvements in bias and precision with NM = 9	4.97	3.45	6.31	5.97
Time-interval decreasing (see Figure 2.9 and Table 2.9)	Largest improvements in bias and precision with NM = 9	4.88	3.40	6.15	5.96
Middle-and-extreme (see Figure 2.11 and Table 2.9)	Largest improvements in bias and precision with NM = 9	6.51	5.55	9.02	7.20

Question 1: Does placing measurements near periods of change increase model performance?

Answer 1: Model performance increases when measurements are placed near periods of change

Question 2: How to space measurements when the nature of change is unknown?

Answer 2: When the nature of change is unknown, model performance is highest with equal spacing

Experiment 2

Question: For each spacing schedule, what combination of measurement number and sample size is needed to obtain high model performance (i.e., low bias, high precision)?

Experiment 2 – Systematic Review

Number of measurements

Spacing of measurements

Experiment 2 – Systematic Review

Number of measurements

Spacing of measurements

Sample size

Experiment 2 – Systematic Review

Number of measurements

Spacing of measurements

Sample size



Few guidelines


Effect	Nonlinear pattern
Main effects	
Number of measurements (NM)	6 studies
Spacing of measurements (SM)	1 study
Sample size (S)	7 studies

Effect	Nonlinear pattern	
Main effects		
Number of measurements (NM)	6 studies	
Spacing of measurements (SM)	1 study	
Sample size (S)	7 studies	
Two-way interactions		
NM x SM	1 study	
NM x S	5 studies	
SM x S	Cell 5 (Exp. 2)	
TS x S	2 studies	

Effect	Nonlinear pattern
Main effects	
Number of measurements (NM)	6 studies
Spacing of measurements (SM)	1 study
Sample size (S)	7 studies
Two-way interactions	
NM x SM	1 study
NM x S	5 studies
SM x S	Cell 5 (Exp. 2)
TS x S	2 studies
Three-way interactions	
NM x SM x S	Cell 9 (Exp. 2)


Effect	Nonlinear pattern
Main effects	
Number of measurements (NM)	6 studies
Spacing of measurements (SM)	1 study
Sample size (S)	7 studies
Two-way interactions	
NM x SM	1 study
NM x S	5 studies
SM x S	Cell 5 (Exp. 2)
TS x S	2 studies
Three-way interactions	
NM x SM x S	Cell 9 (Exp. 2)

Experiment 2 design (Monte Carlo method)



1. Population definition

$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (known)




2. Sample generation


IV 1: Number of measurements (5, 7, 9, 11)

IV 2: Spacing of measurements (equal, time-inc., time dec., middle-extreme)

IV 3: Sample size ($N = 30, 50, 100, 200, 500, 1000$)




$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (estimated)



4. Model performance

Bias + precision



3. Modelling

Structured latent growth model

Experiment 2

Question: For each spacing schedule, what combination of measurement number and sample size is needed to obtain high model performance (i.e., low bias, high precision)?

Experiment 2

Question: For each spacing schedule, what combination of measurement number and sample size is needed to obtain high model performance (i.e., low bias, high precision)?

Answer: The greatest improvements in model performance for each spacing schedule result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.

Answer: The greatest improvements in model performance for each spacing schedule result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.

Spacing Schedule	Qualitative Description
Equal (see Figure 3.2 and Table 3.1)	
Time-interval increasing (see Figure 3.3 and Table 3.3)	
Time-interval decreasing (see Figure 3.4 and Table 3.5)	

Answer: The greatest improvements in model performance for each spacing schedule result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.

Spacing Schedule	Qualitative Description
Equal (see Figure 3.2 and Table 3.1)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$
Time-interval increasing (see Figure 3.3 and Table 3.3)	
Time-interval decreasing (see Figure 3.4 and Table 3.5)	

Answer: The greatest improvements in model performance for each spacing schedule result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.

Spacing Schedule	Qualitative Description
Equal (see Figure 3.2 and Table 3.1)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$
Time-interval increasing (see Figure 3.3 and Table 3.3)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$
Time-interval decreasing (see Figure 3.4 and Table 3.5)	

Answer: The greatest improvements in model performance for each spacing schedule result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.

Spacing Schedule	Qualitative Description
Equal (see Figure 3.2 and Table 3.1)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$
Time-interval increasing (see Figure 3.3 and Table 3.3)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$
Time-interval decreasing (see Figure 3.4 and Table 3.5)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$

Answer: The greatest improvements in model performance for each spacing schedule result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.

Error Bar Summary		
Spacing Schedule	Qualitative Description	β_{fixed}
Equal (see Figure 3.2 and Table 3.1)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	12.67
Time-interval increasing (see Figure 3.3 and Table 3.3)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	13.27
Time-interval decreasing (see Figure 3.4 and Table 3.5)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	13.41

Answer: The greatest improvements in model performance for each spacing schedule result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.



		Error Bar Summary	
Spacing Schedule	Qualitative Description	β_{fixed}	γ_{fixed}
Equal (see Figure 3.2 and Table 3.1)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	12.67	9.79
Time-interval increasing (see Figure 3.3 and Table 3.3)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	13.27	9.69
Time-interval decreasing (see Figure 3.4 and Table 3.5)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	13.41	9.62

Answer: The greatest improvements in model performance for each spacing schedule result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.



		Error Bar Summary		
Spacing Schedule	Qualitative Description	β_{fixed}	γ_{fixed}	β_{random}
Equal (see Figure 3.2 and Table 3.1)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	12.67	9.79	16.02
Time-interval increasing (see Figure 3.3 and Table 3.3)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	13.27	9.69	16.28
Time-interval decreasing (see Figure 3.4 and Table 3.5)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	13.41	9.62	16.16

Answer: The greatest improvements in model performance for each spacing schedule result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.



		Error Bar Summary			
Spacing Schedule	Qualitative Description	β_{fixed}	γ_{fixed}	β_{random}	γ_{random}
Equal (see Figure 3.2 and Table 3.1)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	12.67	9.79	16.02	10.08
Time-interval increasing (see Figure 3.3 and Table 3.3)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	13.27	9.69	16.28	10.15
Time-interval decreasing (see Figure 3.4 and Table 3.5)	Largest improvements in bias and precision using NM = 7 with $N \geq 200$ or NM = 9 with $N \leq 100$	13.41	9.62	16.16	10.32

Experiment 3



Experiment 3

Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

Experiment 3

Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

- 1) What is time structuredness?
- 2) Why is time structuredness important?

Experiment 3


Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

- 1) **What is time structuredness?**
- 2) Why is time structuredness important?

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time


Time-structured data (same response schedule)



Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time


Time-structured data (same response schedule)



Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time


Time-structured data (same response schedule)



Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time


Time-structured data (same response schedule)



Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time


Time-structured data (same response schedule)



Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time


Time-structured data (same response schedule)



Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time

Time-structured data (same response schedule)




Time-unstructured data


Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time

Time-structured data (same response schedule)




Time-unstructured data (different response schedules)




Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time

Time-structured data (same response schedule)




Time-unstructured data (different response schedules)




Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time

Time-structured data (same response schedule)




Time-unstructured data (different response schedules)




Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time

Time-structured data (same response schedule)




Time-unstructured data (different response schedules)




Mehta & West (2000); Mehta & Neale (2005)

Time structuredness: the extent to which the same response schedule characterizes how participants provide data over time

Time-structured data (same response schedule)



Time-unstructured data (different response schedules)



Mehta & West (2000); Mehta & Neale (2005)


Experiment 3


Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

1) What is time structuredness?


The extent to which the same response schedule characterizes how participants provide data over time

2) Why is time structuredness important?





By assuming one response pattern (i.e., time-structured data), individual curves are shifted to fit this response pattern and result in considerable error



Experiment 3

Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

1) What is time structuredness?

The extent to which the same response schedule characterizes how participants provide data over time

2) Why is time structuredness important?

Considerable error can be incurred if a model assuming time-structured data is used to analyse time-unstructured data

Experiment 3 – Systematic Review

- Number of measurements
 - Spacing of measurements
 - Sample size
 - Time structuredness
- 
- Few guidelines


Effect	Nonlinear pattern
Main effects	
Number of measurements (NM)	6 studies
Spacing of measurements (SM)	1 study
Time structuredness (TS)	1 study
Sample size (S)	7 studies

Effect	Nonlinear pattern
Main effects	
Number of measurements (NM)	6 studies
Spacing of measurements (SM)	1 study
Time structuredness (TS)	1 study
Sample size (S)	7 studies
Two-way interactions	
NM x SM	1 study
NM x TS	Cell 1 (Exp. 3)
NM x S	5 studies
SM x TS	Cell 3
SM x S	Cell 5 (Exp. 2)
TS x S	2 studies

Effect	Nonlinear pattern
Main effects	
Number of measurements (NM)	6 studies
Spacing of measurements (SM)	1 study
Time structuredness (TS)	1 study
Sample size (S)	7 studies
Two-way interactions	
NM x SM	1 study
NM x TS	Cell 1 (Exp. 3)
NM x S	5 studies
SM x TS	Cell 3
SM x S	Cell 5 (Exp. 2)
TS x S	2 studies
Three-way interactions	
NM x SM x TS	Cell 7
NM x SM x S	Cell 9 (Exp. 2)
NM x TS x S	Cell 10 (Exp. 3)
SM x TS x S	Cell 12


Effect	Nonlinear pattern
Main effects	
Number of measurements (NM)	6 studies
Spacing of measurements (SM)	1 study
Time structuredness (TS)	1 study
Sample size (S)	7 studies
Two-way interactions	
NM x SM	1 study
NM x TS	Cell 1 (Exp. 3)
NM x S	5 studies
SM x TS	Cell 3
SM x S	Cell 5 (Exp. 2)
TS x S	2 studies
Three-way interactions	
NM x SM x TS	Cell 7
NM x SM x S	Cell 9 (Exp. 2)
NM x TS x S	Cell 10 (Exp. 3)
SM x TS x S	Cell 12

Experiment 3 design (Monte Carlo method)



1. Population definition

$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (known)




2. Sample generation


IV 1: Number of measurements (5, 7, 9, 11)

IV 2: Spacing of measurements (equal, time-inc., time dec., mid-extreme)

IV 3: Time structuredness




$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (estimated)



4. Model performance


Bias + precision



3. Modelling

Structured latent growth model

Time-structured data (High time structuredness)



Momentary response window (immediate response rate)

High

Short

Fast

Time structuredness

Response window length

Response rate


Low

Long


Slow

Coulombe (2016)

*Time-structured data
(High time structuredness)*



Medium time structuredness



*Momentary response window
(immediate response rate)*

High

Short

Fast

*Short response window
(fast response rate)*

Time structuredness

Response window length

Response rate


Low

Long

Slow

Coulombe (2016)

*Time-structured data
(High time structuredness)*




*Momentary response window
(immediate response rate)*

High

Short


Fast

Medium time structuredness



*Short response window
(fast response rate)*

Low time structuredness



*Long response window
(slow response rate)*

Time structuredness
Response window length
Response rate


Low

Long

Slow


Coulombe (2016)

Experiment 3 design (Monte Carlo method)



1. Population definition

$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (known)




2. Sample generation


IV 1: Number of measurements (5, 7, 9, 11)

IV 2: Spacing of measurements (equal, time-inc., time dec., mid-extreme)

IV 3: Time structuredness (time structured, fast response rate, slow response rate)




$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (estimated)



4. Model performance

Bias + precision



3. Modelling

Structured latent growth model

Experiment 3


Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

Experiment 3

Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

Answer: Bias systematically increases as time structuredness decreases

Answer: Bias systematically increases as time structuredness decreases



Answer: Bias systematically increases as time structuredness decreases

High

Fast


*Time structuredness
Response rate*

Low

Slow


*Time structured data
(immediate response rate)*

A : β_{fixed} (Days-to-Halfway Elevation)



Fast response rate

A : β_{fixed} (Days-to-Halfway Elevation)



Sample Size (N)

Number of Measurements	■ 5	● 7	▲ 9	◆ 11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		

Answer: Bias systematically increases as time structuredness decreases


High
Fast

Time structuredness
Response rate

Low
Slow


Time structured data
(immediate response rate)

A : β_{fixed} (Days-to-Halfway Elevation)




Fast response rate

A : β_{fixed} (Days-to-Halfway Elevation)



Slow response rate

A : β_{fixed} (Days-to-Halfway Elevation)



Sample Size (N)

Number of Measurements	■ 5	● 7	▲ 9	◆ 11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		

Answer: Bias systematically increases as time structuredness decreases


High
Fast

Time structuredness
Response rate

Low
Slow


Time structured data
(immediate response rate)

C : β_{random} (Days-to-Halfway Elevation)




Fast response rate

C : β_{random} (Days-to-Halfway Elevation)



Slow response rate


C : β_{random} (Days-to-Halfway Elevation)



Sample Size (N)


Number of Measurements	5	7	9	11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		

Answer: Bias systematically increases as time structuredness decreases




Number of Measurements	5	7	9	11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		


By assuming one response pattern, all individual curves are shifted to fit this response pattern




By assuming one response pattern, all individual curves are shifted to fit this response pattern



By assuming one response pattern, all individual curves are shifted to fit this response pattern




Experiment 3 design (Monte Carlo method)



1. Population definition

$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (known)




2. Sample generation


IV 1: Number of measurements (5, 7, 9, 11)

IV 2: Spacing of measurements (equal, time-inc., time dec., mid-extreme)

IV 3: Time structuredness (time structured, fast response rate, slow response rate)




$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (estimated)



4. Model performance


Bias + precision



3. Modelling


Structured latent growth model

Experiment 3 design (Monte Carlo method)



1. Population definition

$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (known)




2. Sample generation


IV 1: Number of measurements (5, 7, 9, 11)

IV 2: Spacing of measurements (equal, time-inc., time dec., mid-extreme)

Constant: slow response rate




$\theta, \alpha, \beta, \gamma_{\text{fixed + random}}, \varepsilon$ (estimated)



4. Model performance

Bias + precision



3. Modelling

Structured latent growth model +
definition variables

Definition variables can be used to prevent low time structuredness from inflating bias


High
Fast

Time structuredness
Response rate

Low
Slow


Time structured data
(immediate response rate)

A : β_{fixed} (Days-to-Halfway Elevation)




Fast response rate

A : β_{fixed} (Days-to-Halfway Elevation)



Slow response rate

A : β_{fixed} (Days-to-Halfway Elevation)



Sample Size (N)

Number of Measurements	5	7	9	11
Is Unbiased?	■ Yes	<input type="checkbox"/> No		
Is Precise?	— Yes	<input type="line"/> No		

Definition variables can be used to prevent low time structuredness from inflating bias


High
Fast

Time structuredness
Response rate

Low
Slow


Time structured data
(immediate response rate)

A : β_{fixed} (Days-to-Halfway Elevation)




Fast response rate

A : β_{fixed} (Days-to-Halfway Elevation)



Slow response rate +
definition variables

A : β_{fixed} (Days-to-Halfway Elevation)



Sample Size (N)

Number of Measurements	5	7	9	11
Is Unbiased?	■ Yes	<input type="checkbox"/> No		
Is Precise?	— Yes	<input type="blue"/> No		


Definition variables can be used to prevent low time structuredness from inflating bias

High
Fast


Time structuredness
Response rate

Low
Slow


Time structured data
(immediate response rate)



Fast response rate



Slow response rate



Number of Measurements	■ 5	● 7	▲ 9	◆ 11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		


Definition variables can be used to prevent low time structuredness from inflating bias

High
Fast


Time structuredness
Response rate

Low
Slow


Time structured data
(immediate response rate)



Fast response rate



Slow response rate +
definition variables



Number of Measurements	■ 5	● 7	▲ 9	◆ 11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		

Definition variables can be used to prevent low time structuredness from inflating bias


High
Fast

Time structuredness
Response rate

Low
Slow


Time structured data
(immediate response rate)

D : γ_{random} (Triquarter-Halfway Delta)




Fast response rate

D : γ_{random} (Triquarter-Halfway Delta)



Slow response rate

D : γ_{random} (Triquarter-Halfway Delta)



Sample Size (N)

Number of Measurements	5	7	9	11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		

Definition variables can be used to prevent low time structuredness from inflating bias


High
Fast

Time structuredness
Response rate

Low
Slow


Time structured data
(immediate response rate)

D : γ_{random} (Triquarter-Halfway Delta)




Fast response rate

D : γ_{random} (Triquarter-Halfway Delta)



Slow response rate + definition variables

D : γ_{random} (Triquarter-Halfway Delta)



Number of Measurements	5	7	9	11
Is Unbiased?	■ Yes	□ No		
Is Precise?	— Yes	— No		

Experiment 3

Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

Experiment 3

Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

Answer: Bias systematically increases as time structuredness decreases

Experiment 3

Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

Answer: Bias systematically increases as time structuredness decreases

Definition variables can be used to prevent low time structuredness from increasing in bias.

Experiment 3

Question: Do the number of measurements and sample sizes needed to obtain high model performance (i.e., low bias, high precision) increase as time structuredness decreases?

Answer: Bias systematically increases as time structuredness decreases

Definition variables can be used to prevent low time structuredness from increasing in bias.

If using definition variable, the largest improvements in model performance result from using either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.

Experiment 1

Model performance increases when measurements are placed near periods of change

When the nature of change is unknown, model performance is highest with equal spacing

Experiment 1

Model performance increases when measurements are placed near periods of change

When the nature of change is unknown, model performance is highest with equal spacing

Experiment 2

The greatest improvements in model performance result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.

Experiment 1

Model performance increases when measurements are placed near periods of change

When the nature of change is unknown, model performance is highest with equal spacing

Experiment 2

The greatest improvements in model performance result from either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.

Experiment 3

If using definition variables, the largest improvements in model performance result from using either seven measurements with $N \geq 200$ or nine measurements with $N \leq 100$.

The results of my experiments are the first (to my knowledge) to provide recommendations for modelling nonlinear change

The results of my experiments are the first (to my knowledge) to provide recommendations for modelling nonlinear change

Number of measurements

Spacing of measurements

Sample size

Time structuredness



Few guidelines

Limitations & future directions


- 1) Cutoff values
- 2) External validity

Limitations & future directions

1) Cutoff values

2) External validity

I used a percentage-based cutoff rule in my experiments



Bias: difference between average estimated value and population value

Estimates were biased if bias > 10% of population value

Precision: range of values covered by the middle 95% of estimates

Estimation was imprecise if a whisker length > 10% of population value

Muthén et al. (1997)

Percentage-based cutoff values are context dependent

A 10% increase in heart rate does not lead to the same health outcomes as a 10% increase in body temperature



Muthén et al. (1997)

Percentage-based cutoff values are context dependent

A 10% increase in heart rate does not lead to the same health outcomes as a 10% increase in body temperature

10% increase in heart rate = 7–8 beats/minute



10% increase body temperature = 3.7°C



Future research could use smallest cutoff values of interest to account for the context

Future research could use smallest cutoff values of interest to account for the context

Smallest effect sizes of interest → Test for meaningful effect sizes

Future research could use smallest cutoff values of interest to account for the context

Smallest effect sizes of interest → Test for meaningful effect sizes

Smallest cutoff values of interest → Test for meaningful changes in model performance

Limitations & future directions

1) Cutoff values

2) External validity

The current simulation experiments generated data under ideal conditions

- 1) Data generation assumed 100% response rate

The current simulation experiments generated data under ideal conditions

1) Data generation assumed 100% response rate

Data are estimated to be missing 30% of the time, and increase over time

The current simulation experiments generated data under ideal conditions

- 1) Data generation assumed 100% response rate

Data are estimated to be missing 30% of the time, and increase over time

- 2) Model assumed a constant measurement model over time (i.e., measurement invariance)

The current simulation experiments generated data under ideal conditions

1) Data generation assumed 100% response rate

Data are estimated to be missing 30% of the time, and increase over time

2) Model assumed a constant measurement model over time (i.e., measurement invariance)

Measurement invariance is seldom observed

The current simulation experiments generated data under ideal conditions

- 1) Data generation assumed 100% response rate

Data are estimated to be missing 30% of the time, and increase over time

- 2) Model assumed a constant measurement model over time (i.e., measurement invariance)

Measurement invariance is seldom observed

- 3) Error variances assumed to be constant and uncorrelated over time

The current simulation experiments generated data under ideal conditions

- 1) Data generation assumed 100% response rate

Data are estimated to be missing 30% of the time, and increase over time

- 2) Model assumed a constant measurement model over time (i.e., measurement invariance)

Measurement invariance is seldom observed

- 3) Error variances assumed to be constant and uncorrelated over time

Error variances are likely to be heterogeneous and correlated.

McKnight et al. (2007); Newman (2003); Mellenbergh (1989); Van De Schoot et al. (2015); Bliese & Ployhart (2002)

A gap exists between researchers and practitioners whereby research findings seldom make their way into practice



A gap exists between researchers and practitioners whereby research findings seldom make their way into practice

Poor understanding of organizational psychology among managers

A gap exists between researchers and practitioners whereby research findings seldom make their way into practice

Poor understanding of organizational psychology among managers

~55% of practitioners skeptical that organizational psychology can effect positive change

One factor that may contribute to the academic-practitioner gap is the lack of specific recommendations

One factor that may contribute to the academic-practitioner gap is the lack of specific recommendations

Organizational theory typically specifies non-zero relations

One factor that may contribute to the academic-practitioner gap is the lack of specific recommendations

Organizational theory typically specifies non-zero relations




~3% of human resource articles address problems faced by practitioners

Edwards & Berry (2010); Sackett & Larson (1990)

Modelling nonlinear change could provide one avenue for developing specific recommendations

Modelling nonlinear change could provide one avenue for developing specific recommendations



References

- Bliese, P. D., Kautz, J., & Lang, J. W. B. (2020). Discontinuous growth models: Illustrations, recommendations, and an R function for generating the design matrix. In Y. Griep & S. D. Hansen (Eds.), *Handbook on the temporal dynamics of organizational behavior* (pp. 319–350). Edward Elgar Publishing. <https://doi.org/10.4337/9781788974387.00030>
- Bliese, P. D., & Ployhart, R. E. (2002). Growth modeling using random coefficient models: Model building, testing, and illustrations. *Organizational Research Methods*, 5(4), 362–387. <https://doi.org/10.1177/109442802237116>
- Browne, M. W. (1993). Structured latent curve models. In C. M. Cuadras & C. R. Rao (Eds.), *Multivariate analysis: Future directions 2* (1st ed., pp. 171–197). Elsevier. <https://doi.org/10.1016/B978-0-444-81531-6.50016-7>
- Browne, M. W., & du Toit, S. H. C. (1991). Models for learning data. In L. M. Collins & J. L. Horn (Eds.), *Best methods for the analysis of change: Recent advances, unanswered questions, future directions* (pp. 47–68). American Psychological Association. <https://doi.org/10.1037/10099-004>
- Burchinal, M., & Appelbaum, M. I. (1991). Estimating individual developmental functions: Methods and their assumptions. *Child Development*, 62(1), 23–42. <https://doi.org/10.2307/1130702>

References

- Chen, G., Ployhart, R. E., Thomas, H. C., Anderson, N., & Bliese, P. D. (2011). The power of momentum: A new model of dynamic relationships between job satisfaction change and turnover intentions. *Academy of Management Journal*, 54(1), 159–181.
<https://doi.org/10.5465/amj.2011.59215089>
- Christian, M. S., & Ellis, A. P. J. (2011). Examining the effects of sleep deprivation on workplace deviance: A self-regulatory perspective. *Academy of Management Journal*, 54(5), 913–934. <https://doi.org/10.5465/amj.2010.0179>
- Coulombe, P., Selig, J. P., & Delaney, H. D. (2016). Ignoring individual differences in times of assessment in growth curve modeling. *International Journal of Behavioral Development*, 40(1), 76–86. <https://doi.org/10.1177/0165025415577684>
- Edwards, J. R., & Berry, J. W. (2010). The presence of something or the absence of nothing: Increasing theoretical precision in management research. *Organizational Research Methods*, 13(4), 668–689. <https://doi.org/10.1177/1094428110380467>
- Fournier, M., d'Arripe-Longueville, F., Rovere, C., Easthope, C. S., Schwabe, L., El Methni, J., & Radel, R. (2017). Effects of circadian cortisol on the development of a health habit. *Health Psychology*, 36(11), 1059–1064.
<https://doi.org/10.1037/he0000510>

References

- Lakens, D. (2017). Equivalence tests: A practical primer for *t* tests, correlations, and meta-analyses. *Social Psychological and Personality Science*, 8(4), 355–362. <https://doi.org/10.1177/1948550617697177>
- Lakens, D., Scheel, A. M., & Isager, P. M. (2018). Equivalence testing for psychological research: A tutorial. *Advances in Methods and Practices in Psychological Science*, 1(2), 259–269. <https://doi.org/10.1177/2515245918770963>
- Lawrence, T. B., Winn, M. I., & Jennings, P. D. (2001). The temporal dynamics of institutionalization. *Academy of Management Review*, 26(4), 624–644. <https://doi.org/10.5465/amr.2001.5393901>
- Li, L. (2017). Sentiment-enhanced learning model for online language learning system. *Electronic Commerce Research*, 18(1), 23–64. <https://doi.org/10.1007/s10660-017-9284-5>
- Mayer, R. C., Davis, J. H., & Schoorman, F. D. (1995). An integrative model of organizational trust. *The Academy of Management Review*, 20(3), 709. <https://doi.org/10.2307/258792>
- McKnight, P., McKnight, P. K., Sidani, S., & Figueredo, A. J. (2007). *Missing data: A gentle introduction*. Guilford Press. <https://bit.ly/3i5of1W>
- Mehta, P. D., & Neale, M. C. (2005). People are variables too: Multilevel structural equations modeling. *Psychological Methods*, 10(3), 259–284. <https://doi.org/10.1037/1082-989x.10.3.259>

References

- Mehta, P. D., & West, S. G. (2000). Putting the individual back into individual growth curves. *Psychological Methods*, 5(1), 23–43. <https://doi.org/10.1037/1082-989x.5.1.23>
- Mellenbergh, G. J. (1989). Item bias and item response theory. *International Journal of Educational Research*, 13(2), 127–143. [https://doi.org/10.1016/0883-0355\(89\)90002-5](https://doi.org/10.1016/0883-0355(89)90002-5)
- Moran, D. S., & Mendal, L. (2002). Core temperature measurement. *Sports Medicine*, 32(14), 879–885. <https://doi.org/10.2165/00007256-200232140-00001>
- Muthén, B. O., du Toit, S. H. C., & Spisic, D. (1997). *Robust inference using weighted least squares and quadratic estimating equations in latent variable modeling with categorical and continuous outcomes* [Unpublished technical report]. <https://bit.ly/3Q3Yew>
- Nanchen, D. (2018). Resting heart rate: What is normal? *Heart*, 104(13), 1048–1049. <https://doi.org/10.1136/heartjnl-2017-312731>
- Navarro, J., Roe, R. A., & Artiles, M. (2015). Taking time seriously: Changing practices and perspectives in work/organizational psychology. *Journal of Work and Organizational Psychology*, 31(3), 135–145. <https://doi.org/10.1016/j.rpto.2015.07.002>
- Newman, D. A. (2003). Longitudinal modeling with randomly and systematically missing data: A simulation of ad hoc, maximum likelihood, and multiple imputation techniques. *Organizational Research Methods*, 6(3), 328–362. <https://doi.org/10.1177/1094428103254673>

References

- Pickup, M. (2014). *Introduction to time series analysis*. Sage Publications. <https://bit.ly/3WDtdll>
- Rynes, S. L., Colbert, A. E., & Brown, K. G. (2002). HR professionals' beliefs about effective human resource practices: Correspondence between research and practice. *Human Resource Management*, 41(2), 149–174. <https://doi.org/10.1002/hrm.10029>
- Sackett, P. R., & Larson, J., J. R. (1990). Research strategies and tactics in industrial and organizational psychology. In M. D. Dunnette & L. M. Hough (Eds.), *Handbook of industrial and organizational psychology* (pp. 419–489). Consulting Psychologists Press. <https://bit.ly/3GymBiR>
- Sanders, K., van Riemsdijk, M., & Groen, B. (2008). The gap between research and practice: A replication study on the hr professionals' beliefs about effective human resource practices. *The International Journal of Human Resource Management*, 19(10), 1976–1988. <https://doi.org/10.1080/09585190802324304>
- Spears, M., & Bolton, R. (2015). *Evidence-based hr: The bridge between your people and delivering business strategy*. KPMG International.
- Van De Schoot, R., Schmidt, P., De Beuckelaer, A., Lek, K., & Zondervan-Zwijnenburg, M. (2015). Editorial: Measurement invariance. *Frontiers in Psychology*, 6, 1064. <https://doi.org/10.3389/fpsyg.2015.01064>

References

- van der Nest, G., Lima Passos, V., Candel, M. J., & van Breukelen, G. J. (2020). An overview of mixture modelling for latent evolutions in longitudinal data: Modelling approaches, fit statistics and software. *Advances in Life Course Research*, 43, 100323. <https://doi.org/10.1016/j.alcr.2019.100323>
- Vancouver, J. B., Wang, M., & Li, X. (2020). Translating informal theories into formal theories: The case of the dynamic computational model of the integrated model of work motivation. *Organizational Research Methods*, 23(2), 238–274. <https://doi.org/10.1177/1094428118780308>
- Whitley, J. D., & Schoene, L. L. (1987). Comparison of heart rate responses. *Physical Therapy*, 67(10), 1501–1504. <https://doi.org/10.1093/ptj/67.10.1501>
- Yang, Y., & Li, X. (2021). The impact of challenge and hindrance stressors on thriving at work double mediation based on affect and motivation. *Frontiers in Psychology*, 12. <https://doi.org/10.3389/fpsyg.2021.613871>

Break
Slide.
