

MATHS7027 Mathematical Foundations of Data Science Assignment 5

Trimester 2, 2024

Dongju Ma

July 28, 2024

Assignment 5 - Question 1

To find the eigenvalues and eigenspaces of A , we could consider an expression follow:

$$Ax = \lambda x, x \neq 0$$

According to the concept of characteristic equation, we could write:

$$\det(\lambda I - A) = 0$$

And we could get:

$$\begin{vmatrix} \lambda - 1 & 0 & 2 \\ 0 & \lambda - 3 & 0 \\ 2 & 0 & \lambda - 1 \end{vmatrix} = 0$$

Calculate the determinant:

$$\begin{aligned} \begin{vmatrix} \lambda - 1 & 0 & 2 \\ 0 & \lambda - 3 & 0 \\ 2 & 0 & \lambda - 1 \end{vmatrix} &= (\lambda - 1) \begin{vmatrix} \lambda - 3 & 0 \\ 0 & \lambda - 1 \end{vmatrix} - 0 + 2 \begin{vmatrix} 0 & \lambda - 3 \\ 2 & 0 \end{vmatrix} \\ &= (\lambda - 1)(\lambda - 3)(\lambda - 1) - 4(\lambda - 3) \\ &= (\lambda^2 - 2\lambda + 1 - 4)(\lambda - 3) \\ &= (\lambda^2 - 2\lambda - 3)(\lambda - 3) \\ &= (\lambda + 1)(\lambda - 3)(\lambda - 3) \end{aligned}$$

So we could find the eigenvalues of A is $\lambda = -1, 3, 3$

Next we solve the characteristic equations to find the corresponding eigenvectors:

Let $\lambda = -1$. Solve $(\lambda I - A)x = 0$ where $x = (x_1, x_2, x_3)$, and $\lambda = -1$. Which is,

$$\begin{bmatrix} -2 & 0 & 2 \\ 0 & -4 & 0 \\ 2 & 0 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{array}{ccc|c} -2 & 0 & 2 & 0 \\ 0 & -4 & 0 & 0 \\ 2 & 0 & -2 & 0 \\ \hline 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \quad \begin{array}{l} (R_3 = R_3 + R_1) \\ (R_1 = -\frac{1}{2}R_1) \\ (R_2 = -\frac{1}{4}R_2) \end{array}$$

$$\therefore x_1 = x_3 = t$$

$$x_2 = 0$$

The corresponding eigenvalue of $\lambda = -1$ is $x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = t \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$

Let $\lambda = 3$. Solve $(\lambda I - A)x = 0$ where $x = (x_1, x_2, x_3)$, and $\lambda = 3$. Which is,

$$\begin{bmatrix} 2 & 0 & 2 \\ 0 & 0 & 0 \\ 2 & 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{array}{ccc|c} 2 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 \\ 2 & 0 & 2 & 0 \\ \hline 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \quad \begin{array}{l} (R_3 = R_3 - R_1) \\ (R_1 = \frac{1}{2}R_1) \end{array}$$

$$\therefore x_1 = t$$

$$\Rightarrow x_3 = -t$$

$$x_2 = 0$$

The corresponding eigenvalue of $\lambda = 3$ is $x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = t \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$

Assignment 5 - Question 2

2(a)

According to the concept of eigenvectors, we could get equations below

$$Bv_1 = \lambda_1 v_1$$

$$Bv_2 = \lambda_2 v_2$$

So we calculate Bv_1 :

$$Bv_1 = \begin{bmatrix} 2 & a & -1 \\ 0 & 2 & b \\ -1 & -1 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix} = \begin{bmatrix} 2-a \\ -2 \\ 0 \end{bmatrix}$$

Since $Bv_1 = \lambda_1 v_1$, so we have

$$\begin{bmatrix} 2-a \\ -2 \\ 0 \end{bmatrix} = \begin{bmatrix} \lambda_1 \\ -\lambda_1 \\ 0 \end{bmatrix}$$

Then we calculate Bv_2

$$Bv_2 = \begin{bmatrix} 2 & a & -1 \\ 0 & 2 & b \\ -1 & -1 & 3 \end{bmatrix} \begin{bmatrix} -1 \\ -1 \\ 2 \end{bmatrix} = \begin{bmatrix} -4-a \\ 2b-2 \\ 8 \end{bmatrix}$$

Since $Bv_2 = \lambda_2 v_2$, so we have

$$\begin{bmatrix} -4-a \\ 2b-2 \\ 8 \end{bmatrix} = \begin{bmatrix} -\lambda_2 \\ -\lambda_2 \\ 2\lambda_2 \end{bmatrix}$$

We could easily obtain that $\lambda_1 = 2, \lambda_2 = 4$, and substitute them into the equations,

$$2-a=2$$

$$-4-a=-4 \Rightarrow a=0, b=-1$$

$$2b-2=-4$$

So $a=0, b=-1$

2(b)

The trace of B is the sum of its diagonal elements.

So we could get

$$\text{tr}(B) = 2 + 2 + 3 = 7$$

Let λ_3 be the remaining eigenvalue, and the sum of the eigenvalues is equal to the trace.

So

$$\lambda_3 = 7 - 2 - 4 = 1$$

Assignment 5 - Question 3

3(a)

We could add the matrix I to the right of P , and then change it into the form $[I|P^{-1}]$:

$$\left[\begin{array}{ccc|ccc} 1 & -1 & 0 & 1 & 0 & 0 \\ 2 & 2 & 3 & 0 & 1 & 0 \\ 2 & 0 & 1 & 0 & 0 & 1 \end{array} \right]$$

Then we apply the row reduction to change the P to I :

$$\begin{aligned}
& \left[\begin{array}{ccc|ccc} 2 & 0 & 1 & 0 & 0 & 1 \\ 2 & 2 & 3 & 0 & 1 & 0 \\ 1 & -1 & 0 & 1 & 0 & 0 \end{array} \right] & (\text{Swap } R_3 \text{ and } R_1) \\
& \left[\begin{array}{ccc|ccc} 2 & 0 & 1 & 0 & 0 & 1 \\ 0 & 2 & 2 & 0 & 1 & -1 \\ 1 & -1 & 0 & 1 & 0 & 0 \end{array} \right] & (R_2 = R_2 - R_1) \\
& \left[\begin{array}{ccc|ccc} 2 & 0 & 1 & 0 & 0 & 1 \\ 0 & 2 & 2 & 0 & 1 & -1 \\ 0 & -1 & -\frac{1}{2} & 1 & 0 & -\frac{1}{2} \end{array} \right] & (R_3 = R_3 - \frac{1}{2}R_1) \\
& \left[\begin{array}{ccc|ccc} 2 & 0 & 1 & 0 & 0 & 1 \\ 0 & 2 & 2 & 0 & 1 & -1 \\ 0 & 0 & \frac{1}{2} & 1 & \frac{1}{2} & -1 \end{array} \right] & (R_3 = R_3 + \frac{1}{2}R_2) \\
& \left[\begin{array}{ccc|ccc} 2 & 0 & 1 & 0 & 0 & 1 \\ 0 & 2 & 2 & 0 & 1 & -1 \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array} \right] & (R_3 = 2R_3) \\
& \left[\begin{array}{ccc|ccc} 2 & 0 & 1 & 0 & 0 & 1 \\ 0 & 2 & 0 & -4 & -1 & 3 \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array} \right] & (R_2 = R_2 - 2R_3) \\
& \left[\begin{array}{ccc|ccc} 2 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -2 & -\frac{1}{2} & \frac{3}{2} \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array} \right] & (R_2 = \frac{1}{2}R_2) \\
& \left[\begin{array}{ccc|ccc} 2 & 0 & 0 & -2 & -1 & 3 \\ 0 & 1 & 0 & -2 & -\frac{1}{2} & \frac{3}{2} \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array} \right] & (R_1 = R_1 - R_3) \\
& \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -1 & -\frac{1}{2} & \frac{3}{2} \\ 0 & 1 & 0 & -2 & -\frac{1}{2} & \frac{3}{2} \\ 0 & 0 & 1 & 2 & 1 & -2 \end{array} \right] & (R_1 = \frac{1}{2}R_1)
\end{aligned}$$

3(b)

As C is diagonal, we could get:

$$C = PDP^{-1}$$

for any positive integer n , there is C^n :

$$C^n = (PDP^{-1})^n = PD^nP^{-1}$$

$$\text{As } D = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & c \end{bmatrix}, \text{ so,}$$

$$D^n = \begin{bmatrix} 1^n & 0 & 0 \\ 0 & 0^n & 0 \\ 0 & 0 & c^n \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & c^n \end{bmatrix}$$

So C^n should be

$$C^n = PD^nP^{-1} = P \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & c^n \end{bmatrix} P^{-1}$$