



Primary Examination

!! PRACTICE EXAM !!

MATHS 7107 Data Taming

Course Coordinator: Anthony Mays

Writing time: 120 mins

Number of questions: 7 Written, 16 Multiple Choice

Total marks: ??

Instructions for Candidates

- This examination is **closed book**.
- Examination materials must not be removed from the examination room.
- Answer all written questions in the answer book provided.
- Answer all multiple choice questions on the provided multiple choice sheet.

Permitted Materials

- Multiple Choice Question (MCQ) Sheet
- Paper translation dictionary
- Scientific calculator.
- One double-sided A4 page of handwritten or typed notes.

**DO NOT COMMENCE WRITING UNTIL INSTRUCTED TO DO SO.
STOP WRITING IMMEDIATELY WHEN INSTRUCTED**

Written Questions

1. A committee has been established to organise 200-year anniversary celebrations in each Australian capital city. A large statue has been built and needs to be moved to each city for the celebration. The following tibble contains some of the information that the committee has gathered

```
# A tibble: 8 × 5
  Name                population urban_pop `Capital city` Year
  <chr>                <dbl>    <dbl> <fct>         <dbl>
1 New South Wales    7759274    64.8 Sydney         1788
2 Victoria          6179249    76.5 Melbourne      1851
3 Queensland         4848877    48.7 Brisbane       1860
4 Western Australia  2558951    79.0 Perth         1829
5 South Australia    1713054    77.3 Adelaide     1836
6 Tasmania           517588     43.4 Hobart       1826
7 Australian Capital Territory 403468    100 Canberra       1913
8 Northern Territory 245740     59.4 Darwin       1911
```

The variables in the table are:

- Name: the name of the state or territory
- population: the number of people residing in the state
- urban_pop: the percentage of the state's population who lives in the state's capital city
- Capital city: the name of the state's capital city
- Year: the year the city became the state's capital city, which is used for planning where the statue needs to be.

- (a) For each of the variables in the dataset identify the type of variable, ie. is it **quantitative continuous**, **quantitative discrete**, **categorical nominal** or **categorical ordinal**. Make sure you write a short description justifying your choice.
- (b) For each variable state whether the corresponding column in the tibble is the correct data type. If it is incorrect, say what the correct data type should be.




Figure 1: Log-Log plot of data for Question 2.

2. (a) Assume that a data set consists of continuous variables x and y , which are related by the formula

$$y = \alpha x^k.$$

Using the logarithm laws, show that the Log-Log plot of this data will be a straight line.

- (b) Using the straight line in part (a), write down the:
- gradient of the line,
 - the value at which the line cuts the vertical axis.
- (c) Using the Log-Log plot in Figure 1 determine the explicit relationship between x and y . (Make sure you show your working.)

x_1	x_2	x_3	x_4	x_5
1	0	2	-1	3

Table 1: Dataset for Questions 3 and 4.

3. For this question, use the dataset in Table 1.
- Transform the dataset in Table 1 to x^* by applying Min-Max scaling. Calculate your answers to 2 decimal places. **(Make sure you show your working.)**
 - What are the minimum and maximum values of the transformed dataset x^* ? Give exact values for your answers. **(Make sure you show your working.)**
4. For this question, use the dataset in Table 1.
- Find the mean \bar{x} and (sample) standard deviation σ_x of the dataset. Calculate your answers to 2 decimal places. **(Make sure you show your working.)**
 - Standardise the dataset by calculating the z -scores x_j^* for each x_j . Calculate your answers to 2 decimal places. **(Make sure you show your working.)**
 - What is the mean \bar{x}^* and standard deviation σ_{x^*} of the transformed data set. Give exact values for your answers. **(Make sure you show your working.)**
5. Show that when $x \neq 0$ the Box-Cox transformation is continuous at $\lambda = 0$, by showing that

$$\lim_{\lambda \rightarrow 0} \frac{x^\lambda - 1}{\lambda} = \log(x) \quad (\text{for } x \neq 0).$$

6. We are looking to measure the amount of pollen in the air. We have built 31 cubic boxes, each one with a pollen counter in it. The side lengths of each box is recorded in the list (s_1, \dots, s_{31}) , where the side lengths are measured in metres. We collect pollen counts from the air samples inside each box, and record the pollen counts in box j as p_j .

We expect to find a good linear fit with the model

$$p_j = \beta_0 + \beta_1 z_j + \epsilon_j, \quad N(0, \sigma) \quad (1)$$

where $z_j = s_j^3$.

We use R to fit our linear model to the data and we obtain the following output:

```
> pollen_lm <- lm(p ~ z, pollen)
> summary(pollen_lm)
```

Call:

```
lm(formula = p ~ z, data = pollen)
```

Residuals:

Min	1Q	Median	3Q	Max
-22.195	-6.719	-0.049	6.826	20.877

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	6.23517	3.59069	1.736	0.0931 .
z	1.31177	0.05785	22.675	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 11.27 on ##### degrees of freedom

Multiple R-squared: 0.9466, Adjusted R-squared: 0.9448

F-statistic: 514.1 on 1 and ##### DF, p-value: < 2.2e-16

- Note that we have replaced the number degrees of freedom with **#####**, and so you need to calculate this number.

- (a) Use the information in the output to write the equation for the line of best fit in Equation (1) with the appropriate values substituted. (Round off your answer to 4 decimal places.)

- (b) From the output, what is the estimate of the standard deviation of the errors? (Provide your answer to 2 decimal places.)

- (c) With this model, what is the prediction for the pollen count in a box of side length $4.3m$? (Round off your answer to the nearest whole number of pollen grains.)

- (d) Write down the details for the two t -tests performed when fitting this linear model. State the null and alternative hypotheses, and the distribution being tested against (including the number of degrees of freedom).

- (e) What are the relevant P-values for the tests? Are the estimates for the model parameters significant at the 95% level?

- (f) Use the graphs in Figures 2 – 4 to determine if the assumptions for fitting a linear model are satisfied. (Make sure you refer to specific Figures in your answers.) In addition, for the **independence** assumption, come up with at least one reason why the assumption may not be satisfied.




Figure 2: A residuals vs fitted graph for the linear model in Question 6.




Figure 3: A scale-location graph for the linear model in Question 6.




Figure 4: A Q-Q plot of residuals for the linear model in Question 6.

7. A medical test for a certain disease has the following confusion matrix:

	No disease	Has disease
Test negative	100	20
Test positive	50	180

A **true positive** is where a patient has the disease and the test is positive.

- (a) Calculate the sensitivity of the test. Provide an exact answer or 3 significant figures of precision. **(Make sure you show your working.)**
- (b) Calculate the specificity of the test. Provide an exact answer or 3 significant figures of precision. **(Make sure you show your working.)**
- (c) Calculate the accuracy of the test. Provide an exact answer or 3 significant figures of precision. **(Make sure you show your working.)**
- (d) Based on your results from parts (a)–(c), would you estimate that the AUC (area under the curve) for the ROC curve is closest to:
- between 0.65 and 0.90
 - between 0.45 and 0.55
 - between 0.1 and 0.35
 - between -0.35 and -0.1
 - between -0.9 and -0.65

Justify your reasoning.

- (e) What does an AUC value close to 0 tell you about your prediction model? If this was the case, how could we improve the model?

Multiple Choice Questions

Use the multiple choice answer sheet to record your answers.




Figure 5: Plot of city fuel efficiency (`cty`) against highway fuel efficiency (`hwy`) for Multiple Choice Questions 1 and 2.

1. From the data in Figure 5 what is the relationship between city and highway fuel efficiency?
 - (a) A strong positive relationship.
 - (b) A weak negative relationship.
 - (c) No relationship.
 - (d) Impossible to tell from this plot.

2. What R code could have generated the plot in Figure 5? (Select all the correct answers, and only the correct answers.)
 - (a) `mpg %>% ggplot(aes(x=cty, y=hwy))+geom_point()+geom_smooth(method="lm")`
 - (b) `mpg %>% ggplot(aes(x=cty, y=hwy))+geom_point()+geom_smooth()`
 - (c) `ggplot(mpg, aes(x=cty, y=hwy))+geom_smooth()`
 - (d) `ggplot(mpg, aes(x=cty, y=hwy))+geom_point()+geom_smooth()`




Figure 6: RStudio screenshot for Multiple Choice Questions 3–5.

3. Using the RStudio screenshot in Figure 6, what is the name of the Project?
 - (a) `table4a`
 - (b) `Console`
 - (c) `DT_examcode_2024_T1`
 - (d) `Exams`

4. Look at the RStudio screenshot in Figure 6. If we name the tidy tibble `table4a_tidy`, which command will convert it back into wide form?
 - (a) `gather(table4a_tidy, key = "year", value = "TB cases", '1999':'2000')`
 - (b) `inspect_df(table4a_tidy)`
 - (c) `wide_form(table4a_tidy, year, 'TB cases')`
 - (d) `spread(table4a_tidy, key = year, value = 'TB cases')`

5. We would like to obtain summary statistics for the numerical variables in `table4a`. Based on the information in the RStudio screenshot in Figure 6, will the command `inspect_num(table4a)` work for this purpose?
- (a) Yes, `inspect_num` is the correct command.
 - (b) No, `inspect_num` does not calculate summary statistics.
 - (c) No, we must first run the command `library(inspectdf)`.
 - (d) No, we must first run the two commands `install.packages("inspectdf")` and `library(inspectdf)`.
6. Using the RStudio screenshot in Figure 6, what command will return a 1×1 tibble containing just the value 745? (Select all the correct answers, and only the correct answers.)
- (a) `table4a %>% filter(country=="Afghanistan") %>% select("1999")`
 - (b) `filter(table4a, country=="Afghanistan")`
 - (c) `table4a %>% select("country"=="Afghanistan") %>% filter("1999")`
 - (d) `table4a[1,2]`
7. If we run the command `test_string <- "abc (15) xyz"`, what is the command to extract just the number 15? (Select all the correct answers, and only the correct answers.)
- (a) `str_match(test_string, "abc \\((\\d*)")[,2]`
 - (b) `str_match(test_string, "abc \\((\\d)")[,2]`
 - (c) `str_match(test_string, "abc (\\((\\d*)")[,2]`
 - (d) `str_match(test_string, "abc \\((.*)\\)")[,2]`




Figure 7: RStudio screenshot for Multiple Choice Questions 8–9.

8. We have hit an error when knitting the R Markdown file in Figure 7. We definitely have the `tidyverse` package installed, so what problems will prevent our file from knitting successfully?
 - (a) We have not specified the correct directory for the `csv` file and we need to click “Knit on Save”.
 - (b) `library(tidyverse)` is not inside a code chunk, and we have not specified the correct directory for the `csv` file.
 - (c) We should have `warning = FALSE` and we need to click “Knit on Save”.
 - (d) `library(tidyverse)` is not inside a code chunk and we should have `warning = FALSE`.

9. Once we get the R Markdown file in Figure 7 to knit, we find that the output is being displayed in the PDF. However, the code `newdata <- read_csv("StateData.csv")` runs successfully, but produces many warnings that we don't want to see. What changes could we make to prevent R Markdown from showing any warnings in the whole document?
- (a) Include the command `print(!warnings)` after line 16
 - (b) Include the command `print(read_csv == FALSE)` after line 9
 - (c) Change `warning = TRUE` to `warning = FALSE` on line 9
 - (d) Change `echo = FALSE` to `echo = FALSE, warning = FALSE` on line 14

See the attached page “mpg2 data set” for Questions 10 to 16.

For the remaining questions we are going to be working with the mpg2 data set, which is a simplified version of mpg.




Figure 8: Decision tree diagram for Multiple Choice Questions 10–13.

10. In Figure 8 we have fit the model `class ~ year + cyl + drive_type + displ` to a training subset of the mpg2 dataset for predicting the type of car, either small or large. What type of model have we fit?
- (a) Logistic tree
 - (b) General linear model tree
 - (c) Regression tree
 - (d) Classification tree

11. From the diagram in Figure 8, determine how decisions the model made.
- (a) 5
 - (b) 6
 - (c) 11
 - (d) 21
12. From the diagram in Figure 8, what proportion of the whole dataset was chosen for our training set?
- (a) 58%
 - (b) 105%
 - (c) 70%
 - (d) 15%
13. Using the model in Figure 8, what size do we predict for a car with an engine size of 3L, highway fuel efficiency of 23 miles per gallon, four wheel drive and built in 2008?
- (a) small
 - (b) large
 - (c) It can't be determined from this diagram
 - (d) The model was inconclusive




Figure 9: Boxplot of highway fuel efficiency [hwy](#) of a car against the number of cylinders in the car's engine [cyl](#) for Multiple Choice Questions 14–16.

14. According to the box plot in Figure 9, which number of cylinders has the smallest median highway fuel efficiency?
- (a) 4
 - (b) 5
 - (c) 6
 - (d) 8

15. According to the box plot in Figure 9, which number of cylinders has the smallest interquartile range?

- (a) 4
- (b) 5
- (c) 6
- (d) 8

16. According to the box plot in Figure 9, which number of cylinders has the largest number of outliers?

- (a) 4
- (b) 5
- (c) 6
- (d) 8

mpg2 data set

```
> mpg2
```

```
# A tibble: 234 × 11
```

	manufacturer	model	displ	year	cyl	trans	drive_type	cty	hwy	fl	class
	<chr>	<chr>	<dbl>	<int>	<ord>	<chr>	<fct>	<dbl>	<dbl>	<chr>	<fct>
1	audi	a4	1.8	1999	4	auto(l5)	front	18	29	p	small
2	audi	a4	1.8	1999	4	manual(m5)	front	21	29	p	small
3	audi	a4	2	2008	4	manual(m6)	front	20	31	p	small
4	audi	a4	2	2008	4	auto(av)	front	21	30	p	small
5	audi	a4	2.8	1999	6	auto(l5)	front	16	26	p	small
6	audi	a4	2.8	1999	6	manual(m5)	front	18	26	p	small
7	audi	a4	3.1	2008	6	auto(av)	front	18	27	p	small
8	audi	a4 quattro	1.8	1999	4	manual(m5)	four	18	26	p	small
9	audi	a4 quattro	1.8	1999	4	auto(l5)	four	16	25	p	small
10	audi	a4 quattro	2	2008	4	manual(m6)	four	20	28	p	small

```
# i 224 more rows
```

```
> skim_without_charts(mpg2)
```

```
-- Data Summary -----
```

	Values
Name	mpg2
Number of rows	234
Number of columns	11

```
-----  
Column type frequency:
```

character	4
factor	3
numeric	4

```
-----  
Group variables      None
```

```
-- Variable type: character -----
```

	skim_variable	n_missing	complete_rate	min	max	empty	n_unique	whitespace
1	manufacturer	0		1	4	10	0	15
2	model	0		1	2	22	0	38
3	trans	0		1	8	10	0	10
4	fl	0		1	1	1	0	5

```
-- Variable type: factor -----
```

	skim_variable	n_missing	complete_rate	ordered	n_unique	top_counts
1	cyl	0		1 TRUE	4	4: 81, 6: 79, 8: 70, 5: 4
2	drive_type	0		1 FALSE	3	fro: 106, fou: 103, rea: 25
3	class	0		1 FALSE	2	lar: 147, sma: 87

```
-- Variable type: numeric -----
```

	skim_variable	n_missing	complete_rate	mean	sd	p0	p25	p50	p75	p100
1	displ	0		3.47	1.29	1.6	2.4	3.3	4.6	7
2	year	0		2004.	4.51	1999	1999	2004.	2008	2008
3	cty	0		16.9	4.26	9	14	17	19	35
4	hwy	0		23.4	5.95	12	18	24	27	44