

# DONDOOIL®

DONDOOIL® Oral Syrup 28 mil.

## COMPOSITION:

Each 2 mil. / 1 mil. contains only naturals of:

PALM FRUIT (*Elaeis Guineensis*), HONEY (*Apidea Corbiculata* and *NEEM* (*Azadirachta Indica*).

Lauric acid, methyl ester 12:0 R43.71

Oleic acid, methyl ester (z)- 9-(18:1) R11.86

Myristic acid, methyl ester 14:0 R20.19

Palmitic acid, methyl ester 16:0 R7.53

Caprylic acid, methyl ester 8:0 R R7.18

Capric acid, methyl eater 10:0 R5.36

Stearic acid, methyl ester 18:0 R1.69

Linoleic acid, methyl ester 18:2n-6 R1.15

COLOUR - Black

## UNIQUENESS:

DONDOOIL® liquid of high Protein MCTs / MCFAs to enhancing the immune system in the myeloid and lymphoid progenitor to amplify antigen presenting cells to mediate innate and adopt immune response offers the unique ability of preventing, treating, eradicating and maintaining tolerance to healthy self tissue.

Treats: .Autoimmune deficiency .Gastrointestinal .Respiratory .Infection

## INDICATIONS:

Even though DONDOOIL® is seen to boost the immune system, fight Ulceration, Infection, Reproduction, and Immune Deficiency Disorders because of its high level of MCTs and Protein effectively,

DONDOOIL® is indicated for Arthritis, Waist pain, Common cold and Immune System to boost the immune system from the Myeloid and Lymphoid progenitors to amplify Antigen Presenting Cells (APCs) to mediate Innate and Adopt immune cell mediated response

## DOSAGE:

Drink one (1) tea spoon (2 mil.) of DONDOOIL® morning and evening best within an interval of twelve (12) hours in-between. Best taken with empty stomach in the morning and evening fifteen to thirty minutes before meal.

Or as prescribed by physician.

## CONTRADICTIONS:

Make a person tasty to drink at least one cup of water every hour before midnight to avoid weakness of the body and dehydration as a result of cell amplification, cell division and cell differentiation.

## WARNING AND PRECAUTIONS:

Care is advised in the administration of DONDOOIL® to patients with severe anaemia, heavy menstrual flow and other cases of loss of red blood cells, in this case, you are required to combine DONDOOIL® with blood nutrition or blood IV or you can consult a physician.

## SYMPTOMS OF OVERDOSE AND PARTICULARS OF ITS TREATMENT

Initially taking orally up to two table spoons of DONDOOIL® can make a patient dizzy, when this happens the patient should be given water and made to rest.

## SAFETY:

DONDOOIL® is well tolerated and can be safely prescribed to all gender

## STORAGE:

DONDOOIL® should be stored below 30°C, It is better to use the product within two years of it's manufacturing date.

## PACKAGING / PRESENTATION:

While selecting a bottle of DONDOOIL®, make sure you look out for the packaging date to be sure is valid. Opt for DONDOOIL® Registered Trade Mark with approved label and appropriate packaging from DONDO ORIGIN LTD for safety and quality control. Though DONDOOIL® condenses at low temperatures, it can be brought back to liquid form by placing the container in warm area.

## KEEP OUT OF REACH OF CHILDREN.

Carefully read the accompanying instruction before use

Manufactured by DONDO ORIGIN LTD, +2348074747449, +2348033367384