

Variational Methods for Discrete Surface Parameterization. Applications and Implementation.

vorgelegt von
Dipl.-Math. techn. Stefan Sechelmann

von der Fakultät II - Mathematik und Naturwissenschaften
der Technischen Universität Berlin
zur Erlangung des akademischen Grades

Doktor der Naturwissenschaften
– Dr. rer. nat. –

vorgelegte Dissertation

Promotionsausschuss

Vorsitzender: NN
Gutachter/Berichter: Prof. Dr. Alexander I. Bobenko
Prof. Dr. Max Wardetzky

Tag der wissenschaftlichen Aussprache: NN

Berlin, den 12. August 2013

D 83

Contents

Introduction	1
I Uniformization of discrete Riemann surfaces	3
1 Variational Description	5
1.1 Discrete Riemann surfaces	5
1.2 Uniformization	7
1.3 Variational principles for discrete metrics in \mathbb{E}^2 , \mathbb{H}^2 , and \mathbb{S}^2	8
1.4 Quotient spaces and fundamental domains	8
1.4.1 The cut-graph and fuchsian groups	9
1.4.2 Minimal presentation	9
1.4.3 Separated handles	9
1.4.4 Opposite sides identified	9
1.4.5 Canonical Keen Polygons	9
1.5 Uniformization of embedded genus $g = 0$, $g = 1$, and $g > 1$ surfaces	9
2 Examples	11
2.1 Branched coverings of $\hat{\mathbb{C}}$	11
2.1.1 Elliptic curves	11
2.1.2 The moduli space	11
2.1.3 Numerical convergence analysis	11
2.1.4 The modulus of the Wente torus	11
2.1.5 Construction of hyperelliptic surfaces	11
2.1.6 Weierstrass points on hyperelliptic surfaces	12
2.1.7 Canonical domains	12
2.1.8 Lawsons surface	12
2.2 From Schottky to Fuchsian Uniformization	12

2.2.1	Images of isometric circles	13
2.2.2	Hyperelliptic data	13
2.3	Conformal maps to $\hat{\mathbb{C}}$	13
2.3.1	Selection of Branch Data	13
2.3.2	Examples	13
2.4	Surfaces with boundary	13
2.4.1	Variation of edge length	13
2.4.2	Examples	13
2.5	Conformal maps of planar domains	13
2.5.1	Boundary conditions	13
2.5.2	Comparison with examples of the Schwarz-Christoffel community	13
II	Variational Methods for Discrete Surface Parameterization	15
III	Software Packages	17
	Bibliography	19
	Acknowledgements	21

List of Figures

1.1	Discrete surfaces from glued triangles	5
1.2	Euclidean conformal equivalence	6
1.3	Conformal equivalence of Euclidean and hyperbolic/spherical metrics	7
1.4	Angles at a vertex	8
1.5	Hyperbolic flat metric on a genus 2 surface and the axes of the associated hyperbolic motions.	9
2.1	Schottky group generating a Riemann surface of genus 2. The point at infinity is not contained in any of the circles. The parameters A and B lie inside the circles by definition.	12
2.2	Fuchsian uniformization of a Riemann surface (right) given by Schottky data (left).	13
2.3	Square with symmetric slit to the circle	14

Introduction

Part I

Uniformization of discrete Riemann surfaces

Chapter 1

Variational Description

The theory of discrete uniformization presented here is based on the notion of discrete conformal equivalence of triangle meshes. The Euclidean definition was first considered by [6], the variational principle and applications in computer graphics is due to [8, 7, 2]. The notion of conformal equivalence of non-Euclidean metrics and corresponding variational principles were first defined in [2]. [4] investigate the gradient flow of this principle. Most of the material presented here can be found in [3].

1.1 Discrete Riemann surfaces

Definition 1. A discrete surface is a collection of triangles equipped with a metric of constant Gaussian curvature and geodesic edges. Triangles are glued along edges to form a surface.

By glueing triangles equipped with a metric of constant Gaussian curvature we obtain a surface that has constant curvature everywhere except for points where the metric has cone-like singularities (Figure 1.1). A discrete surface is called Euclidean for $K = 0$, hyperbolic for $K < 0$, and spherical if $K > 0$. Remark: In the latter we will use Gaussian curvature and curvature synonymously. Generically a discrete surface can have boundary components where triangles have not been glued. We consider this case in Section 2.5.

A discrete surface consists of vertices, edges, and faces $S = (V, E, F)$. We use single indices for denoting vertices, e.g., $i \in V$, edges are denoted $ij \in E$, and faces $ijk \in F$.




Figure 1.1: Discrete surfaces constructed from glued triangles of constant curvature. Euclidean, hyperbolic, and spherical. Bold edges are identified to create a cone-like singularity at the vertex.




Figure 1.2: Two Euclidean triangles are discretely conformally equivalent if their edge lengths agree after scaling by logarithmic factors u defined on vertices.

Definition 2. The map $l : E \rightarrow \mathbb{R}$ of triangle edge lengths of a discrete surface is called a discrete Euclidean, hyperbolic, or spherical metric respectively.

As in the smooth theory we define what it means for a metric to be conformally equivalent to another metric.

Definition 3. A discrete Euclidean metric with edge lengths l is discretely conformally equivalent to the discrete Euclidean metric \tilde{l} if there is a function $u : V \rightarrow \mathbb{R}$ such that for all edges $ij \in E$ it is

$$l_{ij} = e^{\frac{1}{2}(u_i+u_j)} \tilde{l}_{ij} \quad (1.1)$$

This definition is motivated by the smooth theory of Riemann surfaces where two metrics g and \tilde{g} on a 2-manifold M are conformally equivalent if there is a smooth function $u : M \rightarrow \mathbb{R}$ with

$$g = e^{2u} \tilde{g}.$$

Every discrete Euclidean metric is discretely conformally equivalent to a corresponding discrete hyperbolic or discrete spherical metric by the following

Definition 4. A discrete Euclidean metric l and a discrete hyperbolic or discrete spherical metric \tilde{l} are discretely conformally equivalent if for all edges $ij \in E$

$$l_{ij} = 2 \sinh \frac{\tilde{l}_{ij}}{2} \quad (1.2)$$

$$l_{ij} = 2 \sin \frac{\tilde{l}_{ij}}{2} \quad (1.3)$$

for \tilde{l} hyperbolic or spherical respectively (see Figure 1.3).

Literally this means that in the spherical case if a triangle is fit onto the sphere the conformally equivalent spherical lengths are the lengths of spherical geodesics connecting the triangle vertices. The same intuition holds on the upper sheet of the two-sheeted unit hyperboloid for hyperbolic triangles.

Combining Equations 1.1 and 1.2/1.3 we can define conformal equivalence of Euclidean and hyperbolic/spherical triangulations via transitivity of equivalence.

With this general notion of conformal equivalence of Euclidean, hyperbolic, and spherical metrics we can now define discrete Riemann surfaces.




Figure 1.3: Relations between hyperbolic/spherical lengths \tilde{l} and corresponding Euclidean edge lengths l . See Definition 4.

Definition 5. A discrete Riemann surface is an equivalence class of discretely conformally equivalent metrics.

If one restricts the equivalence class to either Euclidean, hyperbolic, or spherical metrics the length cross-ratio lcr_{ij} at edge ij with opposite vertices k and m is a conformal invariant. Note that this definition depends on the orientation of the surface.

$$\text{lcr}_{ij} = \frac{l_{ik}l_{jm}}{l_{mi}l_{kj}} \quad (1.4)$$

Length cross-ratios are also invariant under Möbius transformations of the ambient space.

1.2 Uniformization

We can now state the uniformization problem: *Given a discrete Riemann surface, find a metric of constant curvature without cone singularities.*

This means that the angles α_{jk}^i of Euclidean, hyperbolic, or spherical triangles around each vertex $i \in V$ sum up to 2π (Figure 1.4). A discrete surface with a cone-like singularity free metric has constant curvature everywhere.

As in the smooth case we expect to find a discrete metric with zero Gaussian curvature for tori, a constant negative curvature metric for surfaces with genus $g > 1$, and a metric with positive constant curvature for spheres. In the latter we will normalize the curvature of the target spaces to have constant Gaussian curvature 0, -1 , or 1.

In this work we calculate discrete uniformizations as minimizers of functionals that fit the target geometry. The variational description of the uniformization problem then amounts to finding a critical point of the functional $E(u)$ where

$$\frac{\partial E}{\partial u_i} = 2\pi - \sum_{ijk \ni i} \alpha_{jk}^i. \quad (1.5)$$




Figure 1.4: Angles at a vertex

1.3 Variational principles for discrete metrics in \mathbb{E}^2 , \mathbb{H}^2 , and \mathbb{S}^2

Construction of discrete flat metrics. A discrete Euclidean flat metric is the minimizer of a convex functional.

$$\lambda_{ij} := 2 \log l_{ij} \quad (1.6)$$

$$\tilde{\lambda}_{ij} := \lambda_{ij} + u_i + u_j \quad (1.7)$$

$$f_{Euc}(u_i, u_j, u_k) := \alpha_i \tilde{\lambda}_{jk} + \alpha_j \tilde{\lambda}_{ki} + \alpha_k \tilde{\lambda}_{ij} + 2(\mathbb{I}(\alpha_i) + \mathbb{I}(\alpha_j) + \mathbb{I}(\alpha_k)) \quad (1.8)$$

Definition 6.

$$E_{Euc}(u) := \sum_{ijk \in F} \left(f_{Euc}(u_i, u_j, u_k) - \frac{\pi}{2} (\tilde{\lambda}_{jk} + \tilde{\lambda}_{ki} + \tilde{\lambda}_{ij}) \right) + \sum_{i \in V} \Theta_i u_i \quad (1.9)$$

This definition and the derivatives can be found in [2]

For the hyperbolic case λ and $\tilde{\lambda}$ are defined as before. Further define

$$\beta_i := \frac{1}{2} (\pi + \alpha_i - \alpha_j - \alpha_k) \quad (1.10)$$

$$\beta_j := \frac{1}{2} (\pi - \alpha_i + \alpha_j - \alpha_k) \quad (1.11)$$

$$\beta_k := \frac{1}{2} (\pi - \alpha_i - \alpha_j + \alpha_k) \quad (1.12)$$

$$f_{Hyp}(u_i, u_j, u_k) := \beta_i \tilde{\lambda}_{jk} + \beta_j \tilde{\lambda}_{ki} + \beta_k \tilde{\lambda}_{ij} \quad (1.13)$$

$$+ \mathbb{I}(\alpha_i) + \mathbb{I}(\alpha_j) + \mathbb{I}(\alpha_k) + \mathbb{I}(\beta_i) + \mathbb{I}(\beta_j) + \mathbb{I}(\beta_k) \quad (1.14)$$

$$+ \mathbb{I} \left(\frac{1}{2} (\pi - \alpha_i - \alpha_j - \alpha_k) \right) \quad (1.15)$$

Definition 7.

$$E_{Hyp}(u) := \sum_{ijk \in F} \left(f_{Hyp}(u_i, u_j, u_k) - \frac{\pi}{2} (\tilde{\lambda}_{jk} + \tilde{\lambda}_{ki} + \tilde{\lambda}_{ij}) \right) + \sum_{i \in V} \Theta_i u_i \quad (1.16)$$

1.4 Quotient spaces and fundamental domains

Every Riemann surface R has a universal cover X , i.e., a simply connected covering space and a corresponding covering map. A metric of constant curvature on a compact 2-manifold can be realized as the quotient of the universal cover over a uniformizing group.




Figure 1.5: Hyperbolic flat metric on a genus 2 surface and the axes of the associated hyperbolic motions.

Triangulated surfaces of genus $g \geq 2$ without boundary can be equipped with a discretely conformally equivalent flat hyperbolic metric [2]. By flat hyperbolic metric we mean that the edge length are hyperbolic and for any vertex the angle sum is 2π . To realize this metric in the hyperbolic plane e.g. in the Poincaré disk model one has to introduce cuts along a basis of the homotopy. This creates a simply connected domain in \mathbb{H}^2 . Matching cut paths are related by a hyperbolic motion i.e. the Möbius transformations that leave the unit disk invariant (Figure 1.5).

1.4.1 The cut-graph and fuchsian groups

Want so say here: the number of transformations generated by the mapping of corresponding edges equals the number of path segments in the homotopy-cut-graph. They generate a fuchsian group with #vertices relations

1.4.2 Minimal presentation

1.4.3 Separated handles

1.4.4 Opposite sides identified

1.4.5 Canonical Keen Polygons

1.5 Uniformization of embedded genus $g = 0$, $g = 1$, and $g > 1$ surfaces

Chapter 2

Examples

2.1 Branched coverings of $\hat{\mathbb{C}}$

In this section we discuss surfaces that arise as branched coverings of $\hat{\mathbb{C}}$. A Riemann surface that can be represented as a double cover of $\hat{\mathbb{C}}$ is called elliptic for $g = 1$ and hyperelliptic for $g > 1$ [5, p. 235].

Let $\lambda_1, \dots, \lambda_{2g+2} \in \hat{\mathbb{C}}, \lambda_i \neq \lambda_j \forall i \neq j$. The algebraic curve

$$C = \{(z, w) \in \mathbb{C}^2 \mid w^2 = \prod_{i=1}^{2g+2} (z - \lambda_i)\} \quad (2.1)$$

is a one dimensional complex manifold. A branched double cover of $\hat{\mathbb{C}}$ is the projection $\pi : C \rightarrow \mathbb{C}, C \ni (z, w) \mapsto z$. A discrete branched cover of $\hat{\mathbb{C}}$ is a triangulation of $\pi(C)$ with vertices at λ_i .

2.1.1 Elliptic curves

2.1.2 The moduli space

2.1.3 Numerical convergence analysis

2.1.4 The modulus of the Wente torus

2.1.5 Construction of hyperelliptic surfaces

Any hyperelliptic Riemann surface can be expressed as an algebraic curve of the form

$$w^2 = \prod_{i=1}^{2g+2} (z - \lambda_i) \quad g \geq 1, \quad \lambda_i \neq \lambda_j \forall i \neq j.$$

Here λ_i are the branch points of the doubly covered Riemann sphere.




Figure 2.1: Schottky group generating a Riemann surface of genus 2. The point at infinity is not contained in any of the circles. The parameters A and B lie inside the circles by definition.

2.1.6 Weierstrass points on hyperelliptic surfaces

A hyperelliptic surface comes together with a holomorphic involution h called the hyperelliptic involution. The branch points are fixed points under this transformation. For a hyperelliptic algebraic curve it is $h(\mu, \lambda) = (-\mu, \lambda)$

2.1.7 Canonical domains

2.1.8 Lawsons surface

2.2 From Schottky to Fuchsian Uniformization

Let $C_1, C'_1, \dots, C_g, C'_g$ be disjoint circles in $\hat{\mathbb{C}}$. A classical Schottky group G is a Kleinian group with generators $\sigma_1, \dots, \sigma_g$ satisfying

$$\frac{\sigma_i z - B_i}{\sigma_i z - A_i} = \mu_i \frac{z - B_i}{z - A_i}, \quad 0 < |\mu_i| < 1,$$

where σ_i maps the exterior of C_i onto the interior of C'_i . The points A_i and B_i lie inside the circles C_i and B_i respectively [1]. Figure 2.1 illustrates this construction. The quotient space $\hat{\mathbb{C}}/G$ is a Riemann surface of genus g . The group G is called the uniformizing Schottky group

Given a Riemann surface of genus g and a classical Schottky uniformization, we calculate a corresponding Fuchsian uniformization using discrete conformal equivalence of spherical and hyperbolic triangle meshes. A fundamental domain of the Schottky uniformization is the Riemann sphere without the interior of the circles.

The triangulation of a fundamental domain has to respect the mappings between the circles. Vertices on identified circles have to be identified as well. It is however not clear a priori what the conformal structure of a triangulated fundamental domain is. The edge length on identified circles differ since there is a Möbius transformation between them.

The idea to define the conformal structure is to use length cross-ratios instead of edge lengths. Both definitions are equivalent and define the conformal structure of a discrete Riemann surface. So given length cross ratios on edges we can calculate representative edge lengths in the discrete




Figure 2.2: Fuchsian uniformization of a Riemann surface (right) given by Schottky data (left).

conformal class of the surface and, more obvious, vice versa. On boundary edges of the fundamental domain the length cross ratios agree since length-cross ratios are Möbius invariant.

2.2.1 Images of isometric circles

2.2.2 Hyperelliptic data

2.3 Conformal maps to $\hat{\mathbb{C}}$

2.3.1 Selection of Branch Data

2.3.2 Examples

2.4 Surfaces with boundary

2.4.1 Variation of edge length

2.4.2 Examples

2.5 Conformal maps of planar domains

2.5.1 Boundary conditions

2.5.2 Comparison with examples of the Schwarz-Christoffel community




Figure 2.3: Square with symmetric slit to the circle

Part II

Variational Methods for Discrete Surface Parameterization

Part III

Software Packages

Bibliography

- [1] Alexander I. Bobenko, *Introduction to compact riemann surfaces*, Computational Approach to Riemann Surfaces, Lecture Notes in Mathematics, no. 2013, Springer, 2011, pp. 3–64.
- [2] Alexander I. Bobenko, Ulrich Pinkall, and Boris Springborn, *Discrete conformal maps and ideal hyperbolic polyhedra*, Preprint; <http://arxiv.org/abs/1005.2698>, 2010.
- [3] Alexander I. Bobenko, Stefan Sechelmann, and Boris Springborn, *Uniformization of discrete Riemann surfaces*, in preparation.
- [4] Ren Guo, *Combinatorial yamabe flow on hyperbolic surfaces with boundary*, Commun. Contemp. Math. **13** (2011), no. 5, 827–842.
- [5] Jürgen Jost, *Compact riemann surfaces*, Universitext (En ligne), Springer-Verlag Berlin Heidelberg, 2007.
- [6] Feng Luo, *Combinatorial yamabe flow on surfaces.*, Commun. Contemp. Math. **6** (2004), no. 5, 765–780 (English).
- [7] Boris Springborn, *Discrete conformal equivalence for triangle meshes*, Oberwolfach Reports, Oberwolfach Reports, vol. 6, 2009, pp. 104–106.
- [8] Boris Springborn, Peter Schröder, and Ulrich Pinkall, *Conformal equivalence of triangle meshes*, ACM Trans. Graph. **27** (2008), no. 3, 77:1–77:11.

Acknowledgements