

Uniformization of discrete Riemann surfaces and Applications

vorgelegt von
Dipl.-Math. techn. Stefan Sechelmann

von der Fakultät II - Mathematik und Naturwissenschaften
der Technischen Universität Berlin
zur erlangung des akademischen Grades

Doktor der Naturwissenschaften
– Dr. rer. nat. –

Promotionsausschuss

Vorsitzender: NN
Gutachter/Berichter: Prof. Dr. Alexander I. Bobenko
NN

Tag der wissenschaftlichen Aussprache: NN

Berlin, den 07. Januar 2013

Contents

1	Introduction	1
2	Discrete Uniformization	3
2.1	Discrete Riemann surfaces	3
2.2	Variational principles	5
2.2.1	Discrete conformal equivalence	5
2.2.2	Variational principles for discrete metrics in \mathbb{E}^2 , \mathbb{H}^2 , and \mathbb{S}^2	5
2.2.3	Realization	6
2.3	Uniformization of surfaces of higher genus	6
2.3.1	The cut-graph and fuchsian groups	6
2.3.2	Minimal presentation	7
2.4	Canonical fundamental domains of fuchsian groups	7
2.4.1	Separated handles	7
2.4.2	Opposite sides identified	7
2.5	Uniformization of elliptic and hyperelliptic surfaces	7
2.5.1	Elliptic Functions	7
2.5.2	The moduli space	7
2.5.3	Numerical convergence analysis	7
2.5.4	The modulus of the Wente torus	7
2.5.5	Construction of hyperelliptic surfaces	7
2.5.6	Weierstrass points on hyperelliptic surfaces	7
2.5.7	Canonical domains	8
2.5.8	Lawsons surface	8
2.6	Meromorphic functions (planned)	8
2.6.1	Conformal maps to $\hat{\mathbb{C}}$ with singularities	8
2.6.2	Selection of Branch Data	8
2.6.3	Examples	8

2.7	Simply and multiply connected domains	8
2.7.1	Variation of edge length	8
2.7.2	Examples	8
2.8	Conformal maps of planar domains	8
2.8.1	Boundary conditions	8
2.8.2	Comparison with examples of the Schwarz-Christoffel community	8
3	Discrete Surface Parameterization	9
3.1	Discrete quasiisothermic parametrizations	9
3.1.1	Discrete quasiisothermic parameterizations	10
3.1.2	Formulation as boundary value problem	10
3.1.3	Global approach	10
3.1.4	Variational principle for S-isothermic surfaces	10
3.1.5	Constructing the associated family of appproximate minimal surfaces	10
3.1.6	A discrete ellipsoid and its dual surface	10
3.1.7	Applications in architecture	10
3.2	Piece-wise projective interpolation for arbitrary parameterizations	10
3.2.1	Approximation by conformal deformation	10
3.2.2	Examples	10
3.3	A variational principle for discrete Tschebyshev nets	10
3.3.1	Tschebyscheff Meshes	10
3.3.2	Variational Principle	10
3.3.3	Applications in architecture	10
	Bibliography	11

List of Figures

2.1	Discrete surfaces constructed from glued triangles of constant curvature. Euclidean, hyperbolic, and spherical. Bold edges are identified to create a cone-like singularity at the vertex.	3
2.2	Two Euclidean triangles are discretely conformally equivalent if their edge lengths can be scaled by logarithmic factors u defined on vertices.	4
2.3	Hyperbolic flat metric on a genus 2 surface and the axes of the associated hyperbolic motions.	6
2.4	Square with symmetric slit to the circle	8

Chapter 1

Introduction

Chapter 2

Discrete Uniformization

The discrete uniformization theory presented here is based on the notion of discrete conformal equivalence of triangle meshes. The Euclidean definition was first considered by [Luo04], the variational principle and applications in computer graphics is due to [SSP08, Spr09, BPS10]. The notion of conformal equivalence of non-Euclidean metrics and corresponding variational principles were first defined in [BPS10]. [Guo11] investigate the gradient flow of this principle. Most of the material presented here can be found in [BSS].

2.1 Discrete Riemann surfaces

Definition 1. A discrete surface is a collection of triangles equipped with a metric of constant Gaussian curvature and geodesic edges. Triangles are glued along edges to form a surface.

By glueing triangles equipped with a metric of constant Gaussian curvature we obtain a surface that has constant curvature everywhere except for points where the metric has cone-like singularities (Figure 2.1). A discrete surface is called Euclidean for $K = 0$, hyperbolic for $K = -1$, and spherical if $K = 1$. Remark: In the latter we will use Gaussian curvature and curvature synonymously. Generically a discrete surface can have boundary components where triangles have not been glued. We consider this case in Section 2.7.

A discrete surface consists of vertices, edges, and faces $S = (V, E, F)$. We use single indices for denoting vertices, e.g., $i \in V$. Edges are denoted by $ij \in E$ and faces $ijk \in F$.




Figure 2.1: Discrete surfaces constructed from glued triangles of constant curvature. Euclidean, hyperbolic, and spherical. Bold edges are identified to create a cone-like singularity at the vertex.




Figure 2.2: Two Euclidean triangles are discretely conformally equivalent if their edge lengths can be scaled by logarithmic factors u defined on vertices.

Definition 2. *The map $l : E \rightarrow \mathbb{R}$ of triangle edge lengths of a discrete surface is called a discrete Euclidean, hyperbolic, or spherical metric respectively.*

As in the smooth theory we define what it means for a metric to be conformally equivalent to another metric.

Definition 3. *A discrete Euclidean metric with edge lengths l is discretely conformally equivalent to the discrete Euclidean metric \tilde{l} if there is a function $u : V \rightarrow \mathbb{R}$ such that for all edges $ij \in E$ it is*

$$l_{ij} = e^{\frac{1}{2}(u_i+u_j)} \tilde{l}_{ij} \quad (2.1)$$

This definition is motivated by the smooth theory of Riemann surfaces where two metrics g and \tilde{g} on a 2-manifold M are conformally equivalent if there is a smooth function $u : M \rightarrow \mathbb{R}$ with

$$g = e^{2u} \tilde{g}.$$

Every Euclidean metric is discretely conformally equivalent to a corresponding hyperbolic or spherical metric by the following 2.3

Definition 4. *A discrete Euclidean metric l and a discrete hyperbolic or discrete spherical metric \tilde{l} are discretely conformally equivalent if for all edges $ij \in E$*

$$l_{ij} = 2 \sinh \frac{\tilde{l}_{ij}}{2} \quad (2.2)$$

$$l_{ij} = 2 \sin \frac{\tilde{l}_{ij}}{2} \quad (2.3)$$

for \tilde{l} hyperbolic or spherical respectively.

Literally this means that in the spherical case each triangle can be fit onto the sphere. The conformally equivalent spherical lengths are the lengths of spherical geodesics connecting the triangle vertices.

2.2 Variational principles

2.2.1 Discrete conformal equivalence

Definition 5. Two Euclidean triangulations T and \tilde{T} are discretely conformally equivalent if there is a map $u : V \rightarrow \mathbb{R}$ such that for any edge ij it is

$$l_{ij} = e^{u_i + u_j} \tilde{l}_{ij}$$

where l_{ij} is the length of the edge ij .

Definition 6. A discrete flat Euclidean metric is a map $l : E \rightarrow \mathbb{R}_+$ such that triangle inequalities are satisfied and angle sums around each inner vertex are equal to 2π .

2.2.2 Variational principles for discrete metrics in \mathbb{E}^2 , \mathbb{H}^2 , and \mathbb{S}^2

Construction of discrete flat metrics. A discrete Euclidean flat metric is the minimizer of a convex functional.

$$\lambda_{ij} := 2 \log l_{ij} \quad (2.4)$$

$$\tilde{\lambda}_{ij} := \lambda_{ij} + u_i + u_j \quad (2.5)$$

$$f_{Euc}(u_i, u_j, u_k) := \alpha_i \tilde{\lambda}_{jk} + \alpha_j \tilde{\lambda}_{ki} + \alpha_k \tilde{\lambda}_{ij} + 2(\mathcal{I}(\alpha_i) + \mathcal{I}(\alpha_j) + \mathcal{I}(\alpha_k)) \quad (2.6)$$

Definition 7.

$$E_{Euc}(u) := \sum_{ijk \in F} \left(f_{Euc}(u_i, u_j, u_k) - \frac{\pi}{2} (\tilde{\lambda}_{jk} + \tilde{\lambda}_{ki} + \tilde{\lambda}_{ij}) \right) + \sum_{i \in V} \Theta_i u_i \quad (2.7)$$

This definition and the derivatives can be found in [BPS10]

For the hyperbolic case λ and $\tilde{\lambda}$ are defined as before. Further define

$$\beta_i := \frac{1}{2} (\pi + \alpha_i - \alpha_j - \alpha_k) \quad (2.8)$$

$$\beta_j := \frac{1}{2} (\pi - \alpha_i + \alpha_j - \alpha_k) \quad (2.9)$$

$$\beta_k := \frac{1}{2} (\pi - \alpha_i - \alpha_j + \alpha_k) \quad (2.10)$$

$$f_{Hyp}(u_i, u_j, u_k) := \beta_i \tilde{\lambda}_{jk} + \beta_j \tilde{\lambda}_{ki} + \beta_k \tilde{\lambda}_{ij} \quad (2.11)$$

$$+ \mathcal{I}(\alpha_i) + \mathcal{I}(\alpha_j) + \mathcal{I}(\alpha_k) + \mathcal{I}(\beta_i) + \mathcal{I}(\beta_j) + \mathcal{I}(\beta_k) \quad (2.12)$$

$$+ \mathcal{I} \left(\frac{1}{2} (\pi - \alpha_i - \alpha_j - \alpha_k) \right) \quad (2.13)$$

Definition 8.

$$E_{Hyp}(u) := \sum_{ijk \in F} \left(f_{Hyp}(u_i, u_j, u_k) - \frac{\pi}{2} (\tilde{\lambda}_{jk} + \tilde{\lambda}_{ki} + \tilde{\lambda}_{ij}) \right) + \sum_{i \in V} \Theta_i u_i \quad (2.14)$$




Figure 2.3: Hyperbolic flat metric on a genus 2 surface and the axes of the associated hyperbolic motions.

2.2.3 Realization

2.3 Uniformization of surfaces of higher genus

Triangulated surfaces of genus $g \geq 2$ without boundary can be equipped with a discretely conformally equivalent flat hyperbolic metric [BPS10]. By flat hyperbolic metric we mean that the edge lengths are hyperbolic and for any vertex the angle sum is 2π . To realize this metric in the hyperbolic plane e.g. in the Poincaré disk model one has to introduce cuts along a basis of the homotopy. This creates a simply connected domain in \mathbb{H}^2 . Matching cut paths are related by a hyperbolic motion i.e. the Möbius transformations that leave the unit disk invariant (Figure 2.3).

2.3.1 The cut-graph and fuchsian groups

Want so say here: the number of transformations generated by the mapping of corresponding edges equals the number of path segments in the homotopy-cut-graph. They generate a fuchsian group with #vertices relations

Proposition 1.

2.3.2 Minimal presentation**2.4 Canonical fundamental domains of fuchsian groups****2.4.1 Separated handles****2.4.2 Opposite sides identified****2.5 Uniformization of elliptic and hyperelliptic surfaces****2.5.1 Elliptic Functions****2.5.2 The moduli space****2.5.3 Numerical convergence analysis****2.5.4 The modulus of the Wente torus****2.5.5 Construction of hyperelliptic surfaces**

Any hyperelliptic Riemann surface can be expressed as an algebraic curve of the form

$$w^2 = \prod_{i=1}^{2g+2} (z - \lambda_i) \quad g \geq 1, \quad \lambda_i \neq \lambda_j \forall i \neq j.$$

Here λ_i are the branch points of the doubly covered Riemann sphere.

2.5.6 Weierstrass points on hyperelliptic surfaces

A hyperelliptic surface comes together with a holomorphic involution h called the hyperelliptic involution. The branch points are fixed points under this transformation. For a hyperelliptic algebraic curve it is $h(\mu, \lambda) = (-\mu, \lambda)$




Figure 2.4: Square with symmetric slit to the circle

2.5.7 Canonical domains

2.5.8 Lawsons surface

2.6 Meromorphic functions (planned)

2.6.1 Conformal maps to $\hat{\mathbb{C}}$ with singularities

2.6.2 Selection of Branch Data

2.6.3 Examples

2.7 Simply and multiply connected domains

2.7.1 Variation of edge length

2.7.2 Examples

2.8 Conformal maps of planar domains

2.8.1 Boundary conditions

2.8.2 Comparison with examples of the Schwarz-Christoffel community

Chapter 3

Discrete Surface Parameterization

3.1 Discrete quasiisothermic parametrizations

The notion of quasiconformal parameterizations

- 3.1.1 Discrete quasiisothermic parameterizations
- 3.1.2 Formulation as boundary value problem
- 3.1.3 Global approach
- 3.1.4 Variational principle for S-isothermic surfaces
- 3.1.5 Constructing the associated family of approximate minimal surfaces
- 3.1.6 A discrete ellipsoid and its dual surface
- 3.1.7 Applications in architecture
- 3.2 Piece-wise projective interpolation for arbitrary parameterizations
 - 3.2.1 Approximation by conformal deformation
 - 3.2.2 Examples
- 3.3 A variational principle for discrete Tschebyshev nets
 - 3.3.1 Tschebyscheff Meshes
 - 3.3.2 Variational Principle
 - 3.3.3 Applications in architecture

Bibliography

- [BPS10] Alexander I. Bobenko, Ulrich Pinkall, and Boris Springborn. Discrete conformal maps and ideal hyperbolic polyhedra. Preprint; <http://arxiv.org/abs/1005.2698>, 2010.
- [BSS] Alexander I. Bobenko, Stefan Sechelmann, and Boris Springborn. Uniformization of discrete Riemann surfaces. in preparation.
- [Guo11] Ren Guo. Combinatorial yamabe flow on hyperbolic surfaces with boundary. *Commun. Contemp. Math.*, 13(5):827–842, 2011.
- [Luo04] Feng Luo. Combinatorial yamabe flow on surfaces. *Commun. Contemp. Math.*, 6(5):765–780, 2004.
- [Spr09] Boris Springborn. Discrete conformal equivalence for triangle meshes. In *Oberwolfach Reports*, volume 6 of *Oberwolfach Reports*, pages 104–106, 2009.
- [SSP08] Boris Springborn, Peter Schröder, and Ulrich Pinkall. Conformal equivalence of triangle meshes. *ACM Trans. Graph.*, 27(3):#77, 1–11, 2008.