

Screen Shot Taken on 11/4/16 of Order set for Post-Partum Hemorrhage

Order Sets		Actions	Orders
Summary	<p>1,000 mg, Intravenous, EVERY 12 HOURS</p> <p><b>MISOPROSTOL FOR CERVICAL RIPENING</b></p> <p><input checked="" type="checkbox"/> misoprostol (CYTOTEK) tablet 25 mcg, Vaginal, ONCE</p> <p><input checked="" type="checkbox"/> misoprostol (CYTOTEK) tablet 50 mcg, Oral, ONCE</p> <p><b>LAMINARIA FOR CERVICAL RIPENING</b></p> <p><input type="checkbox"/> cervical dilator (LAMINARIA) 3 MM device</p> <p><input type="checkbox"/> cervical dilator (LAMINARIA) 5 MM device</p> <p><b>OXYTOCIN FOR POST DELIVERY</b></p> <p><input type="checkbox"/> oxytocin (PITOCIN) 20 units in 1,000 mL Intravenous, PER ADMINISTRATION INSTRUCTIONS</p> <p><input type="checkbox"/> PHYSICIAN TO NURSE - 14 french triple lumen foley catheter with 30cc balloon inserted into cervix by provider for manual cervical dilation. Routine, ONGOING, Starting 11/4/16 for 1 occurrence, 14 french triple lumen foley catheter with 30cc balloon inserted into cervix by provider for manual cervical dilation. When inserted gentle traction hourly or as needed.</p> <p><input type="checkbox"/> PHYSICIAN TO NURSE - Foley bulb inserted into cervix by physician for induction Routine, ONCE, Starting 11/4/16 for 1 occurrence</p> <p><b>OBSTETRICAL HEMORRHAGE RISK ASSESSMENT — Required</b></p> <p><b>Low Risk</b> No risk factors identified</p> <p><b>High Risk</b> Placenta previa, low lying placenta, suspected placenta accreta, increta or percreta, hematocrit less than 30 AND other risk factors, platelets less than or equal to 100,000, active bleeding (greater than normal show) on admit, known coagulopathy with abnormal coag results, greater than 2 previous c-section, greater than 4 vaginal births, multiple gestation greater than 2 fetus, history of previous PPH, positive antibody screen</p> <p><input checked="" type="radio"/> LOW RISK - GP/IR</p> <p><input checked="" type="radio"/> HIGH RISK - GP/IR</p>		<p>Manage Orders Go to Order Sets Options</p> <p>Providers</p> <p>Place new order New</p> <p>Select order mode Next</p> <p>Orders from Order Sets</p> <p><b>IP OB ADMISSION ORDERS</b> Remove</p> <p><b>PNEUMATIC COMPRESSION DEVICE</b> X</p> <p>Routine, CONTINUOUS starting Today at 1100 Unit Specified Placement: Bilateral</p> <p>The provider selected is not authorized to be an ORDERING provider for this order mode. Please enter a physician or mid-level provider in the ORDERING provider field.</p> <p><b>lidocaine (XYLOCAINE) 1 % injection 0.1 mL</b> X</p> <p>0.1 mL, Intradermal, AS NEEDED, FOR IV START, starting Today at 1059, For 1 dose, Routine</p> <p>The provider selected is not authorized to be an ORDERING provider for this order mode. Please enter a physician or mid-level provider in the ORDERING provider field.</p> <p><b>BLOOD GASES CORD VENOUS</b> X</p> <p>LAB STAT ONCE First occurrence Today at 1100 Unit Collect</p> <p>The provider selected is not authorized to be an ORDERING provider for this order mode. Please enter a physician or mid-level provider in the ORDERING provider field.</p> <p><b>BLOOD GASES, CORD, ARTERIAL</b> X</p> <p>LAB STAT ONCE First occurrence Today at 1100 Unit Collect</p> <p>The provider selected is not authorized to be an ORDERING provider for this order mode. Please enter a physician or mid-level provider in the ORDERING provider field.</p> <p><b>COMPLETE BLOOD COUNT</b> X</p> <p>LAB STAT ONCE First occurrence Today at 1100 Unit Collect</p> <p>The provider selected is not authorized to be an ORDERING provider for this order mode. Please enter a physician or mid-level provider in the ORDERING provider field.</p>