

January 9, 2018

## 2 Lecture 2

### 2.1 Taylor series

- We know that  $\sin(30^\circ) = \frac{1}{2}$ . How do we compute the value of  $\sin(31^\circ)$ ?
- We expect that it will be close to  $\frac{1}{2}$ , because  $\sin(x)$  is continuous in  $x$ , but how do we compute an accurate answer?
- This is the sort of problem where Taylor series are useful.
- Basically, we know the value of a function at some point  $x = a$ , i.e. we know  $f(a)$ , and we wish to compute  $f$  for nearby values  $x = a + \varepsilon$ , where  $|\varepsilon|$  is assumed small. Hence we seek to compute the value of  $f(a + \varepsilon)$ .
- If  $a = 0$ , the series is also called a Maclaurin series.
- **(Taylor's theorem)**

If  $f(x)$  and its first  $(n + 1)$  derivatives  $f'(x), f''(x), \dots, f^{(n+1)}(x)$  all exist and are continuous and bounded in some interval  $x_0 \leq x \leq x_1$ , then if  $a \in [x_0, x_1]$ , then we can write, for  $x \in [x_0, x_1]$ ,

$$f(x) = f(a) + (x - a)f'(a) + \frac{(x - a)^2}{2!} f''(a) + \dots + \frac{(x - a)^n}{n!} f^{(n)}(a) + R_n. \quad (2.1.1)$$

Here  $R_n$  is a remainder term whose value is finite and equal to

$$R_n = \frac{(x - a)^{n+1}}{(n + 1)!} f^{(n+1)}(b(x)). \quad (2.1.2)$$

Here  $b \in [x_0, x_1]$  but unfortunately the exact value of  $b$  is not known. At best we can place an upper bound on the value of  $|R_n|$ , i.e. an upper bound on the accuracy of the sum.

- If  $|R_n| \rightarrow 0$  as  $n \rightarrow \infty$ , then we can extend the series to infinity and obtain a power series

$$\begin{aligned} f(x) &= f(a) + (x - a)f'(a) + \frac{(x - a)^2}{2!} f''(a) + \frac{(x - a)^3}{3!} f'''(a) + \dots \\ &= \sum_{n=0}^{\infty} \frac{(x - a)^n}{n!} f^{(n)}(a). \end{aligned} \quad (2.1.3)$$

### 2.1.1 Example

- Compute  $\sin(31^\circ)$ .
- Hence  $f(x) = \sin(x)$ . We set  $a = 30^\circ = \pi/6$ . We know that

$$f'(x) = \cos(x), \quad f''(x) = -\sin(x), \quad f'''(x) = -\cos(x), \quad \dots \quad (2.1.4)$$

- Also  $\sin(\pi/6) = \frac{1}{2}$  and  $\cos(\pi/6) = \frac{\sqrt{3}}{2}$ . Hence

$$\sin(x) = \sin(a) + (x-a)\cos(a) - \frac{(x-a)^2}{2!}\sin(a) - \frac{(x-a)^3}{3!}\cos(a) + \dots \quad (2.1.5)$$

- Substitute  $x - a = 1^\circ = \pi/180$  to obtain

$$\sin(31^\circ) = \frac{1}{2} + \left(\frac{\pi}{180}\right) \frac{\sqrt{3}}{2} - \frac{1}{2!} \left(\frac{\pi}{180}\right)^2 \frac{1}{2} - \frac{1}{3!} \left(\frac{\pi}{180}\right)^3 \frac{\sqrt{3}}{2} + \dots \quad (2.1.6)$$

- Sum the series until the answer converges to a desired tolerance.

## 2.2 Multinomial coefficient

- Recall the **binomial coefficient** is

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}. \quad (2.2.1)$$

- The **multinomial coefficient** is the generalization for  $m > 1$  variables  $k_1, k_2, \dots, k_m$

$$\begin{aligned} \binom{n}{k_1, k_2, \dots, k_m} &= \frac{n!}{k_1! k_2! \dots k_m! (n - k_1 - k_2 - \dots - k_m)!} \\ &= \frac{n(n-1) \dots (n+1 - \sum_i k_i)}{k_1! k_2! \dots k_m!}. \end{aligned} \quad (2.2.2)$$

- We must have  $k_1 + \dots + k_m \leq n$  else the value of the multinomial coefficient is zero.
- The multinomial coefficient has the following combinatorial interpretation. Suppose we have a set of  $n$  objects and we wish to place them in  $m$  bins. The multinomial coefficient counts the number of ways to place  $k_1$  objects in bin 1,  $k_2$  objects in bin 2, etc. and finally  $k_m$  objects in bin  $m$ . The special case  $m = 1$  is the binomial coefficient (“the number of ways to choose  $k$  objects from a set of  $n$  objects”).
- The value of the multinomial coefficient is therefore a positive integer.
- It is obvious that the value of  $n$  can be generalized to a non-integer value. Let  $x$  be a real (or complex) number.

The **generalized multinomial coefficient** is given by the finite product

$$\binom{x}{k_1, k_2, \dots, k_m} = \frac{x(x-1) \dots (x+1 - \sum_i k_i)}{k_1! k_2! \dots k_m!}. \quad (2.2.3)$$

### 2.3 Multinomial theorem (not for examination)

- The multinomial coefficient is used to specify the coefficients in the **multinomial theorem**.
- Let us begin with  $m = 2$  to illustrate.

$$\begin{aligned}
 (1 + x_1 + x_2)^n &= 1 + \binom{n}{1,0} x_1 + \binom{n}{0,1} x_2 \\
 &\quad + \binom{n}{2,0} x_1^2 + \binom{n}{1,1} x_1 x_2 + \binom{n}{0,2} x_2^2 \\
 &\quad + \binom{n}{3,0} x_1^3 + \binom{n}{2,1} x_1^2 x_2 + \binom{n}{1,2} x_1 x_2^2 + \binom{n}{0,3} x_2^3 \\
 &\quad + \dots \\
 &= 1 + \left[ \sum_{k_1=0}^n \sum_{k_2=0}^n \binom{n}{k_1, k_2} x_1^{k_1} x_2^{k_2} \right]_{k_1+k_2=1} \\
 &\quad + \left[ \sum_{k_1=0}^n \sum_{k_2=0}^n \binom{n}{k_1, k_2} x_1^{k_1} x_2^{k_2} \right]_{k_1+k_2=2} \\
 &\quad + \dots \\
 &= 1 + \sum_{j=1}^{\infty} \left[ \sum_{k_1=0}^n \sum_{k_2=0}^n \binom{n}{k_1, k_2} x_1^{k_1} x_2^{k_2} \right]_{k_1+k_2=j}.
 \end{aligned} \tag{2.3.1}$$

- The last line tells us how to generalize to  $m$  terms

$$\begin{aligned}
 (1 + x_1 + x_2 + \dots + x_m)^n \\
 = 1 + \sum_{j=1}^{\infty} \left[ \sum_{k_1=0}^n \dots \sum_{k_m=0}^n \binom{n}{k_1, k_2, \dots, k_m} x_1^{k_1} x_2^{k_2} \dots x_m^{k_m} \right]_{k_1+k_2+\dots+k_m=j}.
 \end{aligned} \tag{2.3.2}$$

- Just as with the binomial theorem, the series terminates if the value of  $n$  is a nonnegative integer, and the series does not terminate otherwise.

## 2.4 Multivariate Taylor series (not for examination)

- The multinomial coefficients appear in the formula for a Taylor series of a function of multiple variables  $x_1, x_2, \dots, x_m$ .
- Again, let us begin with  $m = 2$  to illustrate.

$$\begin{aligned}
 f(x_1 + \varepsilon_1, x_2 + \varepsilon_2) &= f(x_1, x_2) + \binom{1}{1,0} \frac{\partial f}{\partial x_1} \varepsilon_1 + \binom{1}{0,1} \frac{\partial f}{\partial x_2} \varepsilon_2 \\
 &\quad + \frac{1}{2!} \binom{2}{2,0} \frac{\partial^2 f}{\partial x_1^2} \varepsilon_1^2 + \frac{1}{2!} \binom{2}{1,1} \frac{\partial^2 f}{\partial x_1 \partial x_2} \varepsilon_1 \varepsilon_2 + \frac{1}{2!} \binom{2}{0,2} \frac{\partial^2 f}{\partial x_2^2} \varepsilon_2^2 \\
 &\quad + \frac{1}{3!} \binom{3}{3,0} \frac{\partial^3 f}{\partial x_1^3} \varepsilon_1^3 + \frac{1}{3!} \binom{3}{2,1} \frac{\partial^3 f}{\partial x_1^2 \partial x_2} \varepsilon_1^2 \varepsilon_2 \\
 &\quad \quad \quad + \frac{1}{3!} \binom{3}{1,2} \frac{\partial^3 f}{\partial x_1 \partial x_2^2} \varepsilon_1 \varepsilon_2^2 + \frac{1}{3!} \binom{3}{0,3} \frac{\partial^3 f}{\partial x_2^3} \varepsilon_2^3 + \dots \\
 &= f(x_1, x_2) + \frac{\partial f}{\partial x_1} \varepsilon_1 + \frac{\partial f}{\partial x_2} \varepsilon_2 \\
 &\quad + \frac{1}{2!} \frac{2!}{2!0!} \frac{\partial^2 f}{\partial x_1^2} \varepsilon_1^2 + \frac{1}{2!} \frac{2!}{1!1!} \frac{\partial^2 f}{\partial x_1 \partial x_2} \varepsilon_1 \varepsilon_2 + \frac{1}{2!} \frac{2!}{0!2!} \frac{\partial^2 f}{\partial x_2^2} \varepsilon_2^2 \\
 &\quad + \frac{1}{3!} \frac{3!}{3!0!} \frac{\partial^3 f}{\partial x_1^3} \varepsilon_1^3 + \frac{1}{3!} \frac{3!}{2!1!} \frac{\partial^3 f}{\partial x_1^2 \partial x_2} \varepsilon_1^2 \varepsilon_2 \\
 &\quad \quad \quad + \frac{1}{3!} \frac{3!}{1!2!} \frac{\partial^3 f}{\partial x_1 \partial x_2^2} \varepsilon_1 \varepsilon_2^2 + \frac{1}{3!} \frac{3!}{0!3!} \frac{\partial^3 f}{\partial x_2^3} \varepsilon_2^3 + \dots \\
 &= f(x_1, x_2) + \frac{\partial f}{\partial x_1} \varepsilon_1 + \frac{\partial f}{\partial x_2} \varepsilon_2 \\
 &\quad + \frac{1}{2!0!} \frac{\partial^2 f}{\partial x_1^2} \varepsilon_1^2 + \frac{1}{1!1!} \frac{\partial^2 f}{\partial x_1 \partial x_2} \varepsilon_1 \varepsilon_2 + \frac{1}{0!2!} \frac{\partial^2 f}{\partial x_2^2} \varepsilon_2^2 \\
 &\quad + \frac{1}{3!0!} \frac{\partial^3 f}{\partial x_1^3} \varepsilon_1^3 + \frac{1}{2!1!} \frac{\partial^3 f}{\partial x_1^2 \partial x_2} \varepsilon_1^2 \varepsilon_2 \\
 &\quad \quad \quad + \frac{1}{1!2!} \frac{\partial^3 f}{\partial x_1 \partial x_2^2} \varepsilon_1 \varepsilon_2^2 + \frac{1}{0!3!} \frac{\partial^3 f}{\partial x_2^3} \varepsilon_2^3 + \dots \\
 &= f(x_1, x_2) + \sum_{j=1}^{\infty} \left[ \sum_{k_1=0}^n \sum_{k_2=0}^n \frac{1}{k_1! k_2!} \frac{\partial^j f}{\partial x_1^{k_1} \partial x_2^{k_2}} \varepsilon_1^{k_1} \varepsilon_2^{k_2} \right]_{k_1+k_2=j}.
 \end{aligned} \tag{2.4.1}$$

- The last line tells us how to generalize to  $m$  terms

$$\begin{aligned}
 f(x_1 + \varepsilon_1, x_2 + \varepsilon_2, \dots, x_m + \varepsilon_m) &= f(x_1, x_2, \dots, x_m) \\
 &\quad + \sum_{j=1}^{\infty} \left[ \sum_{k_1=0}^n \sum_{k_2=0}^n \dots \sum_{k_m=0}^n \frac{1}{k_1! k_2! \dots k_m!} \frac{\partial^j f}{\partial x_1^{k_1} \partial x_2^{k_2} \dots \partial x_m^{k_m}} \varepsilon_1^{k_1} \varepsilon_2^{k_2} \dots \varepsilon_m^{k_m} \right]_{k_1+k_2+\dots+k_m=j}.
 \end{aligned} \tag{2.4.2}$$