



Disentangling within- and between-female egg size–number trade-offs in the European earwig

Journal:	<i>Current Zoology</i>
Manuscript ID	CZ-2026-0008.R3
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Kelly, Clint; UQAM Departement des sciences biologiques
Keywords:	fecundity, life-history trade off, residual correlation, mixed models, egg size, clutch size

SCHOLARONE™
Manuscripts

Disentangling within- and between-female egg size–number trade-offs in the European earwig.

Clint D. Kelly^{1,✉}

¹ Département des Sciences biologiques, Université du Québec à Montréal, Montréal, Canada

✉ Correspondence: [Clint D. Kelly <kelly.clint@uqam.ca>](mailto:kelly.clint@uqam.ca)

For Review Only

Abstract

Life-history theory predicts a trade-off between offspring size and number, yet empirical studies often fail to detect the expected negative association. This discrepancy may occur because phenotypic correlations confuse variation in resource acquisition with variation in allocation strategy. Using repeated measures of female European earwigs (*Forficula auricularia*), I partitioned covariance between egg size and egg number into among- and within-female components using Bayesian mixed models. The raw phenotypic correlation was near zero. However, females that produced more eggs on average laid smaller eggs, revealing a modest negative association among females that aligns with an allocation trade-off. In contrast, there was no detectable within-female covariance once clutch order was accounted for. Egg number repeatability was effectively zero, suggesting that clutch size varies mainly within rather than among females, while egg size was highly repeatable. Both egg number and egg size decreased in second clutches, and within-clutch egg-size variance increased, indicating a reproductive constraint (i.e., fewer functional ovarioles) under a possibly diminished resource state. Together, these findings demonstrate that the size–number trade-off appears to be expressed among females through consistent differences in allocation strategy, whereas clutch size itself remains flexible. By separating hierarchical sources of variation, this study shows that allocation trade-offs can remain hidden at the phenotypic level and that evolutionary constraints act through stable allocation rules rather than through flexible clutch-size adjustments.

Introduction

Life-history theory predicts that organisms must allocate finite resources among competing fitness components. Because energy and time are limited, investment in one trait constrains investment in others (Roff, 1992; Stearns, 1992). A central expression of this principle is the offspring size–number trade-off: for a given reproductive budget, producing larger offspring reduces the number that can be produced (Parker, 1982; Smith and Fretwell, 1974). Since larger eggs often enhance offspring performance and survival (Clutton-Brock, 1991), selection is expected to favour an optimal balance between egg size and egg number that maximises maternal fitness (Fox and Czesak, 2000). Yet empirical tests of the size–number trade-off often fail to reveal the expected negative relationship. Across individuals within populations, egg size and egg number are often positively correlated (Bernardo, 1996; Fox and Czesak, 2000; Hargitai et al., 2005; Roff, 1992). This apparent contradiction arises because phenotypic correlations conflate two distinct biological processes: variation in resource acquisition and variation in allocation strategy. Van Noordwijk and De Jong (1986) showed that when variance in acquisition exceeds variance in allocation, individuals with greater resource pools can produce both more and larger eggs, generating positive among-individual correlations even if all individuals are constrained by the same underlying allocation trade-off. In such cases, the population-level pattern masks allocation constraints operating within individuals—an outcome conceptually analogous to Simpson’s paradox (Simpson, 1951).

From this perspective, the critical prediction of life-history theory is not simply a negative phenotypic correlation, but a negative within-individual covariance between egg size and egg number once among-individual differences in acquisition are accounted for (Careau and Wilson,

2017). Detecting such a masked trade-off, therefore, requires partitioning phenotypic variance and covariance into hierarchical components. Repeated measures of individuals allow estimation of (i) among-individual variance, reflecting consistent differences among females in total reproductive investment, and (ii) within-individual variance, reflecting how females adjust allocation across reproductive bouts. Careau and Wilson (2017) argue that only by decomposing covariance into these components using multivariate mixed models can we determine whether allocation trade-offs operate within individuals despite positive population-level associations. Crucially, this framework also predicts that traits involved in allocation trade-offs should show substantial repeatability. If females differ consistently in resource acquisition, egg size and egg number should exhibit among-individual variance across clutches. High repeatability would, therefore, indicate persistent differences in total reproductive investment among females, consistent with variation in acquisition. The marked increase in within-clutch egg size variance in second broods further supports a resource constraint interpretation. Under reduced resource availability, females may be less able to maintain uniform provisioning across all eggs, leading to greater heterogeneity in egg size within a clutch. Such increased variance is consistent with diminished allocation precision when reproductive resources are limited, rather than an adaptive diversification of offspring size.

Despite the foundational importance of the size–number trade-off, empirical tests rarely partition covariance into within- and among-individual components to distinguish allocation from acquisition effects. Here, I address this gap using the European earwig, *Forficula auricularia*, a species that allows repeated measures of reproductive investment within females but in which previous work has reported no clear or consistent pattern in the association between egg size and

1
2
3
4 71 egg number (Koch and Meunier, 2014), highlighting the difficulty of interpreting phenotypic
5
6 72 correlations without partitioning hierarchical sources of variation. Female earwigs lay a clutch
7
8 73 within a short temporal window in a subterranean burrow (Meunier, 2024) and provide extended
9
10 74 maternal care (Meunier, 2024; Staerke and Kölliker, 2008). In Québec populations,
11
12 75 environmental constraints typically render females effectively semelparous (Tourneur, 2018), but
13
14 76 under favourable laboratory conditions, they can produce a second clutch. This system, therefore,
15
16 77 enables direct estimation of within-female covariance across successive reproductive bouts. I test
17
18 78 three predictions derived from acquisition–allocation theory (Van Noordwijk and Jong, 1986; see
19
20 79 also Reznick, Nunney, and Tessier, 2000). First, if variance in resource acquisition among
21
22 80 females exceeds variance in allocation strategy, then egg size and egg number should exhibit
23
24 81 substantial repeatability and a positive among-individual correlation, whereas the within-
25
26 82 individual correlation should be negative, revealing a masked allocation trade-off. Second, if
27
28 83 maternal fitness is maximised at a particular offspring size or number (Smith and Fretwell, 1974),
29
30 84 then one trait should remain relatively invariant across clutches while the other adjusts between
31
32 85 reproductive bouts (Winkler and Wallin, 1987). Third, if reproductive investment is structured
33
34 86 differently across bouts, then variance in egg size may differ between first and second clutches at
35
36 87 both the among-female and within-clutch levels (e.g. Hargitai et al., 2005), providing further
37
38 88 insight into how selection shapes the distribution of maternal investment across reproductive
39
40 89 events.

41
42
43
44
45
46
47
48 90 By separating among-female differences in total reproductive investment from within-female
49
50 91 allocation decisions, this study tests a fundamental question in life-history theory: do apparent
51
52 92 positive associations between egg size and number reflect relaxed constraints, or do allocation
53
54
55
56
57
58
59
60

trade-offs persist but remain hidden within individuals? Demonstrating a negative within-female covariance despite positive population-level patterns would show that acquisition can mask, but not erase, the constraints imposed by finite resources. In this way, my study evaluates not only the existence of a trade-off but also the hierarchical structure through which life-history constraints are expressed.

Methods

I collected adult female earwigs (n=42) in September and October 2024 in an urban park (Parc Outremont) in Montréal, Canada. All earwigs were brought into the laboratory, assigned a unique ID, and individually placed in a 90-mm-diameter Petri dish containing a thin layer of moist sand. Females were provided a cotton-plugged water vial and fed TetraMin Tropical fish flakes ad libitum. Females were maintained in an incubator (Percival Scientific, Iowa, USA) at 10°C in constant darkness.

Females were checked daily for oviposition. When a female laid eggs, I recorded the date, collected all eggs from the sand, and placed them on filter paper in a new Petri dish. The eggs were then photographed under a Leica S6D stereo microscope (Leica Microsystems Inc., Concord, ON, Canada) at 0.68x magnification using Enersight software (Leica Microsystems Inc., Concord, ON, Canada), which automatically adds a scale bar to the photo. Eggs were then discarded.

To stimulate females to lay a second clutch of eggs, I recreated a brief spring period during which they were allowed to accumulate resources for egg production and replenish their sperm stores. To achieve this, females that had recently oviposited were placed in a new Petri dish containing a

1
2
3
4 114 cotton-plugged water vial, TetraMin Tropical fish flakes, and a shelter consisting of four 4 cm-
5
6 115 long straws (Blowholes, Markham, Canada) glued together. These Petri dishes were placed in a
7
8 116 23 °C incubator with a 12L:12D light: dark cycle. Females were maintained under these
9
10 117 conditions for two weeks, at which time a haphazardly chosen stock male from our colony was
11
12 118 placed with her. The pair was kept together for a week, after which a new stock male replaced the
13
14 119 original, and the new pair was kept together for another week. Females were then transferred to
15
16 120 the 10 °C incubator and treated as previously until they laid their second clutch, at which point
17
18 121 their eggs were photographed. Females were euthanised by freezing after laying their second
19
20 122 clutch, and their pronotum was photographed as described above. Because female fecundity is
21
22 123 often related to body size in insects (Honek, 1993) and could thus possibly explain a positive
23
24 124 among-female covariance between egg number and size, I measured two proxies of size,
25
26 125 pronotum length and body mass. Females were weighed to the nearest 0.01 mg using a Sartorius
27
28 126 Secura 224-1S analytical balance. Females were weighed three times, with each weighing bout
29
30 127 separated by four days. I used Fiji image analysis software (Schindelin et al., 2012) to measure
31
32 128 the length of the female pronotum to the nearest 0.001 mm three times, with each measurement
33
34 129 separated by 4 days. I used the same procedure to measure the perimeter of each egg.
35
36
37
38
39
40

41 130 **Statistical analysis**

42
43 131 I calculated the repeatability of the body mass, pronotum length and egg perimeter measurements
44
45 132 using the R package *rptR* (Stoffel, Nakagawa, and Schielzeth, 2017). I entered the trait of interest
46
47 133 as the response variable, female ID as a random effect, and assumed a Gaussian distribution.
48
49
50 134 Each analysis was based on 1000 bootstrap samples. The measurements of my two
51
52 135 morphological traits were highly repeatable (pronotum length: R [95% CI] = 0.98 [0.97, 0.99];
53
54
55
56
57
58
59
60

body mass: R [95% CI] = 0.99 [0.988, 0.996]); therefore, I used each female's mean value for each trait where necessary. Because egg perimeter was highly repeatable in both clutches (first clutch: R [95% CI] = 0.977 [0.976, 0.979]; second clutch: R [95% CI] = 0.93 [0.926, 0.937]), I calculated each egg's average perimeter and then used these values to calculate the mean egg size for each clutch for each female.

To reduce the two correlated measures of body size (pronotum length and body mass) to a single structural size metric, I performed a principal components analysis (PCA) on the correlation matrix of standardized variables using the *prcomp* function in the R stats package (R Core Team, 2025). The first principal component (PC1) explained 60.8% of the total variance and was used as an index of structural body size. In addition, female body condition was estimated using the scaled mass index (SMI) following Peig and Green (2010, 2009; see also Kelly, Tawes, and Worthington, 2014), which standardizes body mass relative to structural size (pronotum length).

I first quantified associations among reproductive traits using raw phenotypic correlations. Pearson correlation coefficients were calculated using individual-level trait values, thereby capturing total phenotypic covariation across all sources of variation. These correlations reflect the combined contribution of consistent differences among females, variation among clutches within females, and residual effects. Consequently, raw phenotypic correlations provide a descriptive estimate of overall trait associations but do not distinguish between covariation arising from stable female differences and that arising from within-female (i.e., among-clutch) processes.

1
2
3
4 156 To distinguish covariance arising from within-female plasticity from covariance attributable to
5
6 157 consistent among-female differences, I followed the within–between decomposition approach of
7
8 158 van de Pol and Wright [(2009); see Supplementary Material for details]. This decomposition is
9
10 159 necessary because residual covariance does not isolate within-individual effects and, in models
11
12 160 with different response distributions, does not provide a directly interpretable estimate of within-
13
14 161 female trade-offs. Although mean-centering has been shown to impose assumptions about
15
16 162 underlying biological processes (Westneat et al., 2020), these assumptions are appropriate here
17
18 163 because the biological question concerns how reproductive allocation varies within females
19
20 164 across clutches versus how females differ consistently in allocation strategy. Egg number was
21
22 165 decomposed into two components: a between-female component, defined as each female’s mean
23
24 166 egg number across broods centered on the population mean, and a within-female component,
25
26 167 defined as the deviation of each brood from that female’s own mean. Hereafter, we refer to
27
28 168 “among-female” variation as differences between individuals and “within-female” variation as
29
30 169 differences among clutches within individuals. These two predictors were included
31
32 170 simultaneously in a Bayesian multivariate mixed-effects model in which egg number and mean
33
34 171 egg size were modelled jointly. Egg number was modelled using a negative binomial distribution
35
36 172 with a log link to account for overdispersion, whereas mean egg size was modelled assuming a
37
38 173 Gaussian distribution with an identity link. Clutch (first vs. second clutch) was included as a
39
40 174 fixed effect in both submodels. Female identity was included as a random intercept for both
41
42 175 response variables. This structure allowed estimation of among-female variance in egg number,
43
44 176 among-female variance in egg size, and the covariance (correlation) between female-level
45
46 177 intercepts for the two traits.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 178 This multivariate framework partitions total phenotypic covariance into components arising from
5
6 179 consistent differences among females and from within-female deviations across clutches.
7
8 180 Covariance attributable to consistent differences among females is captured by the female-level
9
10 181 random effects, whereas covariance arising from within-female variation is captured by the fixed
11
12 182 within-female component of egg number. The fixed between-female component estimates how
13
14 183 mean egg size varies with mean egg number across females, thereby capturing systematic (i.e.,
15
16 184 explained) differences in allocation strategy among females. In contrast, the fixed within-female
17
18 185 component tests whether egg size changes plastically when a female produces more or fewer
19
20 186 eggs than her own mean. The correlation between female-level intercepts represents residual
21
22 187 (latent) covariance among baseline trait values that is not explained by the fixed effects, including
23
24 188 the between-female allocation relationship. Thus, the between-female slope tests whether females
25
26 189 diverge in their mean allocation strategy (i.e., whether females that consistently produce more
27
28 190 eggs also produce smaller eggs), whereas the intercept correlation quantifies any remaining
29
30 191 covariance between egg number and egg size after accounting for these effects. Continuous
31
32 192 predictors were standardized prior to analysis, such that standardized coefficients represent the
33
34 193 expected change in egg size (in standard deviation units) associated with a one standard deviation
35
36 194 change in egg number at the relevant hierarchical level.
37
38
39
40
41
42

43 195 To quantify the structure of egg-size variation specifically, I fitted a univariate mixed-effects
44
45 196 model in which egg size (standardised mean egg perimeter) was modelled as a Gaussian response
46
47 197 with female identity included as a random intercept (see Supplementary Material for details).
48
49 198 This formulation partitions total phenotypic variance in egg size into variance attributable to
50
51 199 differences among females and variance occurring within females among eggs within clutches.
52
53
54
55
56
57
58
59
60

1
2
3
4 200 The among-female variance component reflects consistent maternal differences in egg size
5
6 201 allocation, whereas the within-female component captures heterogeneity in provisioning within
7
8 202 clutches and residual variation.
9

10
11 203 Repeatability was calculated on the observation scale for both traits. For egg size (Gaussian
12
13 204 model):
14
15

16
17 205
$$R_{\text{egg size}} = \frac{V_F}{V_F + V_R}$$

18
19

20
21 206 where V_F is the among-female variance (i.e., variance of the female random intercept) and V_R is
22
23 207 the residual variance. Because the Gaussian model is parameterized directly on the observation
24
25 208 scale, this formulation corresponds to both latent- and observation-scale repeatability.
26
27

28
29 209 For egg number, which was modelled using a negative binomial distribution with a log link, the
30
31 210 variance depends on both the expected mean and the dispersion (shape) parameter. Under the
32
33 211 negative binomial parameterization used in *brms*, the observation-level (distribution-specific)
34
35 212 variance is given by:
36
37

38
39 213
$$\text{Var}(Y) = \mu + \frac{\mu^2}{\theta}$$

40
41

42
43 214 where μ is the expected mean egg number and θ is the dispersion (shape) parameter. Because the
44
45 215 model includes a log link, the among-female variance estimated from the random intercept is
46
47 216 expressed on the latent (log) scale and must be transformed to the observation scale before
48
49 217 calculating repeatability. Following Nakagawa et al. (2017), the among-female variance on the
50
51 218 observation scale was therefore calculated as:
52
53
54
55
56
57
58
59
60

$$V_F^{\text{obs}} = \mu^2(\exp(\sigma^2) - 1)$$

where σ^2 is the variance of the random intercept on the latent scale.

Observation-scale repeatability for egg number was then calculated as:

$$R_{\text{egg number}} = \frac{V_F^{\text{obs}}}{V_F^{\text{obs}} + \left(\mu + \frac{\mu^2}{\theta}\right)}$$

Both the transformed among-female variance and the distribution-specific variance were computed for each posterior draw using the corresponding posterior samples of μ , σ^2 , and θ . To account for variation in expected values across observations, these quantities were first calculated at the observation level and then averaged within each posterior draw. This procedure yielded a posterior distribution of repeatability that fully propagates uncertainty from all variance components.

Repeatability for non-Gaussian traits was calculated following the framework of Nakagawa and Schielzeth (2010) and its extensions for generalized linear mixed models (Nakagawa, Johnson, and Schielzeth, 2017). Because distribution-specific variance can contribute substantially to total phenotypic variance in negative binomial models, repeatability estimates for count data are often lower than those obtained under Gaussian assumptions, even when among-individual variance is present (Nakagawa, Johnson, and Schielzeth, 2017; Nakagawa and Schielzeth, 2010). By calculating repeatability on the observation scale, these estimates represent the proportion of total phenotypic variance in observed egg number attributable to consistent differences among females.

I used weakly informative priors to constrain parameters to biologically realistic ranges while allowing the data to drive inference. Regression coefficients were assigned normal priors centered at zero (Normal(0, 0.5–1)), reflecting the expectation that clutch effects on egg traits would be moderate in magnitude. Intercepts were given normal priors scaled to each trait (egg number: Normal(0, 5); mean egg size: Normal(3.5, 1); standardized egg size: Normal(0, 0.5)). Among-female variation was assigned exponential priors (Exponential(1–2)), which favour small but non-zero individual differences. Residual variance (Gaussian models) and the negative binomial dispersion parameter were given Exponential(1) priors. Correlations among female-level random effects were regularized using an LKJ(2) prior to avoid overestimating trait associations. Posterior distributions were obtained using six Markov chains each with 4,000 iterations, including 1,000 warm-up iterations, and convergence was assessed using $\hat{R}$ statistics, effective sample sizes, and trace-plot inspection. Posterior distributions are summarised as posterior means with 95% credible intervals. Data were visualized using *ggplot* (Wickham, 2016) and *tidybayes* (Kay, 2020).

Results

Between- and within-female associations

Across all broods, the raw phenotypic correlation between egg number and egg size was weak and near zero ($r_{raw} = -0.04$, 95% CI [-0.25, 0.19]), indicating no detectable population-level association when hierarchical structure was ignored.

Egg number (clutch one: $\bar{x} = 51.55 \pm 2.49$ eggs; clutch two: $\bar{x} = 34.14 \pm 2.65$ eggs) and egg size (first clutch: $\bar{x} = 3.54 \pm 0.03$ mm; second clutch: $\bar{x} \pm SE = 3.41 \pm 0.03$ mm) both declined in

second clutches. Egg number decreased on the log scale ($\beta = -0.41$, 95% CrI (-0.64, -0.19; Figure 1; Supplementary Table 1), corresponding to an approximate 34% reduction on the natural scale ($e^{-0.413} \approx 0.66$). Mean egg size also declined in the second clutch ($\beta = -0.13$, 95% CrI (-0.24, -0.02)).

The results did not support my prediction of a within-female plastic trade-off between egg number and egg size (Figure 1 a). The within-female deviation in egg number had a slope near zero ($\beta_W = -0.01$, 95% CrI (-0.06, 0.05), indicating that females did not systematically adjust egg size when producing more or fewer eggs relative to their own mean. By comparison, the among-female association (captured by the fixed between-female slope) revealed a negative relationship between egg size and mean female egg number ($\beta_B = -0.05$, 95% CrI (-0.10, 0.00; Figure 1 b). The posterior distribution of the among-female slope was centered on negative values, suggesting weak evidence for divergence in mean allocation strategy among females. The correlation between female-level intercepts for egg number and egg size was highly uncertain (median = -0.02, 95% CrI: -0.81, 0.82), indicating limited information for estimating latent covariance with only two broods per female. With only two repeated measures per female, intercept correlations are weakly identifiable and should be interpreted cautiously.

No measure of female size/condition was positively correlated with the number of eggs in the first brood (pronotum length: $r = 0.14$, 95% CI [-0.19, 0.43]; body mass: $r = 0.12$, 95% CI [-0.21, 0.43]; PC1: $r = -0.17$, 95% CI [-0.44, 0.15]; scaled mass index: $r_{raw} = -0.02$, 95% CI [-0.35, 0.33]) providing no evidence that larger females or those in better condition produce more eggs. I therefore did not include a measure of size/condition in this model. Together, these results show that the size–number trade-off is not apparent at the raw phenotypic level and is not expressed as

within-female plasticity, but emerges as a modest negative association at the among-female level through differences in average allocation strategy.

Egg size variation between females and within clutches

Egg size exhibited substantial variation at both the between-female and within-clutch levels (Figure 2; Supplementary Table 2). Between-female variance was similar across reproductive events, with a median of 0.73 (95% CrI: 0.50, 1.12) in first clutches and 0.61 (95% CrI: 0.41, 0.96) in second clutches. The posterior median difference (second – first) was -0.11 (95% CrI: -0.56, 0.28), and the posterior probability that between-female variance was greater in second clutches was 0.28, indicating no clear evidence of a difference. In contrast, within-clutch (egg-to-egg) variance was substantially higher in second clutches (median = 0.35 (95% CrI: 0.33, 0.38) than in first clutches (median = 0.21 (95% CrI: 0.20, 0.22). The posterior median difference was 0.14 (95% CrI: 0.12, 0.17), with a posterior probability of 1.00, that within-clutch variance was greater in second clutches.

Egg number and egg size repeatability

Repeatability was calculated from posterior samples of the variance components on the observation scale. Egg number exhibited extremely low repeatability. Although there was some among-female variance in egg number (SD = 0.078, 95% CrI [0.003, 0.270]; Supplementary Table 1), the overwhelming majority of variance arose from distribution-specific (negative binomial) variance. Consequently, observation-scale repeatability was near zero ($R = 4.86 \times 10^{-2}$, 95% CrI [3.80×10^{-5} , 2.43×10^{-1}]), indicating that very little of the total phenotypic variation in clutch size was attributable to consistent differences among females.

Because variance components and repeatability were derived from different models, these quantities are reported separately. Among-female variance in egg size from the multivariate model was modest in absolute terms ($SD = 0.082$, 95% CrI [0.006, 0.155]; residual $SD = 0.191$, 95% CrI [0.155, 0.235]; Supplementary Table 1), providing a sense of the trait's physical scale (mm). Repeatability, however, was estimated from the clutch-specific univariate model (Supplementary Table 2), in which among-female variance exceeded residual variance ($SD = 0.855$ vs 0.457 in first broods; 0.784 vs 0.594 in second broods), yielding $R = 0.78$ (95% CrI: $0.70, 0.84$) in first broods and $R = 0.63$ (95% CrI: $0.53, 0.73$) in second broods.

Discussion

Life-history theory predicts a trade-off between offspring size and number because both traits draw from a finite pool of reproductive resources (Lack, 1947; Roff, 1992; Smith and Fretwell, 1974; Stearns, 1992). Yet, as emphasised by van Noordwijk and de Jong (1992), phenotypic correlations need not directly reveal allocation trade-offs, because variation in resource acquisition can obscure underlying allocation rules (Careau and Wilson, 2017; Johnson and Nassrullah, 2024). This acquisition–allocation framework has become central to interpreting life-history covariance (Houle, 1991; Reznick, Nunney, and Tessier, 2000). By partitioning covariance into within- and among-female components, my results show that the size–number trade-off in this population of earwigs is modestly expressed among females but absent within them, illustrating how hierarchical structure shapes the detectability of life-history trade-offs (Careau and Wilson, 2017; Dingemanse and Dochtermann, 2013; Pol and Wright, 2009; Westneat et al., 2020).

1
2
3
4 323 At the phenotypic level, egg size and egg number were uncorrelated in my earwig population,
5
6 324 consistent with Koch and Meunier (2014). Under the framework of van Noordwijk and de Jong
7
8 325 (1986), such phenotypic independence is expected when variation in resource acquisition
9
10 326 obscures underlying allocation trade-offs. After hierarchical decomposition, however, a weak
11
12 327 negative association emerged at the among-female level: females that produced more eggs on
13
14 328 average tended to lay slightly smaller eggs. This pattern is captured by the fixed between-female
15
16 329 slope and reflects differences among females in their mean allocation strategy. In contrast, the
17
18 330 correlation between female-level intercepts—representing residual (latent) covariance after
19
20 331 accounting for this allocation relationship—was highly uncertain and overlapped zero, indicating
21
22 332 little evidence for additional integration between traits beyond that explained by differences in
23
24 333 mean allocation strategy (Roff and Fairbairn, 2007; Stearns, 1989).
25
26
27
28
29 334 In contrast, there was no evidence for a within-female plastic trade-off. Although each female
30
31 335 contributed only two broods, the posterior distribution of the within-female slope was tightly
32
33 336 centered on zero, suggesting that any within-female effect, if present, is small. Females did not
34
35 337 reduce egg size when producing more eggs than their own average; instead, egg number and egg
36
37 338 size shifted together across broods. This pattern is consistent with state-dependent life-history
38
39 339 models in which allocation rules remain relatively stable while total resource state fluctuates
40
41 340 across reproductive bouts (Houston and McNamara, 1999; McNamara and Houston, 1996).
42
43 341 Rather than dynamically reallocating resources between offspring size and number, females
44
45 342 appear to exhibit clutch-level shifts driven by overall energetic state.
46
47
48
49
50
51 343 Notably, egg number was not correlated with female structural size or body condition (i.e. scaled
52
53 344 mass), indicating that persistent differences in morphology do not explain clutch-size variation. If
54
55
56
57
58
59
60

stable acquisition differences were driving among-female variation, we would expect egg number to show repeatability and to covary with these morphological size/condition indicators. The absence of such patterns weakens the interpretation that persistent acquisition differences—at least as reflected by morphology—underlie among-female differences in reproductive output. Instead, clutch size appears predominantly state-sensitive and influenced by clutch-level constraints.

This interpretation also helps explain the pronounced decline in second clutches. That second clutches comprised both fewer and smaller eggs is consistent with previous work in North American and European populations (Koch and Meunier, 2014; Tourneur and Gingras, 1992). Tourneur and Gingras (1992) proposed that some ovarioles become non-functional after the first clutch, thereby reducing egg production capacity. Structural limitation could therefore contribute to smaller second clutches. However, structural reduction alone cannot explain the simultaneous decline in egg size. Under a strict allocation trade-off, a reduction in egg number would be expected to lead to compensatory increases in egg size if total resources remain constant. Instead, both traits declined, indicating that females likely entered the second reproductive bout with diminished resource reserves in addition to any physiological constraint. In the wild, females accumulate resources for months before producing their first clutch, whereas in the laboratory, they had only a brief interval (approx. one month) to replenish reserves before producing the second. Late reproduction, therefore, likely reflects both reduced energetic state and structural limitation.

The hierarchical variance structure supports a constraint-based interpretation of clutch-size variation. Egg size exhibited moderate repeatability across broods, indicating stable differences

1
2
3
4 367 among females in mean per-offspring investment. In contrast, egg number showed extremely low
5
6 368 repeatability, with most of the variance arising from brood effects and distribution-level variation
7
8 369 rather than from persistent female differences. The absence of any association between egg
9
10 370 number and female structural size or condition, together with the near-zero repeatability of clutch
11
12 371 size, suggests that variation in egg number is not driven by stable differences in resource
13
14
15 372 acquisition but instead reflects transient physiological state or clutch-level constraint. Thus,
16
17 373 clutch size appears primarily state-sensitive, whereas egg size reflects a more stable allocation
18
19 374 strategy (Houston and McNamara, 1999; McNamara and Houston, 1996). At the same time,
20
21 375 within-clutch variance increased substantially in second broods, indicating reduced
22
23 376 developmental precision under diminished resource availability. Structural and energetic
24
25 377 constraints, therefore, appear to reduce total reproductive output while leaving mean allocation
26
27 378 rules largely intact (Ricklefs and Wikelski, 2002; Zera and Harshman, 2001).
28
29
30
31 379 These patterns have broader implications for phenotypic integration and evolutionary potential.
32
33 380 Repeatability places an upper bound on heritability (Falconer and MacKay, 1996; Lessells and
34
35 381 Boag, 1987; Nakagawa and Schielzeth, 2013; Nakagawa and Schielzeth, 2010; but see Dohm,
36
37 382 2002). The moderate repeatability of egg size suggests that selection could act on persistent
38
39 383 differences in mean provisioning, whereas the extremely low repeatability of egg number implies
40
41 384 that clutch size may respond primarily to short-term environmental or physiological conditions
42
43 385 (i.e. reduced ovariole function). The weak intercept correlation further indicates that egg size and
44
45 386 egg number are only loosely integrated, consistent with models in which traits differ in their
46
47 387 sensitivity to resource state versus allocation strategy (Houle, 1991). In this system, per-offspring
48
49 388 investment appears more evolutionarily stable than the total number of offspring.
50
51
52
53
54
55
56
57
58
59
60

Viewed hierarchically, my results illustrate a broader conceptual point: life-history trade-offs may operate primarily through stable differences in allocation rules among individuals, while short-term variation reflects state-dependent constraint. Without decomposing within- and among-individual covariance, the modest among-female trade-off suggested here would have remained obscured. More generally, the evolutionary consequences of trade-offs depend not only on their sign but on their hierarchical expression and repeatability. A trade-off that reflects consistent among-individual differences in a repeatable trait is more likely to shape evolutionary responses than one arising primarily from transient, plastic variation within individuals. Future work measuring resource acquisition directly, experimentally manipulating body condition, or estimating genetic correlations between egg size and egg number will help determine whether the allocation divergence observed here reflects adaptive differentiation in allocation strategy or condition-dependent constraint (Lande, 1980; Roff, 1992).

Data availability

Data are available through the OSF repository at https://osf.io/jrs28/overview?view_only=7a78f30e724b497ab3bb62ffa1c97e10.

Author contributions

CDK collected and analyzed the data, and wrote the manuscript.

Funding

The study was funded by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada (NSERC).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conflict of Interest

I declare no conflict of interest.

Acknowledgements

I thank Andréanne Nault and Clara Smythe for assistance in animal care and data collection. I also thank Denis Réale, Pierre-Olivier Montiglio, and two anonymous reviewers for their discussion and feedback.

For Review Only

References

- Bernardo, J., 1996. The particular maternal effect of propagule size, especially egg size: Patterns, models, quality of evidence and interpretations. *Am Zool* **36**, 216–236.
- Careau, V., Wilson, R., 2017. Of uberfleas and krakens: Detecting trade-offs using mixed models. *Integr Comp Biol* **57**, 362–371.
- Clutton-Brock, T.H., 1991. The Evolution of Parental Care. Princeton University Press.
- De Jong, G., Van Noordwijk, A., 1992. Acquisition and allocation of resources: Genetic (co) variances, selection, and life histories. *Am Nat* **139**, 749–770.
- Dingemanse, N.J., Dochtermann, N., 2013. Quantifying individual variation in behaviour: Mixed-effect modelling approaches. *J Anim Ecol* **82**, 39–54.
- Dohm, M.R., 2002. Repeatability estimates do not always set an upper limit to heritability. *Funct Ecol* **16**, 273–280.
- Falconer, D.S., MacKay, T.F.C., 1996. An Introduction Quantitative Genetics. Pearson Education Limited, Essex, England.
- Fox, C.W., Czesak, M.E., 2000. Evolutionary ecology of progeny size in arthropods. *Annu Rev Entomol* **45**, 341–369.
- Hargitai, R., Török, J., Tóth, L., Hegyi, G., Rosivall, B., Szigeti, B., Szöllősi, E., 2005. Effects of environmental conditions and parental quality on inter-and intraclutch egg-size variation in the collared flycatcher (*Ficedula albicollis*). *Auk* **122**, 509–522.
- Honek, A., 1993. Intraspecific variation in body size and fecundity in insects: A general relationship. *Oikos* **483–492**.
- Houle, D., 1991. Genetic covariance of fitness correlates: What genetic correlations are made of and why it matters. *Evolution* **45**, 630–648.
- Houston, A.I., McNamara, J.M., 1999. Models of adaptive behaviour: An approach based on state. Cambridge University Press.
- Johnson, M.T.J., Nassrullah, Z., 2024. The improbability of detecting trade-offs and some practical solutions. *J Evol Biol* **37**, 1205–1214.
- Kay, M., 2020. tidybayes: Tidy Data and Geoms for Bayesian Models [WWW Document]. *R package version 3.0.5*. URL <http://mjskay.github.io/tidybayes/>
- Kelly, C.D., Tawes, B., Worthington, A., 2014. Evaluating indices of body condition in two cricket species. *Ecol Evol* **4**, 4476–4487.

1
2
3
4 447 Koch, L.K., Meunier, J., 2014. Mother and offspring fitness in an insect with maternal care:
5 448 Phenotypic trade-offs between egg number, egg mass and egg care. *BMC Evol Biol* **14**, 125.
6
7 449 Lack, D., 1947. The significance of clutch-size in the partridge (*Perdix perdix*). *J Anim Ecol*
8 450 **16**, 19–25.
9
10 451 Lande, R., 1980. The genetic covariance between characters maintained by pleiotropic
11 452 mutations. *Genetics* **94**, 203–215.
12
13 453 Lessells, C., Boag, P.T., 1987. Unrepeatable repeatabilities: A common mistake. *Auk* **104**,
14 454 116–121.
15
16 455 McNamara, J.M., Houston, A.I., 1996. State-dependent life histories. *Nature* **380**, 215–221.
17
18 456 Meunier, J., 2024. The biology and social life of earwigs (Dermaptera). *Annu Rev Entomol* **69**,
19 457 259–276.
20
21 458 Nakagawa, S., Johnson, P.C.D., Schielzeth, H., 2017. The coefficient of determination r^2 and
22 459 intra-class correlation coefficient from generalized linear mixed-effects models revisited
23 460 and expanded. *J R Soc Interface* **14**, 20170213.
24
25
26 461 Nakagawa, S., Schielzeth, H., 2013. A general and simple method for obtaining R^2 from
27 462 generalized linear mixed-effects models. *Methods Ecol Evol* **4**, 133–142.
28
29 463 Nakagawa, S., Schielzeth, H., 2010. Repeatability for Gaussian and non-Gaussian data: A
30 464 practical guide for biologists. *Biol Rev Camb Philos Soc* **85**, 935–956.
31
32 465 Parker, G.A., 1982. Why are there so many tiny sperm? Sperm competition and the
33 466 maintenance of two sexes. *J Theor Biol* **96**, 281–294.
34
35 467 Peig, J., Green, A.J., 2010. The paradigm of body condition: A critical reappraisal of current
36 468 methods based on mass and length. *Funct Ecol* **24**, 1323–1332.
37
38 469 Peig, J., Green, A.J., 2009. New perspectives for estimating body condition from mass/length
39 470 data: The scaled mass index as an alternative method. *Oikos* **118**, 1883–1891.
40
41 471 Pol, M. van de, Wright, J., 2009. A simple method for distinguishing within- versus between-
42 472 subject effects using mixed models. *Anim Behav* **77**, 753–758.
43
44
45 473 R Core Team, 2025. R: A language and environment for statistical computing.
46
47 474 Reznick, D., Nunney, L., Tessier, A., 2000. Big houses, big cars, superfleas and the costs of
48 475 reproduction. *Trends Ecol Evol* **15**, 421–425.
49
50 476 Ricklefs, R.E., Wikelski, M., 2002. The physiology/life-history nexus. *Trends Ecol Evol* **17**,
51 477 462–468.
52
53
54
55
56
57
58
59
60

- 478 Roff, D., 1992. Evolution of Life Histories: Theory and Analysis. Chapman & Hall, Inc., New
479 York, NY.
- 480 Roff, D.A., Fairbairn, D.J., 2007. The evolution of trade-offs: Where are we? *J Evol Biol* **20**,
481 433–447.
- 482 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
483 Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K.,
484 Tomancak, P., Cardona, A., 2012. Fiji: An open-source platform for biological-image analysis.
485 *Nat Method* **9**, 676–682.
- 486 Simpson, E.H., 1951. The interpretation of interaction in contingency tables. *J. R. Stat. Soc.*
487 *Ser. B Stat. Methodol.* **13**, 238–241.
- 488 Smith, C.C., Fretwell, S.D., 1974. The optimal balance between size and number of offspring.
489 *Am Nat* **108**, 499–506.
- 490 Staerkle, M., Kölliker, M., 2008. Maternal food regurgitation to nymphs in earwigs (*Forficula*
491 *auricularia*). *Ethology* **114**, 844–850.
- 492 Stearns, S.C., 1992. The Evolution of Life Histories. Oxford University Press.
- 493 Stearns, S.C., 1989. Trade-offs in life-history evolution. *Funct Ecol* **3**, 259–268.
- 494 Stoffel, M.A., Nakagawa, S., Schielzeth, H., 2017. rptR: Repeatability estimation and variance
495 decomposition by generalized linear mixed-effects models. *Methods Ecol Evol* **8**, 1639–
496 1644.
- 497 Tourneur, J.-C., 2018. Factors affecting the egg-laying pattern of *Forficula auricularia*
498 (Dermaptera: Forficulidae) in three climatologically different zones of North America. *Can*
499 *Entomol* **150**, 511–519.
- 500 Tourneur, J.-C., Gingras, J., 1992. Egg laying in a northeastern North American (Montréal,
501 Québec) population of *Forficula auricularia* L. (Dermaptera: Forficulidae). *Can Entomol* **124**,
502 1055–1061.
- 503 Van Noordwijk, A.J., Jong, G. de, 1986. Acquisition and allocation of resources: Their
504 influence on variation in life history tactics. *Am Nat* **128**, 137–142.
- 505 Westneat, D.F., Araya-Ajoy, Y.G., Allegate, H., Class, B., Dingemanse, N., Dochtermann, N.A.,
506 Garamszegi, L.Z., Martin, J.G.A., Nakagawa, S., Réale, D., Schielzeth, H., 2020. Collision
507 between biological process and statistical analysis revealed by mean centring. *J Anim Ecol*
508 **89**, 2813–2824.
- 509 Wickham, H., 2016. ggplot2: Elegant graphics for data analysis. Springer, New York.

Winkler, D.W., Wallin, K., 1987. Offspring size and number: A life history model linking effort per offspring and total effort. *Am Nat* **129**, 708–720.

Zera, A.J., Harshman, L.G., 2001. The physiology of life history trade-offs in animals. *Annu Rev Ecol Syst* **32**, 95–126.

##Figures




Figure 1: Partitioning the egg size–number trade-off into within- and between-female components. (a) Within-female (plastic) relationship between egg number and mean egg size. Points represent individual broods (first clutch: open circles; second clutch: closed circles), with grey lines connecting repeated measures from the same female. The x-axis shows standardized within-female deviation in egg number (0 = female mean). The solid black line represents the Bayesian population-level estimate of the within-female slope ($\beta_W = -0.01$, 95% CrI (-0.06, 0.05)). (b) Between-female relationship (i.e., differences in mean allocation strategy) based on female means across broods plotted against standardized between-female deviation in egg number. The black line represents the Bayesian population-level between-female slope ($\beta_B = -0.05$, 95% CrI (-0.10, 0.00)). Together, the panels distinguish plastic allocation within females from strategic differences among females, indicating weak support for a reproductive size–number trade-off.






Figure 2: Clutch-specific variance components of egg size. (a) Between-female variance in egg size for first and second clutches, estimated from a hierarchical Bayesian model with clutch-specific female effects. (b) Within-clutch (egg-to-egg) variance for first and second clutches. Orange points represent posterior draws, black points indicate posterior medians, and vertical lines denote 95% credible intervals. Between-female variance did not differ clearly between clutches, whereas within-clutch variance was greater in second clutches.

516



Partitioning the egg size–number trade-off into within- and between-female components. (a) Within-female (plastic) relationship between egg number and mean egg size. Points represent individual broods (first clutch: open circles; second clutch: closed circles), with grey lines connecting repeated measures from the same female. The x-axis shows standardized within-female deviation in egg number (0 = female mean). The solid black line represents the Bayesian population-level estimate of the within-female slope ($\beta_W = -0.01$, 95% CrI (-0.06, 0.05)). (b) Between-female relationship (i.e., differences in mean allocation strategy) based on female means across broods plotted against standardized between-female deviation in egg number. The black line represents the Bayesian population-level between-female slope ($\beta_B = -0.05$, 95% CrI (-0.10, 0.00)). Together, the panels distinguish plastic allocation within females from strategic differences among females, indicating weak support for a reproductive size–number trade-off.

1146x1146mm (72 x 72 DPI)



Clutch-specific variance components of egg size. (a) Between-female variance in egg size for first and second clutches, estimated from a hierarchical Bayesian model with clutch-specific female effects. (b) Within-clutch (egg-to-egg) variance for first and second clutches. Grey points represent posterior draws, black points indicate posterior medians, and vertical lines denote 95% credible intervals. Between-female variance did not differ clearly between clutches, whereas within-clutch variance was greater in second clutches.

1146x1146mm (72 x 72 DPI)

**Supplemental Material: Disentangling within- and between-
female egg size–number trade-offs in the European earwig**

Clint D. Kelly^{1,✉}

¹ Département des Sciences biologiques, Université du Québec à Montréal, Montréal, Canada

✉ Correspondence: [Clint D. Kelly <kelly.clint@uqam.ca>](mailto:kelly.clint@uqam.ca)

For Review Only

1. Within–between multivariate model

To partition covariance arising from within-female plasticity from covariance attributable to consistent among-female differences, I applied the within–between decomposition of van de Pol and Wright (Pol and Wright, 2009). Although mean-centering can impose assumptions about underlying biological processes (Westneat et al., 2020), these assumptions are appropriate here because the aim was to distinguish plastic allocation within females from persistent differences among females.

Let egg number for female i in brood j be denoted N_{ij} . Egg number was decomposed as

$$N_{ij} = \bar{N}_i + (N_{ij} - \bar{N}_i) \quad (1)$$

The between- and within-female components were defined as

$$N_i^B = \bar{N}_i - \bar{N} \quad (2)$$

$$N_{ij}^W = N_{ij} - \bar{N}_i \quad (3)$$

Egg number submodel

Egg number was modelled using a negative binomial distribution with a log link:

$$N_{ij} \sim \text{NegBinomial}(\mu_{ij}, \theta) \quad (4)$$

$$\log(\mu_{ij}) = \beta_0^N + \beta_1^N \text{Brood}_{ij} + u_{0i}^N \quad (5)$$

where

$$u_{0i}^N \sim \mathcal{N}(0, \sigma_N^2) \quad (6)$$

Mean egg size submodel

Mean egg size was modelled as

$$S_{ij} \sim \mathcal{N}(\mu_{ij}^S, \sigma_S^2) \tag{7}$$

$$\mu_{ij}^S = \beta_0^S + \beta_1^S \text{Brood}_{ij} + \beta_W N_{ij}^W + \beta_B N_i^B + u_{0i}^S \tag{8}$$

with

$$u_{0i}^S \sim \mathcal{N}(0, \sigma_{S,id}^2) \tag{9}$$

Multivariate structure

The two submodels were fit jointly:

$$\begin{pmatrix} u_{0i}^N \\ u_{0i}^S \end{pmatrix} \sim \mathcal{MVN} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \Sigma \right) \tag{10}$$

where

$$\Sigma = \begin{pmatrix} \sigma_N^2 & \sigma_{NS} \\ \sigma_{NS} & \sigma_{S,id}^2 \end{pmatrix} \tag{11}$$

The intercept correlation is

$$\rho_{NS} = \frac{\sigma_{NS}}{\sqrt{\sigma_N^2 \sigma_{S,id}^2}} \tag{12}$$

Relationship between model parameters and total phenotypic covariance

Total phenotypic covariance can be partitioned as

$$\text{Cov}_{\text{total}} = \text{Cov}_{\text{within}} + \text{Cov}_{\text{among}} \tag{13}$$

Within-female covariance

$$\text{Cov}_{\text{within}} = \beta_W \text{Var}(N_{ij}^W) \quad (14)$$

Among-female covariance

$$\text{Cov}_{\text{among, slope}} = \beta_B \text{Var}(N_i^B) \quad (15)$$

$$\text{Cov}_{\text{among, intercept}} = \sigma_{NS} \quad (16)$$

Total among-female covariance is therefore

$$\text{Cov}_{\text{among}} = \beta_B \text{Var}(N_i^B) + \sigma_{NS} \quad (17)$$

Total phenotypic covariance

Combining components:

$$\text{Cov}_{\text{total}} = \beta_W \text{Var}(N_{ij}^W) + \beta_B \text{Var}(N_i^B) + \sigma_{NS} \quad (18)$$

The intercept correlation is

$$\rho_{NS} = \frac{\sigma_{NS}}{\sqrt{\sigma_N^2 \sigma_{S,id}^2}} \quad (19)$$

Corresponding *brms* Specification

```
bf_number <- bf(
  egg.number ~ brood + (1 |p| id),
  family = negbinomial()
)

bf_size <- bf(
  mean.egg.size ~ brood +
    egg_number_within_z +
    egg_number_between_z +
    (1 |p| id),
  family = gaussian()
)
```

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

```
mv_model <- bf_number + bf_size
```

For Review Only

Table S 1: Posterior summaries from the multivariate Bayesian mixed-effects model examining the relationship between egg number and mean egg size. The model jointly estimated a negative binomial model for egg number and a Gaussian model for egg size, including brood effects and partitioning egg number into within- and between-female components. Reported values are posterior medians and 95% credible intervals (CrI). Random effects represent among-female (ID-level) variation, and the correlation term describes the association between female intercepts for egg number and egg size. The dispersion parameter refers to the negative binomial shape parameter.

Parameter	Estimate	Lower 95% CrI	Upper 95% CrI
Egg number: Brood 2 – Brood 1 (log scale)	-0.413	-0.635	-0.189
Egg number: Intercept (Brood 1, log scale)	3.940	3.782	4.107
Egg size: Brood 2 – Brood 1	-0.133	-0.244	-0.024
Egg size: Between-female deviation in egg number (z)	-0.050	-0.104	0.002
Egg size: Within-female deviation in egg number (z)	-0.007	-0.061	0.049
Egg size: Intercept (Brood 1)	3.543	3.468	3.618
cor_id__eggnumber_Intercept__meaneggsize_Intercept	-0.017	-0.813	0.816
SD (Egg number Female ID)	0.078	0.003	0.270
SD (Egg size Female ID)	0.082	0.006	0.155
Residual SD (Egg size)	0.191	0.155	0.235
Dispersion (Egg number NegBin shape)	4.080	2.835	5.853

2. Clutch-Specific Hierarchical Model of Egg Size

Statistical Model

Egg perimeter was standardized prior to analysis (mean = 0, SD = 1).

For female i , egg j , and brood $b \in \{1,2\}$, standardized egg size y_{ijb} was modeled as:

$$y_{ijb} \sim \mathcal{N}(\mu_{ijb}, \sigma_b^2)$$

The conditional mean structure was:

$$\mu_{ijb} = \beta_0 + \beta_1 I(b = 2) + u_{ib}$$

where:

- β_0 is the mean egg size in clutch 1 (reference level),
- β_1 is the fixed difference between clutch 2 and clutch 1,
- u_{ib} is the female-specific deviation for clutch b .

Random Effects Structure

Clutch-specific female deviations were modeled as a correlated bivariate normal vector:

$$\mathbf{u}_i = \begin{pmatrix} u_{i1} \\ u_{i2} \end{pmatrix} \sim \mathcal{MVN} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \Sigma \right)$$

with variance–covariance matrix:

$$\Sigma = \begin{pmatrix} \tau_1^2 & \rho\tau_1\tau_2 \\ \rho\tau_1\tau_2 & \tau_2^2 \end{pmatrix}$$

Here:

- τ_1^2 and τ_2^2 are the among-female variances in clutches 1 and 2, respectively,
- $\rho = \text{Corr}(u_{i1}, u_{i2})$ is the correlation between female deviations across clutches.

Importantly, no clutch-independent random intercept was estimated. The model includes only clutch-specific female deviations (i.e., random slopes of clutch within female), corresponding to the specification `(0 + brood_factor | id)` in *brms*.

Residual Variance Model

Residual variance was allowed to differ between clutches:

$$\log(\sigma_b) = \alpha_0 + \alpha_1 I(b = 2)$$

where:

- σ_1^2 is the residual variance in clutch 1,
- σ_2^2 is the residual variance in clutch 2.

The log link ensures that residual standard deviations remain positive.

Corresponding *brms* Specification

```
bf(
  scale(mean.perim) ~ brood_factor + (0 + brood_factor | id),
  sigma ~ brood_factor
)
```

Table S 2: Posterior medians and 95% credible intervals from a Bayesian mixed model of scaled egg perimeter, including clutch-specific means, clutch-specific female-level random effects, and clutch-specific residual variance.


Parameter	Estimate	Lower 95% CrI	Upper 95% CrI
Brood 2 – Brood 1	-0.535	-0.827	-0.246
Mean (Brood 1)	0.283	0.050	0.530
Correlation (Brood 1, Brood 2 Female ID)	0.250	-0.025	0.488
SD (Brood 1 Female ID)	0.855	0.704	1.058
SD (Brood 2 Female ID)	0.784	0.638	0.982
Residual SD (Brood 1)	0.457	0.444	0.471
Residual SD (Brood 2)	0.594	0.574	0.616

References

Pol, M. van de, Wright, J., 2009. A simple method for distinguishing within- versus between-subject effects using mixed models. *Anim Behav* **77**, 753–758.

Westneat, D.F., Araya-Ajoy, Y.G., Allegue, H., Class, B., Dingemanse, N., Dochtermann, N.A., Garamszegi, L.Z., Martin, J.G.A., Nakagawa, S., Réale, D., Schielzeth, H., 2020. Collision between biological process and statistical analysis revealed by mean centring. *J Anim Ecol* **89**, 2813–2824.

For Review Only



Two-panel conceptual diagram comparing expected versus observed relationships between egg number and egg size. In the left panel (Expected), a positive among-female correlation is shown (red ellipse), with numbered points indicating first and second clutches and a within-female trade-off implied. In the right panel (Observed), a negative among-female relationship is shown (blue ellipse), indicating a trade-off between egg number and size across females, but no within-female trade-off; second clutches are depicted as having fewer and slightly smaller eggs.

268x155mm (300 x 300 DPI)

Disentangling within- and between-female egg size–number trade-offs in the European earwig.

Clint D. Kelly^{1,✉}

¹ Département des Sciences biologiques, Université du Québec à Montréal, Montréal, Canada

✉ Correspondence: [Clint D. Kelly <kelly.clint@uqam.ca>](mailto:kelly.clint@uqam.ca)

For Review Only

Abstract

Life-history theory predicts a trade-off between offspring size and number, yet empirical studies often fail to detect the expected negative association. This discrepancy may occur because phenotypic correlations confuse variation in resource acquisition with variation in allocation strategy. Using repeated measures of female European earwigs (*Forficula auricularia*), I partitioned covariance between egg size and egg number into among- and within-female components using Bayesian mixed models. The raw phenotypic correlation was near zero. However, females that produced more eggs on average laid smaller eggs, revealing a modest negative association among females that aligns with an allocation trade-off. In contrast, there was no detectable within-female covariance once clutch order was accounted for. Egg number repeatability was effectively zero, suggesting that clutch size varies mainly within rather than among females, while egg size was highly repeatable. Both egg number and egg size decreased in second clutches, and within-clutch egg-size variance increased, indicating a reproductive constraint (i.e., fewer functional ovarioles) under a possibly diminished resource state. Together, these findings demonstrate that the size–number trade-off appears to be expressed among females through consistent differences in allocation strategy, whereas clutch size itself remains flexible. By separating hierarchical sources of variation, this study shows that allocation trade-offs can remain hidden at the phenotypic level and that evolutionary constraints act through stable allocation rules rather than through flexible clutch-size adjustments.

Introduction

Life-history theory predicts that organisms must allocate finite resources among competing fitness components. Because energy and time are limited, investment in one trait constrains investment in others (Roff, 1992; Stearns, 1992). A central expression of this principle is the offspring size–number trade-off: for a given reproductive budget, producing larger offspring reduces the number that can be produced (Parker, 1982; Smith and Fretwell, 1974). Since larger eggs often enhance offspring performance and survival (Clutton-Brock, 1991), selection is expected to favour an optimal balance between egg size and egg number that maximises maternal fitness (Fox and Czesak, 2000). Yet empirical tests of the size–number trade-off often fail to reveal the expected negative relationship. Across individuals within populations, egg size and egg number are often positively correlated (Bernardo, 1996; Fox and Czesak, 2000; Hargitai et al., 2005; Roff, 1992). This apparent contradiction arises because phenotypic correlations conflate two distinct biological processes: variation in resource acquisition and variation in allocation strategy. Van Noordwijk and De Jong (1986) showed that when variance in acquisition exceeds variance in allocation, individuals with greater resource pools can produce both more and larger eggs, generating positive among-individual correlations even if all individuals are constrained by the same underlying allocation trade-off. In such cases, the population-level pattern masks allocation constraints operating within individuals—an outcome conceptually analogous to Simpson’s paradox (Simpson, 1951).

From this perspective, the critical prediction of life-history theory is not simply a negative phenotypic correlation, but a negative within-individual covariance between egg size and egg number once among-individual differences in acquisition are accounted for (Careau and Wilson,

1
2
3
4 49 2017). Detecting such a masked trade-off, therefore, requires partitioning phenotypic variance
5
6 50 and covariance into hierarchical components. Repeated measures of individuals allow estimation
7
8 51 of (i) among-individual variance, reflecting consistent differences among females in total
9
10 52 reproductive investment, and (ii) within-individual variance, reflecting how females adjust
11
12 53 allocation across reproductive bouts. Careau and Wilson (2017) argue that only by decomposing
13
14 54 covariance into these components using multivariate mixed models can we determine whether
15
16 55 allocation trade-offs operate within individuals despite positive population-level associations.
17
18
19
20 56 Crucially, this framework also predicts that traits involved in allocation trade-offs should show
21
22 57 substantial repeatability. If females differ consistently in resource acquisition, egg size and egg
23
24 58 number should exhibit among-individual variance across clutches. High repeatability would,
25
26 59 therefore, indicate persistent differences in total reproductive investment among females,
27
28
29 60 consistent with variation in acquisition. The marked increase in within-clutch egg size variance in
30
31 61 second broods further supports a resource constraint interpretation. Under reduced resource
32
33 62 availability, females may be less able to maintain uniform provisioning across all eggs, leading to
34
35 63 greater heterogeneity in egg size within a clutch. Such increased variance is consistent with
36
37 64 diminished allocation precision when reproductive resources are limited, rather than an adaptive
38
39 65 diversification of offspring size.
40
41
42
43
44 66 Despite the foundational importance of the size–number trade-off, empirical tests rarely partition
45
46 67 covariance into within- and among-individual components to distinguish allocation from
47
48 68 acquisition effects. Here, Here, I address this gap using the European earwig, *Forficula*
49
50
51 69 *auricularia*, a species that allows repeated measures of reproductive investment within females
52
53 70 but whose in which previous work has reported no clear or consistent pattern ~~of allocation to in the~~
54
55
56
57
58
59
60

1
2
3
4 71 association between egg size and egg number ~~and size is unclear~~ (Koch and Meunier, 2014).
5
6 72 highlighting the difficulty of interpreting phenotypic correlations without partitioning
7
8 73 hierarchical sources of variation. Female earwigs lay a clutch within a short temporal window in
9
10 74 a subterranean burrow (Meunier, 2024) and provide extended maternal care (Meunier, 2024;
11
12 75 Staerke and Kölliker, 2008). In ~~Quebee~~Québec populations, environmental constraints typically
13
14 76 render females effectively semelparous (Tourneur, 2018), but under favourable laboratory
15
16 77 conditions, they can produce a second clutch. This system, therefore, enables direct estimation of
17
18 78 within-female covariance across successive reproductive bouts. I test three predictions derived
19
20 79 from acquisition–allocation theory (Van Noordwijk and Jong, 1986; see also Reznick, Nunney,
21
22 80 and Tessier, 2000). First, if variance in resource acquisition among females exceeds variance in
23
24 81 allocation strategy, then egg size and egg number should exhibit substantial repeatability and a
25
26 82 positive among-individual correlation, whereas the within-individual correlation should be
27
28 83 negative, revealing a masked allocation trade-off. Second, if maternal fitness is maximised at a
29
30 84 particular offspring size or number (Smith and Fretwell, 1974), then one trait should remain
31
32 85 relatively invariant across clutches while the other adjusts between reproductive bouts (Winkler
33
34 86 and Wallin, 1987). Third, if reproductive investment is structured differently across bouts, then
35
36 87 variance in egg size may differ between first and second clutches at both the among-female and
37
38 88 within-clutch levels (e.g. Hargitai et al., 2005), providing further insight into how selection
39
40 89 shapes the distribution of maternal investment across reproductive events.
41
42
43
44
45
46
47
48 90 By separating among-female differences in total reproductive investment from within-female
49
50 91 allocation decisions, this study tests a fundamental question in life-history theory: do apparent
51
52 92 positive associations between egg size and number reflect relaxed constraints, or do allocation
53
54
55
56
57
58
59
60

trade-offs persist but remain hidden within individuals? Demonstrating a negative within-female covariance despite positive population-level patterns would show that acquisition can mask, but not erase, the constraints imposed by finite resources. In this way, my study evaluates not only the existence of a trade-off but also the hierarchical structure through which life-history constraints are expressed.

Methods

I collected adult female earwigs (n=42) in September and October 2024 in an urban park (Parc Outremont) in Montréal, Canada. All earwigs were brought into the laboratory, assigned a unique ID, and individually placed in a 90-mm-diameter Petri dish containing a thin layer of moist sand. Females were provided a cotton-plugged water vial and fed TetraMin Tropical fish flakes ad libitum. Females were maintained in an incubator (Percival Scientific, Iowa, USA) at 10°C in constant darkness.

Females were checked daily for oviposition. When a female laid eggs, I recorded the date and collected all eggs from the sand, and placed them on filter paper in a new Petri dish. The eggs were then photographed under a Leica S6D stereo microscope (Leica Microsystems Inc., Concord, ON, Canada) at 0.68x magnification using Enersight software (Leica Microsystems Inc., Concord, ON, Canada), which automatically adds a scale bar to the photo. Eggs were then discarded.

To stimulate females to lay a second clutch of eggs, I recreated a brief spring period ~~of spring~~ where during which they were allowed to accumulate resources for egg production and replenish their sperm stores. To achieve this, females that had recently oviposited were placed in a new

1
2
3
4 114 Petri dish containing a cotton-plugged water vial, TetraMin Tropical fish flakes, and a shelter
5
6 115 consisting of four 4 cm-long straws (Blowholes, Markham, Canada) glued together. These Petri
7
8 116 dishes were placed in a 23 °C incubator with a 12L:12D light: dark cycle. Females were
9
10 117 maintained under these conditions for two weeks, at which time a haphazardly chosen stock male
11
12 118 from our colony was placed with her. The pair was kept together for a week, after which a new
13
14 119 stock male replaced the original ~~male~~, and ~~that~~ the new pair was kept together for another week.
15
16
17 120 Females were then transferred to the 10 °C incubator and treated as previously until they laid
18
19 121 their second clutch, at which point their eggs were photographed. Females were euthanised by
20
21 122 freezing after laying their second clutch, and their pronotum was photographed as described
22
23 123 above. Because female fecundity is often related to body size in insects (Honek, 1993) and could
24
25 124 thus possibly explain a positive among-female covariance between egg number and size, I
26
27 125 measured two proxies of size, pronotum length and body mass. Females were weighed to the
28
29 126 nearest 0.01 mg using a Sartorius Secura 224-1S analytical balance. Females were weighed three
30
31 127 times, with each weighing bout separated by four days. I used Fiji image analysis software
32
33 128 (Schindelin et al., 2012) to measure the length of the female pronotum to the nearest 0.001 mm
34
35 129 three times, with each measurement separated by 4 days. I used the same procedure to measure
36
37 130 the perimeter of each egg.
38
39
40
41
42
43

44 131 **Statistical analysis**

45 132 I calculated the repeatability of the body mass, pronotum length and egg perimeter measurements
46
47 133 using the R package *rptR* (Stoffel, Nakagawa, and Schielzeth, 2017). I entered the trait of interest
48
49 134 as the response variable, female ID as a random effect, and assumed a Gaussian distribution.
50
51 135 Each analysis was based on 1000 bootstrap samples. The measurements of my two
52
53
54
55
56
57
58
59
60

1
2
3
4 136 morphological traits were highly repeatable (pronotum length: R [95% CI] = 0.98 [0.97, 0.99];
5
6 137 body mass: R [95% CI] = 0.99 [0.988, 0.996]); therefore, I used each female's mean value for
7
8 138 each trait where necessary. Because egg perimeter was highly repeatable in both clutches (first
9
10 139 clutch: R [95% CI] = 0.977 [0.976, 0.979]; second clutch: R [95% CI] = 0.93 [0.926, 0.937]), I
11
12 140 calculated each egg's average perimeter and then used these values to calculate the mean egg size
13
14 141 for each clutch for each female.
15
16
17
18 142 To reduce the two correlated measures of body size (pronotum length and body mass) to a single
19
20 143 structural size metric, I performed a principal components analysis (PCA) on the correlation
21
22 144 matrix of standardized variables using the *prcomp* function in the R stats package (R Core Team,
23
24 145 2025). The first principal component (PC1) explained 60.8% of the total variance and was used
25
26 146 as an index of structural body size. In addition, female body condition was estimated using the
27
28 147 scaled mass index (SMI) following Peig and Green (2010, 2009; see also Kelly, Tawes, and
29
30 148 Worthington, 2014), which standardizes body mass relative to structural size (pronotum length).
31
32
33
34
35 149 I first quantified associations among reproductive traits using raw phenotypic correlations.
36
37 150 Pearson correlation coefficients were calculated using individual-level trait values, thereby
38
39 151 capturing total phenotypic covariation across all sources of variation. These correlations reflect
40
41 152 the combined contribution of consistent differences among females, variation among clutches
42
43 153 within females, and residual effects. Consequently, raw phenotypic correlations provide a
44
45 154 descriptive estimate of overall trait associations but do not distinguish between covariation
46
47 155 arising from stable female differences and that arising from within-female (i.e., among-clutch)
48
49
50
51 156 processes.
52
53
54
55
56
57
58
59
60

To distinguish covariance arising from within-female plasticity from covariance attributable to consistent among-female differences, I followed the within-between decomposition approach of van de Pol and Wright [(2009); see Supplementary Material for details]. This decomposition is necessary because residual covariance does not isolate within-individual effects and, in models with different response distributions, does not provide a directly interpretable estimate of within-female trade-offs. Although mean-centering has been shown to impose assumptions about underlying biological processes (Westneat et al., 2020), these assumptions are appropriate here because the biological question concerns how reproductive allocation varies within females across clutches versus how females differ consistently in allocation strategy. Egg number was decomposed into two components: a between-female component, defined as each female's mean egg number across broods centered on the population mean, and a within-female component, defined as the deviation of each brood from that female's own mean. Hereafter, we refer to "among-female" variation as differences between individuals and "within-female" variation as differences among clutches within individuals. These two predictors were included simultaneously in a Bayesian multivariate mixed-effects model in which egg number and mean egg size were modelled jointly. Egg number was modelled using a negative binomial distribution with a log link to account for overdispersion, whereas mean egg size was modelled assuming a Gaussian distribution with an identity link. Clutch (first vs. second clutch) was included as a fixed effect in both submodels. Female identity was included as a random intercept for both response variables. This structure allowed estimation of among-female variance in egg number, among-female variance in egg size, and the covariance (correlation) between female-level intercepts for the two traits.

1
2
3
4 179 This multivariate framework partitions total phenotypic covariance into components arising from
5
6 180 consistent differences among females and from within-female deviations across clutches.
7
8 181 Covariance attributable to consistent differences among females is captured by the female-level
9
10 182 random effects, whereas covariance arising from within-female variation is captured by the fixed
11
12 183 within-female component of egg number. The fixed between-female component estimates how
13
14
15 184 mean egg size varies with mean egg number across females, ~~and thus captures~~ thereby capturing
16
17 185 systematic (i.e., explained) differences in allocation strategy among females. In contrast, the fixed
18
19 186 within-female component tests whether egg size changes plastically when a female produces
20
21 187 more or fewer eggs than her own mean. The correlation between female-level intercepts
22
23 188 represents residual (latent) covariance among baseline trait values that is not explained by the
24
25 189 fixed effects, including the between-female allocation relationship. Thus, the between-female
26
27 190 slope tests whether females diverge in their mean allocation strategy (i.e., whether females that
28
29 191 consistently produce more eggs also produce smaller eggs), whereas the intercept correlation
30
31 192 quantifies any remaining covariance between egg number and egg size after accounting for these
32
33 193 effects. Continuous predictors were standardized prior to analysis, such that standardized
34
35 194 coefficients represent the expected change in egg size (in standard deviation units) associated
36
37 195 with a one standard deviation change in egg number at the relevant hierarchical level.
38
39
40
41
42
43 196 To quantify the structure of egg-size variation specifically, I fitted a univariate mixed-effects
44
45 197 model in which egg size (standardised mean egg perimeter) was modelled as a Gaussian response
46
47 198 with female identity included as a random intercept (see Supplementary Material for details).
48
49
50 199 This formulation partitions total phenotypic variance in egg size into variance attributable to
51
52 200 differences among females and variance occurring within females among eggs within clutches.
53
54
55
56
57
58
59
60

The among-female variance component reflects consistent maternal differences in egg size allocation, whereas the within-female component captures heterogeneity in provisioning within clutches and residual variation.

Repeatability was calculated on the observation scale for both traits. For egg size (Gaussian model):

$$R_{\text{egg size}} = \frac{V_F}{V_F + V_R}$$

where V_F is the among-female variance (i.e., variance of the female random intercept) and V_R is the residual variance. Because the Gaussian model is parameterized directly on the observation scale, this formulation corresponds to both latent- and observation-scale repeatability.

For egg number, which was modelled using a negative binomial distribution with a log link, the variance depends on both the expected mean and the dispersion (shape) parameter. Under the negative binomial parameterization used in *brms*, the observation-level (distribution-specific) variance is given by:

$$\text{Var}(Y) = \mu + \frac{\mu^2}{\theta}$$

where μ is the expected mean egg number and θ is the dispersion (shape) parameter. Because the model includes a log link, the among-female variance estimated from the random intercept is expressed on the latent (log) scale and must be transformed to the observation scale before calculating repeatability. Following Nakagawa et al. (2017), the among-female variance on the observation scale was therefore calculated as:

220
$$V_F^{\text{obs}} = \mu^2(\exp(\sigma^2) - 1)$$

221 where σ^2 is the variance of the random intercept on the latent scale.

222 Observation-scale repeatability for egg number was then calculated as:

223
$$R_{\text{egg number}} = \frac{V_F^{\text{obs}}}{V_F^{\text{obs}} + \left(\mu + \frac{\mu^2}{\theta}\right)}$$

224 Both the transformed among-female variance and the distribution-specific variance were
225 computed for each posterior draw using the corresponding posterior samples of μ , σ^2 , and θ . To
226 account for variation in expected values across observations, these quantities were first calculated
227 at the observation level and then averaged within each posterior draw. This procedure yielded a
228 posterior distribution of repeatability that fully propagates uncertainty from all variance
229 components.

230 Repeatability for non-Gaussian traits was calculated following the framework of Nakagawa and
231 Schielzeth (2010) and its extensions for generalized linear mixed models (Nakagawa, Johnson,
232 and Schielzeth, 2017). Because distribution-specific variance can contribute substantially to total
233 phenotypic variance in negative binomial models, repeatability estimates for count data are often
234 lower than those obtained under Gaussian assumptions, even when among-individual variance is
235 present (Nakagawa, Johnson, and Schielzeth, 2017; Nakagawa and Schielzeth, 2010). By
236 calculating repeatability on the observation scale, these estimates represent the proportion of total
237 phenotypic variance in observed egg number attributable to consistent differences among
238 females.

I used weakly informative priors to constrain parameters to biologically realistic ranges while allowing the data to drive inference. Regression coefficients were assigned normal priors centered at zero (Normal(0, 0.5–1)), reflecting the expectation that clutch effects on egg traits would be moderate in magnitude. Intercepts were given normal priors scaled to each trait (egg number: Normal(0, 5); mean egg size: Normal(3.5, 1); standardized egg size: Normal(0, 0.5)). Among-female variation was assigned exponential priors (Exponential(1–2)), which favour small but non-zero individual differences. Residual variance (Gaussian models) and the negative binomial dispersion parameter were given Exponential(1) priors. Correlations among female-level random effects were regularized using an LKJ(2) prior to avoid overestimating trait associations. Posterior distributions were obtained using six Markov chains each with 4,000 iterations ~~per chain~~, including 1,000 warm-up iterations, and convergence was assessed using $\hat{R}$ statistics, effective sample sizes, and trace-plot inspection. Posterior distributions are summarised as posterior means with 95% credible intervals. Data were visualized using *ggplot* (Wickham, 2016) and *tidybayes* (Kay, 2020).

Results

Between- and within-female associations

Across all broods, the raw phenotypic correlation between egg number and egg size was weak and near zero ($r_{raw} = -0.04$, 95% CI [-0.25, 0.19]), indicating no detectable population-level association when hierarchical structure was ignored.

Egg number (clutch one: $\bar{x} = 51.55 \pm 2.49$ eggs; clutch two: $\bar{x} = 34.14 \pm 2.65$ eggs) and egg size (first clutch: $\bar{x} = 3.54 \pm 0.03$ mm; second clutch: $\bar{x} \pm SE = 3.41 \pm 0.03$ mm) both declined in

1
2
3
4 260 second clutches. Egg number decreased on the log scale ($\beta = -0.41$, 95% CrI (-0.64, -0.19;
5
6 261 [Figure 1](#); Supplementary Table 1), corresponding to an approximate 34% reduction on the natural
7
8 262 scale ($e^{-0.413} \approx 0.66$). Mean egg size also declined in the second clutch ($\beta = -0.13$, 95% CrI (-
9
10 263 0.24, -0.02)).
11
12
13 264 The results did not support my prediction of a within-female plastic trade-off between egg
14
15 265 number and egg size ([Figure 1 a](#)). The within-female deviation in egg number had a slope near
16
17 266 zero ($\beta_W = -0.01$, 95% CrI (-0.06, 0.05), indicating that females did not systematically adjust egg
18
19 267 size when producing more or fewer eggs relative to their own mean. By comparison, the among-
20
21 268 female association (captured by the fixed between-female slope) revealed a negative relationship
22
23 269 between egg size and mean female egg number ($\beta_B = -0.05$, 95% CrI (-0.10, 0.00; [Figure 1 b](#)).
24
25 270 The posterior distribution of the among-female slope was centered on negative values, suggesting
26
27 271 weak evidence for divergence in mean allocation strategy among females. The correlation
28
29 272 between female-level intercepts for egg number and egg size was highly uncertain (median = -
30
31 273 0.02, 95% CrI: -0.81, 0.82), indicating limited information for estimating latent covariance with
32
33 274 only two broods per female. With only two repeated measures per female, intercept correlations
34
35 275 are weakly identifiable and should be interpreted cautiously.
36
37
38
39
40
41 276 No measure of female size/condition was positively correlated with the number of eggs in the
42
43 277 first brood (pronotum length: $r = 0.14$, 95% CI [-0.19, 0.43]; body mass: $r = 0.12$, 95% CI [-0.21,
44
45 278 0.43]; PC1: $r = -0.17$, 95% CI [-0.44, 0.15]; scaled mass index: $r_{raw} = -0.02$, 95% CI [-0.35,
46
47 279 0.33]) providing no evidence that larger females or those in better condition produce more eggs. I
48
49 280 therefore did not include a measure of size/condition in this model. Together, these results show
50
51 281 that the size–number trade-off is not apparent at the raw phenotypic level and is not expressed as
52
53
54
55
56
57
58
59
60

within-female plasticity, but emerges as a modest negative association at the among-female level through differences in average allocation strategy.

Egg size variation between females and within clutches

Egg size exhibited substantial variation at both the between-female and within-clutch levels (Figure 2; Supplementary Table 2). Between-female variance was similar across reproductive events, with a median of 0.73 (95% CrI: 0.50, 1.12) in first clutches and 0.61 (95% CrI: 0.41, 0.96) in second clutches. The posterior median difference (second – first) was -0.11 (95% CrI: -0.56, 0.28), and the posterior probability that between-female variance was greater in second clutches was 0.28, indicating no clear evidence of a difference. In contrast, within-clutch (egg-to-egg) variance was substantially higher in second clutches (median = 0.35 (95% CrI: 0.33, 0.38) than in first clutches (median = 0.21 (95% CrI: 0.20, 0.22). The posterior median difference was 0.14 (95% CrI: 0.12, 0.17), with a posterior probability of 1.00, that within-clutch variance was greater in second clutches.

Egg number and egg size repeatability

Repeatability was calculated from posterior samples of the variance components on the observation scale. Egg number exhibited extremely low repeatability. Although there was some among-female variance in egg number (SD = 0.078, 95% CrI [0.003, 0.270]; Supplementary Table 1), the overwhelming majority of variance arose from distribution-specific (negative binomial) variance. Consequently, observation-scale repeatability was near zero ($R = 4.86 \times 10^{-2}$, 95% CrI-[3.80×10^{-5} , 2.43×10^{-1}]), indicating that very little of the total phenotypic variation in clutch size was attributable to consistent differences among females.

~~In contrast, egg size showed moderate~~Because variance components and repeatability ~~on the~~
~~observation scale.~~were derived from different models, these quantities are reported separately.
Among-female variance in egg size from the multivariate model was modest in absolute terms
(SD = 0.082, 95% CrI [0.006, 0.155]; ~~Supplementary Table 1~~) ~~was substantial relative to residual~~
~~variance~~ (SD = 0.191, 95% CrI [0.155, 0.235]); ~~Supplementary Table 1~~), providing a sense of
the trait's physical scale (mm). Repeatability, however, was estimated from the clutch-specific
univariate model (Supplementary Table 2), in which among-female variance exceeded residual
variance (SD = 0.855 vs 0.457 in first broods; 0.784 vs 0.594 in second broods), yielding $R =$
0.78 (95% CrI: 0.70, 0.84) in first broods and $R = 0.63$ (95% CrI: 0.53, 0.73) in second broods.
~~These results indicate that females differed consistently in their mean egg size across broods,~~
~~whereas clutch size was largely context dependent.~~

Discussion

Life-history theory predicts a trade-off between offspring size and number because both traits
draw from a finite pool of reproductive resources (Lack, 1947; Roff, 1992; Smith and Fretwell,
1974; Stearns, 1992). Yet, as emphasised by van Noordwijk and de Jong (1992), phenotypic
correlations need not directly reveal allocation trade-offs, because variation in resource
acquisition can obscure underlying allocation rules (Careau and Wilson, 2017; Johnson and
Nassrullah, 2024). This acquisition–allocation framework has become central to interpreting life-
history covariance (Houle, 1991; Reznick, Nunney, and Tessier, 2000). By partitioning
covariance into within- and among-female components, my results show that the size–number
trade-off in this population of earwigs is ~~at most weakly~~modestly expressed among females

~~and~~but absent within them, illustrating how hierarchical structure shapes the detectability of life-history trade-offs (Careau and Wilson, 2017; Dingemanse and Dochtermann, 2013; Pol and Wright, 2009; Westneat et al., 2020).

At the phenotypic level, egg size and egg number were uncorrelated in my earwig population, consistent with Koch and Meunier (2014). Under the framework of van Noordwijk and de Jong (1986), such phenotypic independence is expected when variation in resource acquisition obscures underlying allocation trade-offs. After hierarchical decomposition, however, a weak negative association emerged at the among-female level: females that produced more eggs on average tended to lay slightly smaller eggs. This pattern is captured by the fixed between-female slope and reflects ~~systematic~~ differences among females in their mean allocation strategy. In contrast, the correlation between female-level intercepts—representing residual (latent) covariance after accounting for this allocation relationship—was highly uncertain and overlapped zero, indicating little evidence for additional integration between traits beyond that explained by differences in mean allocation strategy (Roff and Fairbairn, 2007; Stearns, 1989).

~~Crucially, these results must be interpreted in light of the variance structure of egg number. Egg number exhibited near-zero repeatability, indicating that there is very little residual among-female variation in clutch size once distribution-level variance is taken into account. At first glance, this may appear inconsistent with the presence of an among-female association. However, repeatability and the between-female slope quantify different components of variation. Repeatability reflects the proportion of unexplained (random-effect) variance attributable to consistent differences among females, whereas the between-female slope represents a systematic (fixed-effect) relationship among female means. Thus, even when residual among-female~~

~~variance is very small, it is still possible to detect a weak association among female means. In this case, the near-zero repeatability indicates that clutch size does not vary consistently among females, while the between-female slope suggests only minor differences in average reproductive output that covary weakly with egg size. Taken together, these results indicate that the apparent among-female trade-off is weak and arises from small differences in female means rather than from substantial or repeatable variation in clutch size. Accordingly, there is limited evidence for stable divergence in allocation strategy among females.~~

In contrast, there was no evidence for a within-female plastic trade-off. Although each female contributed only two broods, the posterior distribution of the within-female slope was tightly centered on zero, suggesting that any within-female effect, if present, is small. Females did not reduce egg size when producing more eggs than their own average; instead, egg number and egg size shifted together across broods. This pattern is consistent with state-dependent life-history models in which allocation rules remain relatively stable while total resource state fluctuates across reproductive bouts (Houston and McNamara, 1999; McNamara and Houston, 1996). Rather than dynamically reallocating resources between offspring size and number, females appear to exhibit clutch-level shifts driven by overall energetic state. ~~Together, these results suggest that clutch size is primarily state-dependent and influenced by transient physiological or structural constraints, whereas egg size reflects a more stable component of reproductive allocation.~~

Notably, egg number was not correlated with female structural size or body condition (i.e. scaled mass), indicating that persistent differences in morphology do not explain clutch-size variation. If stable acquisition differences were driving among-female variation, we would expect egg number

to show repeatability and to covary with these morphological size/condition indicators. The absence of such patterns weakens the interpretation that persistent acquisition differences—at least as reflected by morphology—underlie among-female differences in reproductive output. Instead, clutch size appears predominantly state-sensitive and influenced by clutch-level constraints.

This interpretation also helps explain the pronounced decline in second clutches. That second clutches comprised both fewer and smaller eggs is consistent with previous work in North American and European populations (Koch and Meunier, 2014; Tourneur and Gingras, 1992). Tourneur and Gingras (1992) proposed that some ovarioles become non-functional after the first clutch, thereby reducing egg production capacity. Structural limitation could therefore contribute to smaller second clutches. However, structural reduction alone cannot explain the simultaneous decline in egg size. Under a strict allocation trade-off, a reduction in egg number would be expected to lead to compensatory increases in egg size if total resources remain constant. Instead, both traits declined, indicating that females likely entered the second reproductive bout with diminished resource reserves in addition to any physiological constraint. In the wild, females accumulate resources for months before producing their first clutch, whereas in the laboratory, they had only a brief interval (approx. one month) to replenish reserves before producing the second. Late reproduction, therefore, likely reflects both reduced energetic state and structural limitation.

The hierarchical variance structure supports a constraint-based interpretation of clutch-size variation. Egg size exhibited moderate repeatability across broods, indicating stable differences among females in mean per-offspring investment. In contrast, egg number showed extremely low

repeatability, with most of the variance arising from brood effects and distribution-level variance rather than from persistent female differences. The absence of any association between egg number and female structural size or condition, together with the near-zero repeatability of clutch size, suggests that variation in egg number is not driven by stable differences in resource acquisition but instead reflects transient physiological state or clutch-level constraint. Thus, clutch size appears primarily state-sensitive, whereas egg size reflects a more stable allocation strategy (Houston and McNamara, 1999; McNamara and Houston, 1996). At the same time, within-clutch variance increased substantially in second broods, indicating reduced developmental precision under diminished resource availability. Structural and energetic constraints, therefore, appear to reduce total reproductive output while leaving mean allocation rules largely intact (Ricklefs and Wikelski, 2002; Zera and Harshman, 2001).

These patterns have broader implications for phenotypic integration and evolutionary potential. Repeatability places an upper bound on heritability (Falconer and MacKay, 1996; Lessells and Boag, 1987; Nakagawa and Schielzeth, 2013; Nakagawa and Schielzeth, 2010; but see Dohm, 2002). The moderate repeatability of egg size suggests that selection could act on persistent differences in mean provisioning, whereas the extremely low repeatability of egg number implies that clutch size may respond primarily to short-term environmental or physiological conditions (i.e. reduced ovariole function). The weak intercept correlation further indicates that egg size and egg number are only loosely integrated, consistent with models in which traits differ in their sensitivity to resource state versus allocation strategy (Houle, 1991). In this system, per-offspring investment appears more evolutionarily stable than the total number of offspring ~~number~~.

Viewed hierarchically, my results illustrate a broader conceptual point: life-history trade-offs may operate primarily through systematic differences in mean-allocation rules among individuals, while most short-term variation reflects state-dependent constraint. ~~However, when among-individual variance in one trait is minimal—as is the case for egg number here—such trade-offs will necessarily be weak and difficult to detect.~~ Without decomposing within- and among-individual covariance, ~~even this~~ the modest pattern among-female trade-off suggested here would have remained obscured. More generally, the evolutionary consequences of trade-offs depend not only on their sign but on their hierarchical expression and repeatability. A trade-off that reflects consistent among-individual differences in a repeatable trait is more likely to shape evolutionary responses than one arising primarily from transient, plastic variation within individuals. Future work measuring resource acquisition directly, experimentally manipulating body condition, or estimating genetic correlations between egg size and egg number will help determine whether the limited-allocation divergence observed here reflects adaptive differentiation in allocation strategy or condition-dependent constraint (Lande, 1980; Roff, 1992).

Data availability

Data are available through the OSF repository at https://osf.io/jrs28/overview?view_only=7a78f30e724b497ab3bb62ffa1c97e10.

Author contributions

CDK collected and analyzed the data, and wrote the manuscript.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Funding

The study was funded by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada (NSERC).

Conflict of Interest

I declare no conflict of interest.

Acknowledgements

I thank Andréanne Nault and Clara Smythe for assistance in animal care and data collection. I also thank Denis Réale, Pierre-Olivier Montiglio, and two anonymous reviewers for their discussion and feedback.

Funding

~~The study was funded by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada (NSERC).~~

Conflict of Interest

~~I declare no conflict of interest.~~

Data availability

~~Data are available through the OSF repository at 10.17605/OSF.IO/KWCS8.~~

Author contributions

CDK collected and analyzed the data, and wrote the manuscript.

Ethics Statement

All data were collected using the European earwig (*Forficula auricularia*), an invertebrate species for which institutional animal care approval and collection permits are not required under current regulations. All procedures adhered to standard ethical guidelines for the care and use of animals in research.

Artificial intelligence (AI)-assisted technologies

Alt text for the two figures was generated with the assistance of AI tools and reviewed and verified by the author for accuracy and completeness.

References

Bernardo, J., 1996. The particular maternal effect of propagule size, especially egg size: Patterns, models, quality of evidence and interpretations. *Am Zool* **36**, 216–236.

Careau, V., Wilson, R., 2017. Of uberfleas and krakens: Detecting trade-offs using mixed models. *Integr Comp Biol* **57**, 362–371.

Clutton-Brock, T.H., 1991. The Evolution of Parental Care. Princeton University Press.

De Jong, G., Van Noordwijk, A., 1992. Acquisition and allocation of resources: Genetic (co) variances, selection, and life histories. *Am Nat* **139**, 749–770.

Dingemanse, N.J., Dochtermann, N., 2013. Quantifying individual variation in behaviour: Mixed-effect modelling approaches. *J Anim Ecol* **82**, 39–54.

Dohm, M.R., 2002. Repeatability estimates do not always set an upper limit to heritability. *Funct Ecol* **16**, 273–280.

Falconer, D.S., MacKay, T.F.C., 1996. An Introduction Quantitative Genetics. Pearson Education Limited, Essex, England.

Fox, C.W., Czesak, M.E., 2000. Evolutionary ecology of progeny size in arthropods. *Annu Rev Entomol* **45**, 341–369.

Hargitai, R., Török, J., Tóth, L., Hegyi, G., Rosivall, B., Szigeti, B., Szöllősi, E., 2005. Effects of environmental conditions and parental quality on inter-and intraclutch egg-size variation in the collared flycatcher (*Ficedula albicollis*). *Auk* **122**, 509–522.

Honek, A., 1993. Intraspecific variation in body size and fecundity in insects: A general relationship. *Oikos* **483**–492.

Houle, D., 1991. Genetic covariance of fitness correlates: What genetic correlations are made of and why it matters. *Evolution* **45**, 630–648.

Houston, A.I., McNamara, J.M., 1999. Models of adaptive behaviour: An approach based on state. Cambridge University Press.

Johnson, M.T.J., Nassrullah, Z., 2024. The improbability of detecting trade-offs and some practical solutions. *J Evol Biol* **37**, 1205–1214.

Kay, M., 2020. tidybayes: Tidy Data and Geoms for Bayesian Models [WWW Document]. *R package version 3.0.5*. URL <http://mjskay.github.io/tidybayes/>

Kelly, C.D., Tawes, B., Worthington, A., 2014. Evaluating indices of body condition in two cricket species. *Ecol Evol* **4**, 4476–4487.

- 488 Koch, L.K., Meunier, J., 2014. Mother and offspring fitness in an insect with maternal care:
489 Phenotypic trade-offs between egg number, egg mass and egg care. *BMC Evol Biol* **14**, 125.
- 490 Lack, D., 1947. The significance of clutch-size in the partridge (*Perdix perdix*). *J Anim Ecol*
491 **16**, 19–25.
- 492 Lande, R., 1980. The genetic covariance between characters maintained by pleiotropic
493 mutations. *Genetics* **94**, 203–215.
- 494 Lessells, C., Boag, P.T., 1987. Unrepeatable repeatabilities: A common mistake. *Auk* **104**,
495 116–121.
- 496 McNamara, J.M., Houston, A.I., 1996. State-dependent life histories. *Nature* **380**, 215–221.
- 497 Meunier, J., 2024. The biology and social life of earwigs (Dermaptera). *Annu Rev Entomol* **69**,
498 259–276.
- 499 Nakagawa, S., Johnson, P.C.D., Schielzeth, H., 2017. The coefficient of determination r^2 and
500 intra-class correlation coefficient from generalized linear mixed-effects models revisited
501 and expanded. *J R Soc Interface* **14**, 20170213.
- 502 Nakagawa, S., Schielzeth, H., 2013. A general and simple method for obtaining R^2 from
503 generalized linear mixed-effects models. *Methods Ecol Evol* **4**, 133–142.
- 504 Nakagawa, S., Schielzeth, H., 2010. Repeatability for Gaussian and non-Gaussian data: A
505 practical guide for biologists. *Biol Rev Camb Philos Soc* **85**, 935–956.
- 506 Parker, G.A., 1982. Why are there so many tiny sperm? Sperm competition and the
507 maintenance of two sexes. *J Theor Biol* **96**, 281–294.
- 508 Peig, J., Green, A.J., 2010. The paradigm of body condition: A critical reappraisal of current
509 methods based on mass and length. *Funct Ecol* **24**, 1323–1332.
- 510 Peig, J., Green, A.J., 2009. New perspectives for estimating body condition from mass/length
511 data: The scaled mass index as an alternative method. *Oikos* **118**, 1883–1891.
- 512 Pol, M. van de, Wright, J., 2009. A simple method for distinguishing within- versus between-
513 subject effects using mixed models. *Anim Behav* **77**, 753–758.
- 514 R Core Team, 2025. R: A language and environment for statistical computing.
- 515 Reznick, D., Nunney, L., Tessier, A., 2000. Big houses, big cars, superfleas and the costs of
516 reproduction. *Trends Ecol Evol* **15**, 421–425.
- 517 Ricklefs, R.E., Wikelski, M., 2002. The physiology/life-history nexus. *Trends Ecol Evol* **17**,
518 462–468.

1
2
3
4 519 Roff, D., 1992. Evolution of Life Histories: Theory and Analysis. Chapman & Hall, Inc., New
5 520 York, NY.
6
7 521 Roff, D.A., Fairbairn, D.J., 2007. The evolution of trade-offs: Where are we? *J Evol Biol* **20**,
8 522 433–447.
9
10 523 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
11 524 Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K.,
12 525 Tomancak, P., Cardona, A., 2012. Fiji: An open-source platform for biological-image analysis.
13 526 *Nat Method* **9**, 676–682.
14
15 527 Simpson, E.H., 1951. The interpretation of interaction in contingency tables. *J. R. Stat. Soc.*
16 528 *Ser. B Stat. Methodol.* **13**, 238–241.
17
18 529 Smith, C.C., Fretwell, S.D., 1974. The optimal balance between size and number of offspring.
19 530 *Am Nat* **108**, 499–506.
20
21 531 Staerkle, M., Kölliker, M., 2008. Maternal food regurgitation to nymphs in earwigs (*Forficula*
22 532 *auricularia*). *Ethology* **114**, 844–850.
23
24 533 Stearns, S.C., 1992. The Evolution of Life Histories. Oxford University Press.
25
26 534 Stearns, S.C., 1989. Trade-offs in life-history evolution. *Funct Ecol* **3**, 259–268.
27
28 535 Stoffel, M.A., Nakagawa, S., Schielzeth, H., 2017. RptR: Repeatability estimation and variance
29 536 decomposition by generalized linear mixed-effects models. *Methods Ecol Evol* **8**, 1639–
30 537 1644.
31
32 538 Tourneur, J.-C., 2018. Factors affecting the egg-laying pattern of *Forficula auricularia*
33 539 (Dermaptera: Forficulidae) in three climatologically different zones of North America. *Can*
34 540 *Entomol* **150**, 511–519.
35
36 541 Tourneur, J.-C., Gingras, J., 1992. Egg laying in a northeastern North American (Montréal,
37 542 Québec) population of *Forficula auricularia* L. (Dermaptera: Forficulidae). *Can Entomol* **124**,
38 543 1055–1061.
39
40 544 Van Noordwijk, A.J., Jong, G. de, 1986. Acquisition and allocation of resources: Their
41 545 influence on variation in life history tactics. *Am Nat* **128**, 137–142.
42
43 546 Westneat, D.F., Araya-Ajoy, Y.G., Allegate, H., Class, B., Dingemanse, N., Dochtermann, N.A.,
44 547 Garamszegi, L.Z., Martin, J.G.A., Nakagawa, S., Réale, D., Schielzeth, H., 2020. Collision
45 548 between biological process and statistical analysis revealed by mean centring. *J Anim Ecol*
46 549 **89**, 2813–2824.
47
48 550 Wickham, H., 2016. ggplot2: Elegant graphics for data analysis. Springer, New York.
49
50
51
52
53
54
55
56
57
58
59
60

- 551 Winkler, D.W., Wallin, K., 1987. Offspring size and number: A life history model linking
552 effort per offspring and total effort. *Am Nat* **129**, 708–720.
- 553 Zera, A.J., Harshman, L.G., 2001. The physiology of life history trade-offs in animals. *Annu*
554 *Rev Ecol Syst* **32**, 95–126.

555

For Review Only

556 ##Figures




Figure 1: Partitioning the egg size–number trade-off into within- and between-female components. (a) Within-female (plastic) relationship between egg number and mean egg size. Points represent individual broods (first clutch: open circles; second clutch: closed circles), with grey lines connecting repeated measures from the same female. The x-axis shows standardized within-female deviation in egg number (0 = female mean). The solid black line represents the Bayesian population-level estimate of the within-female slope ($\beta_W = -0.01$, 95% CrI (-0.06, 0.05)). (b) Between-female relationship (i.e., differences in mean allocation strategy) based on female means across broods plotted against standardized between-female deviation in egg number. The black line represents the Bayesian population-level between-female slope (β_B

= -0.05, 95% CrI (-0.10, 0.00). ~~The~~Together, the panels distinguish plastic allocation within females from strategic differences among females, indicating weak support for a reproductive size–number trade-off.

~~Alt text: Two-panel scatterplot showing relationships between egg number and mean egg size.~~

~~Panel (a) plots within-female deviations, with paired points for first and second clutches~~

~~connected by lines; the trend line is nearly flat, indicating little within-female trade-off. Panel (b)~~

~~plots between-female deviations, with a downward-sloping regression line indicating that females~~

~~producing more eggs tend to have smaller eggs.~~




Figure 2: Clutch-specific variance components of egg size. (a) Between-female variance in egg size for first and second clutches, estimated from a hierarchical Bayesian model with clutch-specific female effects. (b) Within-clutch (egg-to-egg) variance for first and second clutches. Orange points represent posterior draws, black points indicate posterior medians, and vertical lines denote 95% credible intervals. Between-female variance did not differ clearly between clutches, whereas within-clutch variance was greater in second clutches.

Alt-text: Two-panel dot plot showing variation in egg size across clutch order. The left panel shows between-female variation, with distributions for first and second clutches; variance is higher for first clutches than second clutches, with mean values indicated by black points

and vertical error bars. The right panel shows within-clutch variation, with second clutches exhibiting greater variance than first clutches; means and error bars are shown in black.

For Review Only