



Disentangling within- and between-female egg size–number trade-offs in the European earwig

Journal:	<i>Current Zoology</i>
Manuscript ID	CZ-2026-0008.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Kelly, Clint; UQAM Departement des sciences biologiques
Keywords:	fecundity, life-history trade off, residual correlation, mixed models, egg size, clutch size

SCHOLARONE™
Manuscripts

Disentangling within- and between-female egg size–number trade-offs in the European earwig.

Clint D. Kelly^{1,✉}

¹ Département des Sciences biologiques, Université du Québec à Montréal, Montréal, Canada

✉ Correspondence: [Clint D. Kelly <kelly.clint@uqam.ca>](mailto:kelly.clint@uqam.ca)

For Review Only

Abstract

Life-history theory predicts a trade-off between offspring size and number, yet empirical studies often fail to detect the expected negative association. This discrepancy may occur because phenotypic correlations confuse variation in resource acquisition with variation in allocation strategy. Using repeated measures of female European earwigs (*Forficula auricularia*), I partitioned covariance between egg size and egg number into among- and within-female components using Bayesian mixed models. The raw phenotypic correlation was near zero. However, females that produced more eggs on average laid smaller eggs, revealing a modest negative association among females that aligns with an allocation trade-off. In contrast, there was no detectable within-female covariance once clutch order was accounted for. Egg number repeatability was effectively zero, suggesting that clutch size varies mainly within rather than among females, while egg size was highly repeatable. Both egg number and egg size decreased in second clutches, and within-clutch egg-size variance increased, indicating a reproductive constraint (i.e., fewer functional ovarioles) under a possibly diminished resource state. Together, these findings demonstrate that the size–number trade-off is primarily expressed among females through consistent differences in allocation strategy, whereas clutch size itself remains flexible. By separating hierarchical sources of variation, this study shows that allocation trade-offs can remain hidden at the phenotypic level and that evolutionary constraints act through stable allocation rules rather than through flexible clutch-size adjustments.

Introduction

Life-history theory predicts that organisms must allocate finite resources among competing fitness components. Because energy and time are limited, investment in one trait constrains investment in others (Roff, 1992; Stearns, 1992). A central expression of this principle is the offspring size–number trade-off: for a given reproductive budget, producing larger offspring reduces the number that can be produced (Parker, 1982; Smith and Fretwell, 1974). Since larger eggs often enhance offspring performance and survival (Clutton-Brock, 1991), selection is expected to favour an optimal balance between egg size and egg number that maximises maternal fitness (Fox and Czesak, 2000). Yet empirical tests of the size–number trade-off often fail to reveal the expected negative relationship. Across individuals within populations, egg size and egg number are often positively correlated (Bernardo, 1996; Fox and Czesak, 2000; Hargitai et al., 2005; Roff, 1992). This apparent contradiction arises because phenotypic correlations conflate two distinct biological processes: variation in resource acquisition and variation in allocation strategy. Van Noordwijk and De Jong (1986) showed that when variance in acquisition exceeds variance in allocation, individuals with greater resource pools can produce both more and larger eggs, generating positive among-individual correlations even if all individuals are constrained by the same underlying allocation trade-off. In such cases, the population-level pattern masks allocation constraints operating within individuals—an outcome conceptually analogous to Simpson’s paradox (Simpson, 1951).

From this perspective, the critical prediction of life-history theory is not simply a negative phenotypic correlation, but a negative within-individual covariance between egg size and egg number once among-individual differences in acquisition are accounted for (Careau and Wilson,

2017). Detecting such a masked trade-off, therefore, requires partitioning phenotypic variance and covariance into hierarchical components. Repeated measures of individuals allow estimation of (i) among-individual variance, reflecting consistent differences among females in total reproductive investment, and (ii) within-individual variance, reflecting how females adjust allocation across reproductive bouts. Careau and Wilson (2017) argue that only by decomposing covariance into these components using multivariate mixed models can we determine whether allocation trade-offs operate within individuals despite positive population-level associations. Crucially, this framework also predicts that traits involved in allocation trade-offs should show substantial repeatability. If females differ consistently in resource acquisition, egg size and egg number should exhibit among-individual variance across clutches. High repeatability would therefore indicate persistent differences in total reproductive investment among females, consistent with variation in acquisition. At the same time, detecting a negative within-individual covariance would indicate that, conditional on their resource pool, females trade off egg size against egg number across reproductive events. Thus, variance partitioning and repeatability are not merely statistical descriptors, but tests of the underlying acquisition–allocation structure predicted by life-history theory.

Despite the foundational importance of the size–number trade-off, empirical tests rarely partition covariance into within- and among-individual components to distinguish allocation from acquisition effects. Here, I address this gap using the European earwig, *Forficula auricularia*, a species that allows repeated measures of reproductive investment within females but whose pattern of allocation to egg number and size is unclear (Koch and Meunier, 2014). Female earwigs lay a clutch within a short temporal window in a subterranean burrow (Meunier, 2024)

1
2
3
4 71 and provide extended maternal care (Meunier, 2024; Staerkle and Kölliker, 2008). In Quebec
5
6 72 populations, environmental constraints typically render females effectively semelparous
7
8 73 (Tourneur, 2018), but under favourable laboratory conditions, they can produce a second clutch.
9
10 74 This system, therefore, enables direct estimation of within-female covariance across successive
11
12 75 reproductive bouts. I test three predictions derived from acquisition–allocation theory (Van
13
14 76 Noordwijk and Jong, 1986; see also Reznick, Nunney, and Tessier, 2000). First, if variance in
15
16 77 resource acquisition among females exceeds variance in allocation strategy, then egg size and egg
17
18 78 number should exhibit substantial repeatability and a positive among-individual correlation,
19
20 79 whereas the within-individual correlation should be negative, revealing a masked allocation
21
22 80 trade-off. Second, if maternal fitness is maximised at a particular offspring size or number (Smith
23
24 81 and Fretwell, 1974), then one trait should remain relatively invariant across clutches while the
25
26 82 other adjusts between reproductive bouts (Winkler and Wallin, 1987). Third, if reproductive
27
28 83 investment is structured differently across bouts, then variance in egg size may differ between
29
30 84 first and second clutches at both the among-female and within-clutch levels (e.g. Hargitai et al.,
31
32 85 2005), providing further insight into how selection shapes the distribution of maternal investment
33
34 86 across reproductive events.
35
36
37
38
39
40
41 87 By separating among-female differences in total reproductive investment from within-female
42
43 88 allocation decisions, this study tests a fundamental question in life-history theory: do apparent
44
45 89 positive associations between egg size and number reflect relaxed constraints, or do allocation
46
47 90 trade-offs persist but remain hidden within individuals? Demonstrating a negative within-female
48
49 91 covariance despite positive population-level patterns would show that acquisition can mask, but
50
51 92 not erase, the constraints imposed by finite resources. In this way, my study evaluates not only
52
53
54
55
56
57
58
59
60

the existence of a trade-off but also the hierarchical structure through which life-history constraints are expressed.

Methods

I collected adult female earwigs (n=42) in September and October 2024 in an urban park (Parc Outremont) in Montréal, Canada. All earwigs were brought into the laboratory, assigned a unique ID, and individually placed in a 90-mm-diameter Petri dish containing a thin layer of moist sand. Females were provided a cotton-plugged water vial and fed TetraMin Tropical fish flakes *ad libitum*. Females were maintained in an incubator (Percival Scientific, Iowa, USA) at 10°C in constant darkness.

Females were checked daily for oviposition. When a female laid eggs, I recorded the date and collected all eggs from the sand and placed them on filter paper in a new Petri dish. The eggs were then photographed under a Leica S6D stereo microscope (Leica Microsystems Inc., Concord, ON, Canada) at 0.68x magnification using Enersight software (Leica Microsystems Inc., Concord, ON, Canada), which automatically adds a scale bar to the photo. Eggs were then discarded.

To stimulate females to lay a second clutch of eggs, I recreated a brief period of spring where they were allowed to accumulate resources for egg production and replenish their sperm stores.

To achieve this, females that had recently oviposited were placed in a new Petri dish containing a cotton-plugged water vial, TetraMin Tropical fish flakes, and a shelter consisting of four 4 cm-long straws (Blowholes, Markham, Canada) glued together. These Petri dishes were placed in a 23 °C incubator with a 12L:12D light: dark cycle. Females were maintained under these

conditions for two weeks, at which time a haphazardly chosen stock male from our colony was placed with her. The pair was kept together for a week, after which a new stock male replaced the original male, and that pair was kept together for another week. Females were then transferred to the 10 °C incubator and treated as previously until they laid their second clutch, at which point their eggs were photographed. Females were euthanised by freezing after laying their second clutch, and their pronotum was photographed as described above. Because female fecundity is often related to body size in insects (Honek, 1993) and could thus possibly explain a positive among-female covariance between egg number and size, I measured two proxies of size, pronotum length and body mass. Females were weighed to the nearest 0.01 mg using a Sartorius Secura 224-1S analytical balance. Females were weighed three times, with each weighing bout separated by four days. I used Fiji image analysis software (Schindelin et al., 2012) to measure the length of the female pronotum to the nearest 0.001 mm three times, with each measurement separated by 4 days. I used the same procedure to measure the perimeter of each egg.

Statistical analysis

I calculated the repeatability of the body mass, pronotum length and egg perimeter measurements using the R package *rptR* (Stoffel, Nakagawa, and Schielzeth, 2017). I entered the trait of interest as the response variable, female ID as a random effect, and assumed a Gaussian distribution. Each analysis was based on 1000 bootstrap samples. The measurements of my two morphological traits were highly repeatable (pronotum length: R [95% CI] = 0.98 [0.97, 0.99]; body mass: R [95% CI] = 0.99 [0.988, 0.996]); therefore, I used each female's mean value for each trait where necessary. Because egg perimeter was highly repeatable in both clutches (first clutch: R [95% CI] = 0.977 [0.976, 0.979]; second clutch: R [95% CI] = 0.93 [0.926, 0.937]), I

calculated each egg's average perimeter and then used these values to calculate the mean egg size for each clutch for each female.

To reduce the two correlated measures of body size (pronotum length and body mass) to a single structural size metric, I performed a principal components analysis (PCA) on the correlation matrix of standardized variables using the *prcomp* function in the R stats package (R Core Team, 2025). The first principal component (PC1) explained 60.8% of the total variance and was used as an index of structural body size. In addition, female body condition was estimated using the scaled mass index (SMI) following Peig and Green (2010, 2009; see also Kelly, Tawes, and Worthington, 2014), which standardizes body mass relative to structural size (pronotum length).

I first quantified associations among reproductive traits using raw phenotypic correlations. Pearson correlation coefficients were calculated using individual-level trait values, thereby capturing total phenotypic covariation across all sources of variation. These correlations reflect the combined contribution of consistent differences among females, within-clutch variation, and residual effects. Consequently, raw phenotypic correlations provide a descriptive estimate of overall trait associations but do not distinguish between covariation arising from stable female differences and that arising from within-clutch processes.

To distinguish covariance arising from within-female plasticity from covariance attributable to consistent among-female differences, I followed the within-between decomposition approach of van de Pol and Wright [(2009); see Supplementary Material for details]. Although mean-centering has been shown to impose assumptions about underlying biological processes (Westneat et al., 2020), these assumptions are appropriate here because the biological question

1
2
3
4 157 concerns how reproductive allocation varies within females across clutches versus how females
5
6 158 differ consistently in allocation strategy. Egg number was decomposed into two components: a
7
8 159 between-female component, defined as each female's mean egg number across broods centered
9
10 160 on the population mean, and a within-female component, defined as the deviation of each brood
11
12 161 from that female's own mean. These two predictors were included simultaneously in a Bayesian
13
14 162 multivariate mixed-effects model in which egg number and mean egg size were modelled jointly.
15
16 163 Egg number was modelled using a negative binomial distribution with a log link to account for
17
18 164 overdispersion, whereas mean egg size was modelled assuming a Gaussian distribution with an
19
20 165 identity link. Clutch (first vs. second clutch) was included as a fixed effect in both submodels.
21
22 166 Female identity was included as a random intercept for both response variables. This structure
23
24 167 allowed estimation of among-female variance in egg number, among-female variance in egg size,
25
26 168 and the covariance (correlation) between female-level intercepts for the two traits.
27
28
29
30
31
32 169 This multivariate framework partitions total phenotypic covariance into two components:
33
34 170 covariance attributable to consistent differences among females is captured by the female-level
35
36 171 random effects, whereas covariance arising from deviations within females across clutches is
37
38 172 captured by the fixed within-female component of egg number. The fixed between-female
39
40 173 component quantifies how mean egg size differs among females based on their average egg
41
42 174 number, thereby estimating the among-female (strategic) association. In contrast, the fixed
43
44 175 within-female component estimates whether egg size changes plastically when a female produces
45
46 176 more or fewer eggs than her own mean. The correlation between female-level intercepts
47
48 177 represents covariance in baseline trait values among females after accounting for all fixed effects.
49
50
51
52 178 Thus, the between-female slope tests whether females diverge in their mean allocation strategy
53
54
55
56
57
58
59
60

(i.e., whether females that consistently produce more eggs also consistently produce smaller eggs). In contrast, the intercept correlation tests for latent covariance between baseline egg number and egg size after accounting for brood effects and the within–between decomposition, reflecting residual integration not explained by mean allocation differences. Continuous predictors were standardized prior to analysis. Standardized coefficients, therefore, represent the expected change in egg size, measured in standard deviation units, associated with a one standard deviation change in egg number at the relevant hierarchical level.

To quantify the structure of egg-size variation specifically, I fitted a univariate mixed-effects model in which egg size (standardised mean egg perimeter) was modelled as a Gaussian response with female identity included as a random intercept (see Supplementary Material for details). This formulation partitions total phenotypic variance in egg size into variance attributable to differences among females and variance occurring within females among eggs within clutches. The among-female variance component reflects consistent maternal differences in egg size allocation, whereas the within-female component captures heterogeneity in provisioning within clutches and residual variation.

Repeatability was calculated on the observation scale for both traits. For egg size (Gaussian model):

$$R_{\text{egg size}} = \frac{V_F}{V_F + V_R}$$

where V_F is the among-female variance (i.e., variance of the female random intercept) and V_R is the residual variance. Because the Gaussian model is parameterized directly on the observation scale, this formulation corresponds to both latent- and observation-scale repeatability.

For egg number, which was modelled using a negative binomial distribution, the variance depends on both the mean and the dispersion (shape) parameter. Under the negative binomial parameterization used in *brms*, the observation-level variance is given by:

$$\text{Var}(Y) = \mu + \frac{\mu^2}{\theta}$$

where μ is the expected mean egg number and θ is the dispersion (shape) parameter. Observation-scale repeatability for egg number was therefore calculated as:

$$R_{\text{egg number}} = \frac{V_F}{V_F + \left(\mu + \frac{\mu^2}{\theta}\right)}$$

where V_F is the among-female variance estimated from the random intercept. The distribution-specific variance term $\mu + \mu^2/\theta$ was calculated for each posterior draw using the corresponding posterior samples of μ and θ , thereby generating a posterior distribution of repeatability.

Repeatability for non-Gaussian traits was calculated following the framework of Nakagawa and Schielzeth (2010) and the extensions for generalized linear mixed models described by Nakagawa et al. (2017). Because distribution-specific variance can contribute substantially to total phenotypic variance in negative binomial models, repeatability estimates for count data are often lower than those obtained under Gaussian assumptions, even when among-individual variance is present (Nakagawa, Johnson, and Schielzeth, 2017; Nakagawa and Schielzeth, 2010).

By calculating repeatability on the observation scale for both traits, these estimates represent the proportion of total phenotypic variance in the observed trait attributable to consistent differences among females (Nakagawa and Schielzeth, 2010). Derived repeatability values were calculated for each posterior draw to propagate uncertainty from all variance components (Villemereuil, Gimenez, and Doligez, 2013).

I used weakly informative priors to constrain parameters to biologically realistic ranges while allowing the data to drive inference. Regression coefficients were assigned normal priors centered at zero ($\text{Normal}(0, 0.5-1)$), reflecting the expectation that clutch effects on egg traits would be moderate in magnitude. Intercepts were given normal priors scaled to each trait (egg number: $\text{Normal}(0, 5)$; mean egg size: $\text{Normal}(3.5, 1)$; standardized egg size: $\text{Normal}(0, 0.5)$).

Among-female variation was assigned exponential priors ($\text{Exponential}(1-2)$), which favour small but non-zero individual differences. Residual variance (Gaussian models) and the negative binomial dispersion parameter were given $\text{Exponential}(1)$ priors. Correlations among female-level random effects were regularized using an $\text{LKJ}(2)$ prior to avoid overestimating trait associations. Posterior distributions were obtained using six Markov chains with 4,000 iterations per chain, including 1,000 warm-up iterations, and convergence was assessed using $\hat{R}$ statistics, effective sample sizes, and trace-plot inspection. Posterior distributions are summarised as posterior means with 95% credible intervals. Data were visualized using *ggplot* (Wickham, 2016) and *tidybayes* (Kay, 2020).

Results

Between- and within-female associations

Across all broods, the raw phenotypic correlation between egg number and egg size was weak and near zero ($r_{raw} = -0.04$, 95% CI [-0.25, 0.19]), indicating no detectable population-level association when hierarchical structure was ignored.

Egg number (clutch one: $\bar{x} = 51.55 \pm 2.49$ eggs; clutch two: $\bar{x} = 34.14 \pm 2.65$ eggs) and egg size (first clutch: $\bar{x} = 3.54 \pm 0.03$ mm; second clutch: $\bar{x} \pm SE = 3.41 \pm 0.03$ mm) both declined in second clutches. Egg number decreased on the log scale ($\beta = -0.41$, 95% CrI (-0.64, -0.19; [Figure 1](#); Supplementary Table 1), corresponding to an approximate 34% reduction on the natural scale ($e^{-0.413} \approx 0.66$). Mean egg size also declined in the second clutch ($\beta = -0.13$, 95% CrI (-0.24, -0.02)).

The results did not support my prediction of a within-female plastic trade-off between egg number and egg size ([Figure 1 a](#)). The within-female deviation in egg number had a slope near zero ($\beta_W = -0.01$, 95% CrI (-0.06, 0.05), indicating that females did not systematically adjust egg size when producing more or fewer eggs relative to their own mean. In contrast, the between-female component revealed a negative association between egg size and mean female egg number ($\beta_B = -0.05$, 95% CrI (-0.10, 0.00; [Figure 1 b](#)). The posterior distribution of the among-female slope was centered on negative values, suggesting weak evidence for divergence in allocation strategy among females. The correlation between female-level intercepts for egg number and egg size was highly uncertain (median = -0.02, 95% CrI: -0.81, 0.82), indicating limited information for estimating latent covariance with only two broods per female. With only

two repeated measures per female, intercept correlations are weakly identifiable and should be interpreted cautiously.

No measure of female size/condition was positively correlated with the number of eggs in the first brood (pronotum length: $r = 0.14$, 95% CI [-0.19, 0.43]; body mass: $r = 0.12$, 95% CI [-0.21, 0.43]; PC1: $r = -0.17$, 95% CI [-0.44, 0.15]; scaled mass index: $r_{raw} = -0.02$, 95% CI [-0.35, 0.33]) providing no evidence that larger females or those in better condition produce more eggs. I therefore did not include a measure of size/condition in this model. Together, these results show that the size–number trade-off is not apparent at the raw phenotypic level and is not expressed as within-female plasticity, but emerges as a modest negative association at the among-female level through differences in average allocation strategy.

Egg size variation between females and within clutches

Egg size exhibited substantial variation at both the between-female and within-clutch levels (Figure 2; Supplementary Table 2). Between-female variance was similar across reproductive events, with a median of 0.73 (95% CrI: 0.50, 1.12) in first clutches and 0.61 (95% CrI: 0.41, 0.96) in second clutches. The posterior median difference (second – first) was -0.11 (95% CrI: -0.56, 0.28), and the posterior probability that between-female variance was greater in second clutches was 0.28, indicating no clear evidence of a difference. In contrast, within-clutch (egg-to-egg) variance was substantially higher in second clutches (median = 0.35 (95% CrI: 0.33, 0.38) than in first clutches (median = 0.21 (95% CrI: 0.20, 0.22)). The posterior median difference was 0.14 (95% CrI: 0.12, 0.17), with a posterior probability of 1.00, that within-clutch variance was greater in second clutches.

Egg number and egg size repeatability

Repeatability was calculated from posterior samples of the variance components on the observation scale. Egg number exhibited extremely low repeatability. Although there was some among-female variance in egg number (SD = 0.078, 95% CrI [0.003, 0.270]; Supplementary Table 1), the overwhelming majority of variance arose from distribution-specific (negative binomial) variance. Consequently, observation-scale repeatability was near zero ($R = 2.97 \times 10^{-5}$, 95% CrI [2.01×10^{-8} , 1.61×10^{-4}]), indicating that very little of the total phenotypic variation in clutch size was attributable to consistent differences among females.

In contrast, egg size showed moderate repeatability on the observation scale. Among-female variance (SD = 0.082, 95% CrI [0.006, 0.155]; Supplementary Table 2) was substantial relative to residual variance (SD = 0.191, 95% CrI [0.155, 0.235]), yielding $R = 0.63$ (95% CrI: 0.53, 0.73) in first broods and $R = 0.78$ (95% CrI: 0.70, 0.84) in second broods. These results indicate that females differed consistently in their mean egg size across broods, whereas clutch size was largely context-dependent.

Discussion

Life-history theory predicts a trade-off between offspring size and number because both traits draw from a finite pool of reproductive resources (Lack, 1947; Roff, 1992; Smith and Fretwell, 1974; Stearns, 1992). Yet, as emphasised by van Noordwijk and de Jong (1992), phenotypic correlations need not directly reveal allocation trade-offs, because variation in resource acquisition can obscure underlying allocation rules (Careau and Wilson, 2017; Johnson and Nassrullah, 2024). This acquisition–allocation framework has become central to interpreting life-

history covariance (Houle, 1991; Reznick, Nunney, and Tessier, 2000). By partitioning covariance into within- and among-female components, my results show that the size–number trade-off in this population of earwigs is modestly expressed among females but absent within them, illustrating how hierarchical structure shapes the detectability of life-history trade-offs (Careau and Wilson, 2017; Dingemanse and Dochtermann, 2013; Pol and Wright, 2009; Westneat et al., 2020).

At the phenotypic level, egg size and egg number were uncorrelated in my earwig population, consistent with Koch and Meunier (2014). Under the framework of van Noordwijk and de Jong (1986), such phenotypic independence is expected when variation in resource acquisition obscures underlying allocation trade-offs. After hierarchical decomposition, however, a weak negative association emerged at the among-female level: females that produced more eggs on average tended to lay slightly smaller eggs. This pattern corresponds to the estimated correlation between female-specific intercepts in the multivariate model and suggests modest divergence in average allocation strategy among females. Nevertheless, the credible interval surrounding this intercept correlation overlapped zero, indicating considerable uncertainty in the strength of among-female covariance. Thus, rather than forming a tightly integrated life-history syndrome, egg size and number appear to vary along a shallow axis of allocation divergence embedded within substantial variation in resource acquisition (Roff and Fairbairn, 2007; Stearns, 1989).

In contrast, there was no evidence for a within-female plastic trade-off. Although each female contributed only two broods, the posterior distribution of the within-female slope was tightly centered on zero, suggesting that this absence reflects genuine biological stability rather than limited statistical power. Females did not reduce egg size when producing more eggs than their

1
2
3
4 320 own average; instead, egg number and egg size shifted together across broods. This pattern is
5
6 321 consistent with state-dependent life-history models in which allocation rules remain relatively
7
8 322 stable while total resource state fluctuates across reproductive bouts (Houston and McNamara,
9
10 323 1999; McNamara and Houston, 1996). Rather than dynamically reallocating resources between
11
12 324 offspring size and number, females appear to exhibit clutch-level shifts driven by overall
13
14 325 energetic state.
15
16
17
18 326 Notably, egg number was not correlated with female structural size or body condition (i.e. scaled
19
20 327 mass), indicating that persistent differences in morphology do not explain clutch-size variation. If
21
22 328 stable acquisition differences were driving among-female variation, we would expect egg number
23
24 329 to show repeatability and to covary with these morphological size/condition indicators. The
25
26 330 absence of such patterns weakens the interpretation that persistent acquisition differences—at
27
28 331 least as reflected by morphology—underlie among-female differences in reproductive output.
29
30 332 Instead, clutch size appears predominantly state-sensitive and influenced by clutch-level
31
32 333 constraints.
33
34
35
36
37 334 This interpretation also helps explain the pronounced decline in second clutches. That second
38
39 335 clutches comprised both fewer and smaller eggs is consistent with previous work in North
40
41 336 American and European populations (Koch and Meunier, 2014; Tourneur and Gingras, 1992).
42
43 337 Tourneur and Gingras (1992) proposed that some ovarioles become non-functional after the first
44
45 338 clutch, thereby reducing egg production capacity. Structural limitation could therefore contribute
46
47 339 to smaller second clutches. However, structural reduction alone cannot explain the simultaneous
48
49 340 decline in egg size. Under a strict allocation trade-off, a reduction in egg number would be
50
51 341 expected to lead to compensatory increases in egg size if total resources remain constant. Instead,
52
53
54
55
56
57
58
59
60

both traits declined, indicating that females likely entered the second reproductive bout with diminished resource reserves in addition to any physiological constraint. In the wild, females accumulate resources for months before producing their first clutch, whereas in the laboratory, they had only a brief interval (approx. one month) to replenish reserves before producing the second. Late reproduction, therefore, likely reflects both reduced energetic state and structural limitation.

The hierarchical variance structure supports a constraint-based interpretation of clutch-size variation. Egg size exhibited moderate repeatability across broods, indicating stable differences among females in mean per-offspring investment. In contrast, egg number showed extremely low repeatability, with most variance arising from brood effects and distribution-level variance rather than persistent female differences. The absence of any association between egg number and female structural size or condition, together with the near-zero repeatability of clutch size, suggests that variation in egg number is not driven by stable differences in resource acquisition but instead reflects transient physiological state or clutch-level constraint. Thus, clutch size appears primarily state-sensitive, whereas egg size reflects a more stable allocation strategy (Houston and McNamara, 1999; McNamara and Houston, 1996). At the same time, within-clutch variance increased substantially in second broods, indicating reduced developmental precision under diminished resource availability. Structural and energetic constraints, therefore, appear to reduce total reproductive output while leaving mean allocation rules largely intact (Ricklefs and Wikelski, 2002; Zera and Harshman, 2001).

These patterns have broader implications for phenotypic integration and evolutionary potential. Repeatability places an upper bound on heritability (Falconer and MacKay, 1996; Lessells and

1
2
3
4 364 Boag, 1987; Nakagawa and Schielzeth, 2013; Nakagawa and Schielzeth, 2010; but see Dohm,
5
6 365 2002). The moderate repeatability of egg size suggests that selection could act on persistent
7
8 366 differences in mean provisioning, whereas the extremely low repeatability of egg number implies
9
10 367 that clutch size may respond primarily to short-term environmental or physiological conditions
11
12 368 (i.e. reduced ovariole function). The weak intercept correlation further indicates that egg size and
13
14 369 egg number are only loosely integrated, consistent with models in which traits differ in their
15
16
17 370 sensitivity to resource state versus allocation strategy (Houle, 1991). In this system, per-offspring
18
19 371 investment appears more evolutionarily stable than total offspring number.
20
21
22 372 Viewed hierarchically, my results illustrate a broader conceptual point: life-history trade-offs
23
24 373 may operate primarily through stable differences in allocation rules among individuals, while
25
26
27 374 short-term variation reflects state-dependent constraint. Without decomposing within- and
28
29 375 among-individual covariance, the modest among-female trade-off detected here would have
30
31 376 remained obscured. More generally, the evolutionary consequences of trade-offs depend not only
32
33 377 on their sign but on their hierarchical expression and repeatability. A trade-off that reflects
34
35 378 consistent among-individual differences in a repeatable trait is more likely to shape evolutionary
36
37 379 responses than one arising primarily from transient, plastic variation within individuals. Future
38
39 380 work measuring resource acquisition directly, experimentally manipulating body condition, or
40
41 381 estimating genetic correlations between egg size and egg number will help determine whether the
42
43 382 allocation divergence observed here reflects adaptive differentiation in allocation strategy or
44
45 383 condition-dependent constraint (Lande, 1980; Roff, 1992).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Data availability

Data are available through the OSF repository at 10.17605/OSF.IO/KWCS8.

Author contributions

CDK collected and analyzed the data, and wrote the manuscript.

Funding

The study was funded by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada (NSERC).

Conflict of Interest

I declare no conflict of interest.

Acknowledgements

I thank Andréanne Nault and Clara Smythe for assistance in animal care and data collection. I also thank Denis Réale and Pierre-Olivier Montiglio for their discussion and feedback.

References

Bernardo, J., 1996. The particular maternal effect of propagule size, especially egg size: Patterns, models, quality of evidence and interpretations. *Am Zool* **36**, 216–236.

Careau, V., Wilson, R., 2017. Of uberfleas and krakens: Detecting trade-offs using mixed models. *Integr Comp Biol* **57**, 362–371.

Clutton-Brock, T.H., 1991. The Evolution of Parental Care. Princeton University Press.

De Jong, G., Van Noordwijk, A., 1992. Acquisition and allocation of resources: Genetic (co) variances, selection, and life histories. *Am Nat* **139**, 749–770.

Dingemanse, N., Dochtermann, N., 2013. Quantifying individual variation in behaviour: Mixed-effect modelling approaches. *J Anim Ecol* **82**, 39–54.

Dohm, M.R., 2002. Repeatability estimates do not always set an upper limit to heritability. *Funct Ecol* **16**, 273–280.

Falconer, D.S., MacKay, T.F.C., 1996. An Introduction Quantitative Genetics. Pearson Education Limited, Essex, England.

Fox, C.W., Czesak, M.E., 2000. Evolutionary ecology of progeny size in arthropods. *Annu Rev Entomol* **45**, 341–369.

Hargitai, R., Török, J., Tóth, L., Hegyi, G., Rosivall, B., Szigeti, B., Szöllősi, E., 2005. Effects of environmental conditions and parental quality on inter-and intraclutch egg-size variation in the collared flycatcher (*Ficedula albicollis*). *The Auk* **122**, 509–522.

Honek, A., 1993. Intraspecific variation in body size and fecundity in insects: A general relationship. *Oikos* **483–492**.

Houle, D., 1991. Genetic covariance of fitness correlates: What genetic correlations are made of and why it matters. *Evolution* **45**, 630–648.

Houston, A.I., McNamara, J.M., 1999. Models of adaptive behaviour: An approach based on state. Cambridge University Press.

Johnson, M.T.J., Nassrullah, Z., 2024. The improbability of detecting trade-offs and some practical solutions. *J Evol Biol* **37**, 1205–1214.

Kay, M., 2020. tidybayes: Tidy Data and Geoms for Bayesian Models [WWW Document]. *R package version 3.0.5*. URL <http://mjskay.github.io/tidybayes/>

Kelly, C.D., Tawes, B., Worthington, A., 2014. Evaluating indices of body condition in two cricket species. *Ecology and Evolution* **4**, 4476–4487.

- 428 Koch, L.K., Meunier, J., 2014. Mother and offspring fitness in an insect with maternal care:
429 Phenotypic trade-offs between egg number, egg mass and egg care. *BMC Evol Biol* **14**, 125.
- 430 Lack, D., 1947. The significance of clutch-size in the partridge (*perdix perdix*). *The Journal of*
431 *Animal Ecology* 19–25.
- 432 Lande, R., 1980. The genetic covariance between characters maintained by pleiotropic
433 mutations. *Genetics* **94**, 203–215.
- 434 Lessells, C., Boag, P.T., 1987. Unrepeatable repeatabilities: A common mistake. *The Auk* **104**,
435 116–121.
- 436 McNamara, J.M., Houston, A.I., 1996. State-dependent life histories. *Nature* **380**, 215–221.
- 437 Meunier, J., 2024. The biology and social life of earwigs (dermaptera). *Annu Rev Entomol* **69**,
438 259–276.
- 439 Nakagawa, S., Johnson, P.C.D., Schielzeth, H., 2017. The coefficient of determination r^2 and
440 intra-class correlation coefficient from generalized linear mixed-effects models revisited
441 and expanded. *J R Soc Interface* **14**, 20170213.
- 442 Nakagawa, S., Schielzeth, H., 2013. A general and simple method for obtaining R^2 from
443 generalized linear mixed-effects models. *Methods Ecol Evol* **4**, 133–142.
- 444 Nakagawa, S., Schielzeth, H., 2010. Repeatability for gaussian and non-gaussian data: A
445 practical guide for biologists. *Biol Rev Camb Philos Soc* **85**, 935–956.
- 446 Parker, G.A., 1982. Why are there so many tiny sperm? Sperm competition and the
447 maintenance of two sexes. *J Theor Biol* **96**, 281–294.
- 448 Peig, J., Green, A.J., 2010. The paradigm of body condition: A critical reappraisal of current
449 methods based on mass and length. *Funct Ecol* **24**, 1323–1332.
- 450 Peig, J., Green, A.J., 2009. New perspectives for estimating body condition from mass/length
451 data: The scaled mass index as an alternative method. *Oikos* **118**, 1883–1891.
- 452 Pol, M. van de, Wright, J., 2009. A simple method for distinguishing within- versus between-
453 subject effects using mixed models. *Anim Behav* **77**, 753–758.
- 454 R Core Team, 2025. R: A language and environment for statistical computing.
- 455 Reznick, D., Nunney, L., Tessier, A., 2000. Big houses, big cars, superfleas and the costs of
456 reproduction. *Trends Ecol Evol* **15**, 421–425.
- 457 Ricklefs, R.E., Wikelski, M., 2002. The physiology/life-history nexus. *Trends Ecol Evol* **17**,
458 462–468.

- 459 Roff, D., 1992. Evolution of Life Histories: Theory and Analysis. Chapman & Hall, Inc., New
460 York, NY.
- 461 Roff, D.A., Fairbairn, D.J., 2007. The evolution of trade-offs: Where are we? *J Evol Biol* **20**,
462 433–447.
- 463 Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
464 Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K.,
465 Tomancak, P., Cardona, A., 2012. Fiji: An open-source platform for biological-image analysis.
466 *Nat Methods* **9**, 676–682.
- 467 Simpson, E.H., 1951. The interpretation of interaction in contingency tables. *Journal of the*
468 *Royal Statistical Society: Series B (Methodological)* **13**, 238–241.
- 469 Smith, C.C., Fretwell, S.D., 1974. The optimal balance between size and number of offspring.
470 *Am Nat* **108**, 499–506.
- 471 Staerkle, M., Kölliker, M., 2008. Maternal food regurgitation to nymphs in earwigs (*forficula*
472 *auricularia*). *Ethology* **114**, 844–850.
- 473 Stearns, S.C., 1992. The Evolution of Life Histories. Oxford University Press.
- 474 Stearns, S.C., 1989. Trade-offs in life-history evolution. *Funct Ecol* **3**, 259–268.
- 475 Stoffel, M.A., Nakagawa, S., Schielzeth, H., 2017. RptR: Repeatability estimation and variance
476 decomposition by generalized linear mixed-effects models. *Methods Ecol Evol* **8**, 1639–
477 1644.
- 478 Tourneur, J.-C., 2018. Factors affecting the egg-laying pattern of *forficula auricularia*
479 (dermaptera: Forficulidae) in three climatologically different zones of north america. *The*
480 *Canadian Entomologist* **150**, 511–519.
- 481 Tourneur, J.-C., Gingras, J., 1992. Egg laying in a northeastern North American (montréal,
482 québec) population of *forficula auricularia* l. (Dermaptera: Forficulidae). *The Canadian*
483 *Entomologist* **124**, 1055–1061.
- 484 Van Noordwijk, A.J., Jong, G. de, 1986. Acquisition and allocation of resources: Their
485 influence on variation in life history tactics. *Am Nat* **128**, 137–142.
- 486 Villemereuil, P. de, Gimenez, O., Doligez, B., 2013. Comparing parent–offspring regression
487 with frequentist and bayesian animal models to estimate heritability in wild populations: A
488 simulation study for gaussian and binary traits. *Methods Ecol Evol* **4**, 260–275.
- 489 Westneat, D.F., Araya-Ajoy, Y.G., Allegue, H., Class, B., Dingemanse, N., Dochtermann, N.A.,
490 Garamszegi, L.Z., Martin, J.G.A., Nakagawa, S., Réale, D., Schielzeth, H., 2020. Collision
491 between biological process and statistical analysis revealed by mean centring. *J Anim Ecol*
492 **89**, 2813–2824.

- Wickham, H., 2016. *ggplot2: Elegant graphics for data analysis*. Springer, New York.
- Winkler, D.W., Wallin, K., 1987. Offspring size and number: A life history model linking effort per offspring and total effort. *Am Nat* **129**, 708–720.
- Zera, A.J., Harshman, L.G., 2001. The physiology of life history trade-offs in animals. *Annu Rev Ecol Syst* **32**, 95–126.

For Review Only

512 ##Figures




Figure 1: Partitioning the egg size–number trade-off into within- and between-female components. (a) Within-female (plastic) relationship between egg number and mean egg size. Points represent individual broods (first clutch: open circles; second clutch: closed circles), with grey lines connecting repeated measures from the same female. The x-axis shows standardized within-female deviation in egg number (0 = female mean). The solid black line represents the Bayesian population-level estimate of the within-female slope ($\beta_W = -0.01$, 95% CrI (-0.06, 0.05)). (b) Between-female (strategic) relationship based on female means across broods plotted against standardized between-female deviation in egg number. The black line represents the Bayesian population-level between-female slope ($\beta_B = -0.05$, 95% CrI (-0.10, 0.00)). Together, the panels distinguish plastic allocation within females from strategic differences among females, indicating weak support for a reproductive size–number trade-off.

513




Figure 2: Clutch-specific variance components of egg size. (a) Between-female variance in egg size for first and second clutches, estimated from a hierarchical Bayesian model with clutch-specific female effects. (b) Within-clutch (egg-to-egg) variance for first and second clutches. Grey points represent posterior draws, black points indicate posterior medians, and vertical lines denote 95% credible intervals. Between-female variance did not differ clearly between clutches, whereas within-clutch variance was greater in second clutches.

514

**Supplemental Material: Disentangling within- and between-
female egg size–number trade-offs in the European earwig**

Clint D. Kelly^{1,✉}

¹ Département des Sciences biologiques, Université du Québec à Montréal, Montréal, Canada

✉ Correspondence: [Clint D. Kelly <kelly.clint@uqam.ca>](mailto:kelly.clint@uqam.ca)

For Review Only

1. Within–between multivariate model

To partition covariance arising from within-female plasticity from covariance attributable to consistent among-female differences, I applied the within–between decomposition of van de Pol and Wright (Pol and Wright, 2009). Although mean-centering can impose assumptions about underlying biological processes (Westneat et al., 2020), these assumptions are appropriate here because the aim was to distinguish plastic allocation within females from persistent differences among females.

Let egg number for female i in brood j be denoted N_{ij} . Egg number was decomposed as

$$N_{ij} = \bar{N}_i + (N_{ij} - \bar{N}_i) \quad (1)$$

The between- and within-female components were defined as

$$N_i^B = \bar{N}_i - \bar{N} \quad (2)$$

$$N_{ij}^W = N_{ij} - \bar{N}_i \quad (3)$$

Egg number submodel

Egg number was modelled using a negative binomial distribution with a log link:

$$N_{ij} \sim \text{NegBinomial}(\mu_{ij}, \theta) \quad (4)$$

$$\log(\mu_{ij}) = \beta_0^N + \beta_1^N \text{Brood}_{ij} + u_{0i}^N \quad (5)$$

where

$$u_{0i}^N \sim \mathcal{N}(0, \sigma_N^2) \quad (6)$$

Mean egg size submodel

Mean egg size was modelled as

$$S_{ij} \sim \mathcal{N}(\mu_{ij}^S, \sigma_S^2) \tag{7}$$

$$\mu_{ij}^S = \beta_0^S + \beta_1^S \text{Brood}_{ij} + \beta_W N_{ij}^W + \beta_B N_i^B + u_{0i}^S \tag{8}$$

with

$$u_{0i}^S \sim \mathcal{N}(0, \sigma_{S,id}^2) \tag{9}$$

Multivariate structure

The two submodels were fit jointly:

$$\begin{pmatrix} u_{0i}^N \\ u_{0i}^S \end{pmatrix} \sim \mathcal{MVN}\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \Sigma\right) \tag{10}$$

where

$$\Sigma = \begin{pmatrix} \sigma_N^2 & \sigma_{NS} \\ \sigma_{NS} & \sigma_{S,id}^2 \end{pmatrix} \tag{11}$$

The intercept correlation is

$$\rho_{NS} = \frac{\sigma_{NS}}{\sqrt{\sigma_N^2 \sigma_{S,id}^2}} \tag{12}$$

Relationship between model parameters and total phenotypic covariance

Total phenotypic covariance can be partitioned as

$$\text{Cov}_{\text{total}} = \text{Cov}_{\text{within}} + \text{Cov}_{\text{among}} \tag{13}$$

Within-female covariance

$$\text{Cov}_{\text{within}} = \beta_W \text{Var}(N_{ij}^W) \quad (14)$$

Among-female covariance

$$\text{Cov}_{\text{among, slope}} = \beta_B \text{Var}(N_i^B) \quad (15)$$

$$\text{Cov}_{\text{among, intercept}} = \sigma_{NS} \quad (16)$$

Total among-female covariance is therefore

$$\text{Cov}_{\text{among}} = \beta_B \text{Var}(N_i^B) + \sigma_{NS} \quad (17)$$

Total phenotypic covariance

Combining components:

$$\text{Cov}_{\text{total}} = \beta_W \text{Var}(N_{ij}^W) + \beta_B \text{Var}(N_i^B) + \sigma_{NS} \quad (18)$$

The intercept correlation is

$$\rho_{NS} = \frac{\sigma_{NS}}{\sqrt{\sigma_N^2 \sigma_{S,id}^2}} \quad (19)$$

Corresponding *brms* Specification

```
bf_number <- bf(
  egg.number ~ brood + (1 |p| id),
  family = negbinomial()
)

bf_size <- bf(
  mean.egg.size ~ brood +
    egg_number_within_z +
    egg_number_between_z +
    (1 |p| id),
  family = gaussian()
)
```

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

```
mv_model <- bf_number + bf_size
```

For Review Only

Table S 1: Posterior summaries from the multivariate Bayesian mixed-effects model examining the relationship between egg number and mean egg size. The model jointly estimated a negative binomial model for egg number and a Gaussian model for egg size, including brood effects and partitioning egg number into within- and between-female components. Reported values are posterior medians and 95% credible intervals (CrI). Random effects represent among-female (ID-level) variation, and the correlation term describes the association between female intercepts for egg number and egg size. The dispersion parameter refers to the negative binomial shape parameter.

Parameter	Estimate	Lower 95% CrI	Upper 95% CrI
Egg number: Brood 2 – Brood 1 (log scale)	-0.413	-0.635	-0.189
Egg number: Intercept (Brood 1, log scale)	3.940	3.782	4.107
Egg size: Brood 2 – Brood 1	-0.133	-0.244	-0.024
Egg size: Between-female deviation in egg number (z)	-0.050	-0.104	0.002
Egg size: Within-female deviation in egg number (z)	-0.007	-0.061	0.049
Egg size: Intercept (Brood 1)	3.543	3.468	3.618
cor_id__eggnumber_Intercept__meaneggsize_Intercept	-0.017	-0.813	0.816
SD (Egg number Female ID)	0.078	0.003	0.270
SD (Egg size Female ID)	0.082	0.006	0.155
Residual SD (Egg size)	0.191	0.155	0.235
Dispersion (Egg number NegBin shape)	4.080	2.835	5.853

2. Clutch-Specific Hierarchical Model of Egg Size

Statistical Model

Egg perimeter was standardized prior to analysis (mean = 0, SD = 1).

For female i , egg j , and brood $b \in \{1,2\}$, standardized egg size y_{ijb} was modeled as:

$$y_{ijb} \sim \mathcal{N}(\mu_{ijb}, \sigma_b^2)$$

The conditional mean structure was:

$$\mu_{ijb} = \beta_0 + \beta_1 I(b = 2) + u_{ib}$$

where:

- β_0 is the mean egg size in clutch 1 (reference level),
- β_1 is the fixed difference between clutch 2 and clutch 1,
- u_{ib} is the female-specific deviation for clutch b .

Random Effects Structure

Clutch-specific female deviations were modeled as a correlated bivariate normal vector:

$$\mathbf{u}_i = \begin{pmatrix} u_{i1} \\ u_{i2} \end{pmatrix} \sim \mathcal{MVN} \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \Sigma \right)$$

with variance–covariance matrix:

$$\Sigma = \begin{pmatrix} \tau_1^2 & \rho\tau_1\tau_2 \\ \rho\tau_1\tau_2 & \tau_2^2 \end{pmatrix}$$

Here:

- τ_1^2 and τ_2^2 are the among-female variances in clutches 1 and 2, respectively,
- $\rho = \text{Corr}(u_{i1}, u_{i2})$ is the correlation between female deviations across clutches.

Importantly, no clutch-independent random intercept was estimated. The model includes only clutch-specific female deviations (i.e., random slopes of clutch within female), corresponding to the specification `(0 + brood_factor | id)` in *brms*.

Residual Variance Model

Residual variance was allowed to differ between clutches:

$$\log(\sigma_b) = \alpha_0 + \alpha_1 I(b = 2)$$

where:

- σ_1^2 is the residual variance in clutch 1,
- σ_2^2 is the residual variance in clutch 2.

The log link ensures that residual standard deviations remain positive.

Corresponding *brms* Specification

```
bf(
  scale(mean.perim) ~ brood_factor + (0 + brood_factor | id),
  sigma ~ brood_factor
)
```

Table S 2: Posterior medians and 95% credible intervals from a Bayesian mixed model of scaled egg perimeter, including clutch-specific means, clutch-specific female-level random effects, and clutch-specific residual variance.

Parameter	Estimate	Lower 95% CrI	Upper 95% CrI
Brood 2 – Brood 1	-0.535	-0.827	-0.246
Mean (Brood 1)	0.283	0.050	0.530
Correlation (Brood 1, Brood 2 Female ID)	0.250	-0.025	0.488
SD (Brood 1 Female ID)	0.855	0.704	1.058
SD (Brood 2 Female ID)	0.784	0.638	0.982
Residual SD (Brood 1)	0.457	0.444	0.471
Residual SD (Brood 2)	0.594	0.574	0.616

References

Pol, M. van de, Wright, J., 2009. A simple method for distinguishing within- versus between-subject effects using mixed models. *Anim Behav* **77**, 753–758.

Westneat, D.F., Araya-Ajoy, Y.G., Allegue, H., Class, B., Dingemanse, N., Dochtermann, N.A., Garamszegi, L.Z., Martin, J.G.A., Nakagawa, S., Réale, D., Schielzeth, H., 2020. Collision between biological process and statistical analysis revealed by mean centring. *J Anim Ecol* **89**, 2813–2824.

For Review Only




Figure 1: Partitioning the egg size–number trade-off into within- and between-female components. (a) Within-female (plastic) relationship between egg number and mean egg size. Points represent individual broods (first clutch: open circles; second clutch: closed circles), with grey lines connecting repeated measures from the same female. The x-axis shows standardized within-female deviation in egg number (0 = female mean). The solid black line represents the Bayesian population-level estimate of the within-female slope ($\beta_W = -0.01$, 95% CrI (-0.06, 0.05)). (b) Between-female (strategic) relationship based on female means across broods plotted against standardized between-female deviation in egg number. The black line represents the Bayesian population-level between-female slope ($\beta_B = -0.05$, 95% CrI (-0.10, 0.00)). Together, the panels distinguish plastic allocation within females from strategic differences among females, indicating weak support for a reproductive size–number trade-off.

1146x1146mm (72 x 72 DPI)





Figure 2: Clutch-specific variance components of egg size. (a) Between-female variance in egg size for first and second clutches, estimated from a hierarchical Bayesian model with clutch-specific female effects. (b) Within-clutch (egg-to-egg) variance for first and second clutches. Grey points represent posterior draws, black points indicate posterior medians, and vertical lines denote 95% credible intervals. Between-female variance did not differ clearly between clutches, whereas within-clutch variance was greater in second clutches.

1146x1146mm (72 x 72 DPI)



268x155mm (300 x 300 DPI)