

LECTURE NOTES: LIMITS AND CONTINUITY

S. PETER JEONG

ABSTRACT. A brief introductory exposition on limits, continuity, and related theorems.

1. LIMIT OF A FUNCTION

1.1. Examples of Limits. We begin with an intuitive definition of limits of real-valued functions.

Definition 1.1. For a real-valued function $f : D \rightarrow \mathbb{R}$ defined on some subset $D \subseteq \mathbb{R}$, we say that *the limit of $f(x)$ as x approaches a is L* if the value of $f(x)$ gets arbitrarily close to L as x gets closer and closer to a , without necessarily reaching a .

That is, the existence of a limit depends only on the behavior of the function near the point a , regardless of whether $f(a)$ is defined. In particular, the limit $\lim_{x \rightarrow a} f(x)$ exists if and only if any sequence of numbers $\{x_n\}$ approaching a within some neighborhood of a produces a sequence of numbers $\{f(x_n)\}$ approaching the same value L . Since we can approach a from both sides (either right or left) on the real number line, for the overall limit to exist, the *right-hand limit* $\lim_{x \rightarrow a^+} f(x)$ and *left-hand limit* $\lim_{x \rightarrow a^-} f(x)$ must both exist and equal the same value L .

Here we look at some “nontrivial” examples (or counterexamples) of limits of functions.

Example 1.2. Let $f(x) = \sin \frac{1}{x}$. If we ever try to estimate the limit $\lim_{x \rightarrow 0} f(x)$ by either graphing or calculating $f(x)$ for very small values of x , we quickly run into an issue; the function seems to “fluctuate” across many different values as we take $x = 0.01, 0.00001, 0.0000001$, and so on.

In fact, the function indeed oscillates *infinitely often* on any open interval containing the point 0. In particular, if we let

$$x_1 = \frac{1}{\frac{1}{2}\pi + 2n\pi}, \quad x_2 = \frac{1}{\frac{3}{2}\pi + 2m\pi},$$

then any open interval around 0 contains the points x_1 and x_2 for sufficiently large n and m , while $f(x_1) = 1$ and $f(x_2) = -1$, meaning that the function oscillates between these two values (which are indeed upper and lower bounds of f).

Example 1.3. Let $f(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$

Since there is always an irrational number arbitrarily close to any real number, the function oscillates between 0 and 1 infinitely often on any given interval on $\mathbb{R}$ and the limit $\lim_{x \rightarrow a} f(x)$ does not exist for any $a \in \mathbb{R}$. This function is also known as the *Dirichlet function*.

Example 1.4. Let $f(x) = \begin{cases} \frac{1}{q} & \text{if } x = \frac{p}{q} \text{ in lowest terms,} \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$

Here f is commonly known as the *Thomae function* or the *popcorn function*. Take some $a \in (0, 1)$. Then for any sequence of real numbers $\{x_n\}$ approaching a , we can split the sequence in two parts: (1) a sequence of irrational numbers $\{z'_n\}$ approaching a , and (2) a sequence of rational numbers $\{z_n\}$ approaching a . The first sequence $\{z'_n\}$ of irrational numbers produces a sequence of function values $\{f(z'_n)\}$ that approaches 0.

The second sequence $\{z_n\}$ requires a little more thinking. If we write $z_n = \frac{p_n}{q_n}$ in lowest terms, then the function values $f(z'_n)$ form a sequence of form $\{\frac{1}{q_n}\}$. Now the question becomes: can we find a sequence of rational numbers $\{z_n\}$ approaching a such that the denominators q_n are bounded above by some constant M ? If not, then the function values $f(z_n) = \frac{1}{q_n}$ must approach 0. The answer is no; if we take M to be any positive integer, then there are only finitely many rational numbers in $(0, 1)$ with denominators at most M , so we cannot find a sequence of rational numbers approaching a with bounded denominators. Thus the function values $f(z_n)$ must approach 0 as well, and we conclude that $\lim_{x \rightarrow a} f(x) = 0$ for any $a \in (0, 1)$.

Example 1.5. Now let $f(x) = \frac{\sin \theta}{\theta}$ and consider the limit of $f(\theta)$ as θ approaches 0. Geometrically, $\sin \theta$ is the length of the vertical segment of the unit vector at angle θ from the positive x -axis. We see that the triangle formed by the origin, the point on the unit circle at angle θ , and the unit vector $(1, 0)$ has area $\frac{1}{2} \sin \theta$, while the sector of the unit circle at angle θ has area $\frac{1}{2} \theta$. Since the triangle is contained in the sector, we have

$$\frac{\sin \theta}{2} \leq \frac{\theta}{2}, \quad \text{for } 0 < \theta < \frac{\pi}{2}.$$

On the other hand, the triangle formed by the origin, the unit vector $(1, 0)$, and the point $(1, \tan \theta)$ has area $\frac{1}{2}$. Since this larger right triangle contains the sector, we have the inequality

$$\frac{\theta}{2} \leq \frac{\tan \theta}{2}.$$

Combining the two inequalities, we have

$$\frac{\sin \theta}{2} \leq \frac{\theta}{2} \leq \frac{\tan \theta}{2}.$$

Multiplying through by $\frac{2}{\sin \theta}$, we have

$$1 \leq \frac{\theta}{\sin \theta} \leq \frac{1}{\cos \theta}.$$

for $\theta \in (0, \pi/2)$. Thus $\lim_{x \rightarrow 0^+} \frac{\sin \theta}{\theta} = 1$.

To show that $\lim_{x \rightarrow 0^-} \frac{\sin \theta}{\theta} = 1$, notice that $\sin \theta$ is an odd function. Thus after changing the sign of θ we obtain

$$\begin{aligned} \lim_{x \rightarrow 0^-} \frac{\sin \theta}{\theta} &= \lim_{x \rightarrow 0^+} \frac{\sin(-\theta)}{-\theta} \\ &= \lim_{x \rightarrow 0^+} \frac{-\sin \theta}{-\theta} \\ &= \lim_{x \rightarrow 0^+} \frac{\sin \theta}{\theta} \\ &= 1. \end{aligned}$$

Thus we have $\lim_{x \rightarrow 0} \frac{\sin \theta}{\theta} = 1$.

1.2. The Squeeze Theorem. So far, we have seen examples of limits of non-polynomial or non-rational functions that require a more sophisticated definition of a limit. We continue to examine limits of functions that cannot be computed directly through standard algebraic manipulation.

Theorem 1.6 (Squeeze Theorem). *Let f, g, h be real-valued functions defined on some open interval containing a , except possibly at a itself. If $f(x) \leq g(x) \leq h(x)$ for all x in the interval, and if $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L$, then $\lim_{x \rightarrow a} g(x) = L$ as well.*

Essentially, we are trapping the tricky function g between two functions f and h whose limits are well-defined and have the same value. In that case, the limit of the trapped function g is forced to approach the same value as well.

Example 1.7. Consider the limit $\lim_{x \rightarrow 0} x^2 \sin \frac{1}{x}$. Notice that we cannot algebraically compute this limit since

$$\lim_{x \rightarrow 0} \sin \frac{1}{x}$$

does not exist (example 1.2). We can apply the squeeze theorem by noting that $-1 \leq \sin \frac{1}{x} \leq 1$ for all x , so we have

$$-x^2 \leq x^2 \sin \frac{1}{x} \leq x^2.$$

Since $\lim_{x \rightarrow 0} -x^2 = \lim_{x \rightarrow 0} x^2 = 0$, we conclude that $\lim_{x \rightarrow 0} x^2 \sin \frac{1}{x} = 0$ as well even though the limit of $\sin \frac{1}{x}$ does not exist.