

Automorphic forms and L -functions for the unitary group^{*}

Re- \TeX ed by Seewoo Lee[†]

Stephen Gelbart
Department of Mathematics
Cornell University
Ithaca, New York 14853/USA

and

Ilya Piatetski-Shapiro
Department of Mathematics
Yale University, New Haven, CT. 06520/USA
Tel Aviv University, Ramat-Aviv, Israel

Last updated: October 19, 2023

Introduction

Our purpose is to define and analyze L -functions attached to automorphic cusp forms on the unitary group $G = \mathrm{U}_{2,1}$ and a six-dimensional representation

$$\rho : {}^L G \rightarrow \mathrm{GL}_6(\mathbb{C})$$

of its L -group.

^{*}Notes based on the lectures by S. G. at the University of Maryland Special Year on Lie Group Representations, 1982-83.

[†]seewoo5@berkeley.edu

The motivation for this work is three fold.

Firstly, we use these L -functions to analyze the lifting of cusp forms from $U_{1,1}$ to $U_{2,1}$; here the model for our work is Waldspurger's L -function theoretic characterization of the image of Shimura's map for modular forms of half-integral weight (cf. [Wald]).

A second motivation comes from the need to relate the poles of the L -functions for G , to integrals of cusp forms over cycles coming from $U_{1,1}$. The prototype here is the recent proof of Tate's conjecture for Hilbert modular surfaces due to Harder, Langlands, and Rapaport.

Thirdly, we view this work as a special contribution to the general program of constructing local L and ε factors of Langlands type for representations of arbitrary reductive groups. In [PS1], such a program was sketched generalizing classical methods of Hecke, Rankin–Selberg, and Shimura. Related developments are discussed in [Jacquet], [Novod], [PS2], and [PS3]. For the unitary group $U_{2,1}$ the present paper extends the developments initiated in [PS3].

Contents

Whittaker Models (Ordinary and Generalized)	3
Some Fourier Expansions and Hypercuspidality	6
L-functions à la Rankin–Selberg–Jacquet	10

Notation

- (i) F is a field (sometimes local, sometimes a global field), E is a quadratic extension of F with Galois involution $z \mapsto \bar{z}$.
- (ii) V is a 3-dimensional vector space over E , with basis $\{\ell_{-1}, \ell_0, \ell_1\}$. $(-, -)_V$ is a Hermitian form on V , with matrix

$$\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

with respect to $\{\ell_{-1}, \ell_0, \ell_1\}$.

- (iii) $G = U_{2,1} = U(V)$ is the unitary group for the form $(-, -)_V$. P =parabolic

subgroup stabilizing the isotropic line through $\ell_{-1} = MN$ with

$$M = \left\{ \begin{bmatrix} \delta & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \bar{\delta}^{-1} \end{bmatrix} : \delta \in E^\times, \beta \in E^1 = \{z : z\bar{z} = 1\} \right\}$$

and unipotent radical

$$N = \left\{ \begin{bmatrix} 1 & b & z \\ 0 & 1 & -\bar{b} \\ 0 & 0 & 1 \end{bmatrix} : z, b \in E, z + \bar{z} = -b\bar{b} \right\}.$$

The center of N is

$$Z = \left\{ \begin{bmatrix} 1 & 0 & z \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} : \bar{z} = -z \right\}$$

❖ Whittaker Models (Ordinary and Generalized)

Some kind of Whittaker model is needed in order to introduce L -functions on G .

Fix F local (not of characteristic two), and suppose (π, H_π) is an irreducible admissible representation of G . Naively, we should look for functionals on H_π which transform under N according to a one-dimensional representation. However, since such functionals need not exist in general, and since there are irreducible representations of N which are not 1-dimensional, it is natural to pursue a more general approach.

1.1

Recall N is the maximal unipotent subgroup of G and E is a quadratic extension of F . We fix, once and for all, an element i in E such that $\bar{i} = -i$, so $\Im(z) = (z - \bar{z})/2i$. Regarding E as a 2-dimensional symplectic space over F with skew-form $\langle z_1, z_2 \rangle = \Im(z_1 \bar{z}_2)$ we have

$$N = \left\{ \begin{bmatrix} 1 & b & z \\ 0 & 1 & -\bar{b} \\ 0 & 0 & 1 \end{bmatrix} : z, b \in E, z + \bar{z} = -b\bar{b} \right\} \simeq H(E),$$

the Heisenberg group attached to E over F . In particular, N is non-abelian, with commutator subgroup

$$[N, N] = \left\{ \begin{bmatrix} 1 & 0 & z \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right\} = Z,$$

the center of N . The maximal abelian subgroup of N is

$$N' = \left\{ \begin{bmatrix} 1 & b & z \\ 0 & 1 & -b \\ 0 & 0 & 1 \end{bmatrix} \in N : b \in F \right\}.$$

1.2

The irreducible representations of the Heisenberg group, and hence those of N , are well known:

(i) σ is 1-dimensional.

In this case, σ must be trivial on

$$Z = [N, N]$$

and define a character of N/Z . So

$$N/Z \simeq \left\{ \begin{bmatrix} 1 & a & 0 \\ 0 & 1 & -\bar{a} \\ 0 & 0 & 1 \end{bmatrix} \right\} \simeq E$$

implies σ corresponds to a character of E , i.e.

$$\sigma = \psi_N \left(\begin{bmatrix} 1 & a & z \\ 0 & 1 & -\bar{a} \\ 0 & 0 & 1 \end{bmatrix} \right) = \psi(\Im a)$$

with ψ a character of F .

(ii) σ is infinite-dimensional.

In this case (by the Stone-von Neumann uniqueness theorem), σ is completely determined by its “central” character. In particular, if

$$\sigma \left(\begin{bmatrix} 1 & 0 & z \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) = \psi(\Im z)I$$

for some (additive) character ψ of F , then

$$\sigma = \rho_\psi = \text{Ind}_{N'}^N \psi_{N'},$$

with $\psi_{N'}$ the character of (the maximal abelian subgroup) N' obtained by trivially extending ψ from Z to N' .

1.3

Definition. By a (generalized) Whittaker functional for (π, H_π) we understand N -map from N_π to some irreducible representation of (σ, L_σ) of N (possibly infinite dimensional).

1.4

Remark. The torus

$$T = \left\{ \begin{bmatrix} \delta & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & \bar{\delta}^{-1} \end{bmatrix} : \delta \in E^\times \right\}$$

acts by conjugation on N , taking

$$\begin{bmatrix} 1 & b & z \\ 0 & 1 & -\bar{b} \\ 0 & 0 & 1 \end{bmatrix} \quad \text{to} \quad \begin{bmatrix} 1 & \delta b & \delta \bar{\delta} z \\ 0 & 1 & -\bar{\delta} \bar{b} \\ 0 & 0 & 1 \end{bmatrix}.$$

So if ψ_N denotes the 1-dimensional representation of N corresponding to the fixed character of F as in 1.2 (i), Pontrygin duality for $E \simeq N/Z$ implies that any other 1-dimensional representation is trivial or of the form

$$\psi_N^\delta(n) = \psi_N \left(\begin{bmatrix} \delta & & \\ & 1 & \\ & & \bar{\delta}^{-1} \end{bmatrix} n \begin{bmatrix} \delta & & \\ & 1 & \\ & & \bar{\delta}^{-1} \end{bmatrix}^{-1} \right)$$

for some $\delta \in E^\times$.

1.5

If σ is a one-dimensional representation of N of the form ψ_N , a given irreducible admissible representation (π, H_π) need not possess a nontrivial ψ_N -Whittaker

functional \mathcal{L} . However, if it does, then by 1.4 it possesses a σ -Whittaker functional for any one-dimensional representation ψ_N^δ , given by the formula

$$\mathcal{L}^\delta(v) = \mathcal{L}\left(\pi\left(\begin{bmatrix} \delta & & \\ & 1 & \\ & & \bar{\delta}^{-1} \end{bmatrix}v\right)\right), \quad v \in H_\pi.$$

In this case, we call (π, H_π) non-degenerate. By a well-known theorem of Shalika and Gelfand-Kazhdan (cf. [Sha1]), the space of such σ -Whittaker functionals is one-dimensional. In particular, the corresponding Whittaker models

$$\mathcal{W}(\pi, \psi) = \{W(g) = \mathcal{L}(\pi(g)v) : v \in H_\pi\}$$

are unique.

1.6

In general, (π, H_π) is not non-degenerate, examples being provided by the Weil representations discussed in §??. Thus it is necessary to consider σ -Whittaker models for infinite dimensional σ as well. Such σ , however, are completely determined by their central character ψ_Z , so it is convenient to work with a slight thickening of N . More precisely, consider the stabilizer R in P of the central character ψ_Z of Z . Because $\begin{bmatrix} \delta & & \\ & \beta & \\ & & \bar{\delta}^{-1} \end{bmatrix}$ conjugates $\begin{bmatrix} 1 & 0 & z \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ to $\begin{bmatrix} 1 & 0 & \delta\bar{\delta}z \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$,

$$R = \left\{ \begin{bmatrix} \delta & * & * \\ 0 & \beta & * \\ 0 & 0 & \delta \end{bmatrix} \in P : \delta, \beta \in E^1 \right\} \simeq (E^1 \times E^1) \ltimes N.$$

In particular, each irreducible infinite dimensional representation ρ_ψ of N extends to a like representation ρ_ψ^α of R with α a character of $E^1 \times E^1$.

Theorem (Existence and Uniqueness of Generalized Whittaker Models: [PS3].) Any (π, H_π) possesses a ρ_ψ^α -Whittaker functional for some choice of ρ_ψ^α ; moreover, the space of such functionals is at most one dimensional.

We shall discuss this result in more detail in the global context of §??.

❖ Some Fourier Expansions and Hypercuspidality

Now F is a global field not of characteristic 2, and π is an automorphic cuspidal representation of $G(\mathbb{A})$ which we suppose realized in some subspace of cusp

forms H_π in $L_0^2(G(F)\backslash G(\mathbb{A}))$. To attach an L -function to π , it is useful to take forms f in H_π and examine their Fourier coefficients along the maximal unipotent subgroup N . When such coefficients are non-zero, π is non-degenerate, and we are led back to the local Whittaker models of 1.5; in this case, we can (and eventually do) introduce L -functions using Jacquet's generalization of the "Rankin-Selberg method".

On the other hand, if these Fourier coefficients represent zero, then π is hypercuspidal; in this case, looking at Fourier expansions along Z will bring us back to the generalized Whittaker models of 1.6, and ultimately allow us to introduce an L -function for π using the so-called "Shimura method".

Henceforth, let us fix a non-trivial character ψ of $F\backslash\mathbb{A}$, and define characters ψ_N and ψ_Z of $N = N(\mathbb{A})$ and $Z = Z(\mathbb{A})$ by

$$\psi_N \left(\begin{bmatrix} 1 & a & z \\ 0 & 1 & -\bar{a} \\ 0 & 0 & 1 \end{bmatrix} \right) = \psi(\Im a)$$

and

$$\psi_Z \left(\begin{bmatrix} 1 & 0 & z \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) = \psi(\Im z).$$

2.1

Fix f in H_π . To obtain a Fourier expansion of f "along N ", we introduce the familiar ψ -th coefficient

$$W_f^\psi(g) = \int_{N(F)\backslash N(\mathbb{A})} f(n g) \overline{\psi_N(n)} dn.$$

The transitivity of $T(\mathbb{A}) = \left\{ \begin{bmatrix} \delta & & \\ & 1 & \\ & & \bar{\delta}^{-1} \end{bmatrix} \right\}$ acting on $Z(\mathbb{A})\backslash N(\mathbb{A})$ implies - as in the local theory - that

$$\begin{aligned} W_f^{\psi^\delta}(g) &= \int_{N(F)\backslash N(\mathbb{A})} f(n g) \overline{\psi_N^\delta(n)} dn \\ &= \int_{N(F)\backslash N(\mathbb{A})} f(n g) \psi_N \left(\begin{bmatrix} \delta & & \\ & 1 & \\ & & \bar{\delta}^{-1} \end{bmatrix} n \begin{bmatrix} \delta & & \\ & 1 & \\ & & \bar{\delta}^{-1} \end{bmatrix}^{-1} \right) dn \\ &= W_f^\psi \left(\begin{bmatrix} \delta & & \\ & 1 & \\ & & \bar{\delta}^{-1} \end{bmatrix} g \right). \end{aligned}$$

In other words, knowing W_f^ψ determines $W_f^{\psi^\delta}$ for all $\psi^\delta, \delta \in E^\times$.

However, through $N(F) \backslash N(\mathbb{A})$ is compact, it is not abelian; to obtain a nice Fourier expansion, we must bring into play the compact abelian group $N(F)Z(\mathbb{A}) \backslash N(\mathbb{A})$.

2.2

We compute

$$\begin{aligned} W_f^\psi(g) &= \int_{N(F) \backslash N(\mathbb{A})} f(n g) \overline{\psi_N(n)} dn \\ &= \int_{N(F)Z(\mathbb{A}) \backslash N(\mathbb{A})} \int_{Z(F) \backslash Z(\mathbb{A})} f(n z g) dz \overline{\psi_N(n)} dn \\ &= \int_{N(F)Z(\mathbb{A}) \backslash N(\mathbb{A})} f_{00}(n g) \overline{\psi_N(n)} dn \end{aligned}$$

with

$$f_{00}(g) = \int_{Z(F) \backslash Z(\mathbb{A})} f(z g) dz \quad (1)$$

the constant term (in the Fourier expansion) of $f(z g)$ along Z .

Fix g in $G(\mathbb{A})$. As a function on the compact abelian group $N(F)Z(\mathbb{A}) \backslash N(\mathbb{A})$, $f_{00}(n g)$ has a Fourier expansion

$$f_{00}(g) = \sum_{\delta \in E^\times} W_F^{\psi^\delta}(g) + \int_{N(F)Z(\mathbb{A}) \backslash N(\mathbb{A})} f_{00}(n' g) dn'. \quad (2)$$

Indeed, the last paragraph says precisely that $W_f^\psi(g)$ is the ψ -th Fourier coefficient of $f_{00}(n g)$ along $Z \backslash N \simeq E$. Moreover, the constant term is actually zero since f cuspidal implies

$$\begin{aligned} \int_{N(F)Z(\mathbb{A}) \backslash N(\mathbb{A})} f_{00}(n' g) dn' &= \int_{N(F)Z(\mathbb{A}) \backslash N(\mathbb{A})} \int_{Z(F) \backslash Z(\mathbb{A})} f(z n' g) dz dn' \\ &= \int_{N(F) \backslash N(\mathbb{A})} f(n g) dn = 0. \end{aligned}$$

2.3

Let $\mathcal{W}(\pi, \psi)$ denote the space of ψ -th Fourier coefficients $W_f^\psi(g)$, $f \in H_\pi$.

Proposition 2.1. The vanishing or nonvanishing of $\mathcal{W}(\pi, \psi)$ is independent of ψ ; in particular, $\mathcal{W}(\pi, \psi) = 0$ if and only if

$$f_{00}(g) = 0 \quad \forall f \in H_\pi.$$

Proof. According to (1) and (2),

$$\begin{aligned} f_{00}(g) &= \sum_{\delta \in E^\times} W_f^{\psi^\delta}(g) \\ &= \sum_{\delta \in E^\times} W_f^\psi \left(\begin{bmatrix} \delta & & \\ & 1 & \\ & & \bar{\delta}^{-1} \end{bmatrix} g \right) \end{aligned} \quad (3)$$

with

$$W_f^\psi(g) = \int_{N(F)Z(\mathbb{A}) \backslash N(\mathbb{A})} f_{00}(ng) \overline{\psi_N(n)} dn.$$

□

2.4

Definition 2.2. We call (π, H_π) hypercuspidal if $f \in H_\pi$ implies $f_{00} = 0$.

Proposition 2.3. Let $L_{0,1}^2$ be the orthogonal complement in L_0^2 of all cusp forms. Then

- (i) $L_{0,1}^2$ has multiplicity 1.
- (ii) each $(\pi, H_\pi) \subset L_{0,1}^2$ is non-degenerate, and
- (iii) for any $f \in H_\pi \subset L_{0,1}^2$, the constant term

$$f_{00}(g) = \sum_{\delta \in E^\times} W_f^{\psi^\delta}(g)$$

completely determines f .

Proof. We start with (iii). Suppose f and f' are in H_π such that $f_{00} = f'_{00}$. Then $(f - f')_{00} = 0$ implies $f - f' = 0$ (by the hypothesis $H_\pi \in L_{0,1}^2$). This proves (iii). To prove (i) and (ii), suppose there exists $H'_\pi \subset L_{0,1}^2$ such that the right regular representation restricted to H'_π again realizes π . If $f \in H_\pi$ and $f' \in H'_\pi$, then

$$f_{00}(g) = \sum_{\delta \in E^\times} W_f^\psi \left(\begin{bmatrix} \delta & & \\ & 1 & \\ & & \bar{\delta}^{-1} \end{bmatrix} g \right) \quad (4)$$

and

$$f'_{00}(g) = \sum_{\delta \in E^\times} W_{f'}^\psi \left(\begin{bmatrix} \delta & & \\ & 1 & \\ & & \bar{\delta}^{-1} \end{bmatrix} g \right).$$

Note that each W_f^ψ (or $W_{f'}^\psi$) satisfies the condition $W_f^\psi(n g) = \psi(n) W_f^\psi(g)$, $n \in N$, i.e. the spaces (W_f^ψ) and $(W_{f'}^\psi)$ afford Whittaker models for π . But by §2.3 these spaces are nonzero (which proves (ii)) and by the uniqueness of Whittaker models quoted in §1.5, these spaces coincide. Thus by (4), the spaces (f_{00}) and (f'_{00}) coincide; by (iii) the spaces $H_\pi = (f)$ and $H'_\pi = (f')$ also coincide, thereby proving (i). \square

2.5

- Remark.** (i) It is conjectured (c.f. [Flicker]) that multiplicity one holds for the entire space of cusp forms; however, at the present time, we can prove this only for $L_{0,1}^2$.
- (ii) Hypercusps do exist; again, the examples are provided by the Weil representation discussed in §??.
- (iii) Although $\mathcal{W}(\pi, \psi) \neq \{0\}$ implies π non-degenerate (in the sense that an abstract functional exists), the converse is not clear. Indeed, the work of [Wald] indicates that characterizing the nonvanishing of a space of Fourier coefficients is a delicate matter.

❖ L -functions à la Rankin–Selberg–Jacquet

We are now ready to attach (global) L -functions to (non-degenerate) cuspidal representations π of $G(\mathbb{A})$. The method used goes back to [Rankin] and [Selberg] who used it to analytically continue the convolution of Dirichlet series corresponding to classical holomorphic modular forms. The reformulation of their construction in the language of representation theory was carried out in detail by [Jacquet], for $GL_2 \times GL_2$, and by [PS 1] in general (but without details or explicit computation). In this Section (and the next), we carry out the construction for $G = U_{2,1}$.

3.1

Recall V is a 3-dimensional vector space over E/F and $(-, -)_V$ is a hermitian form on V whose matrix with respect to the basis $\{\ell_{-1}, \ell_0, \ell_1\}$ is $\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$.