

=E16+\$F\$13

G17

=COUNTIFS(\$B\$13:\$B\$32, ">="&\$E17,\$B\$13:\$B\$32, "<="&\$F17)

	B	C	D	E	F	G	H	I	J	K	
1	Numerical variables. Frequency distribution table										
2											
3	Background	You are given a dataset.									
4	Task 1	Given that we want to divide the numbers into 6 intervals of equal width, calculate that interval width. Round up to the nearest whole number, bigger than the result that you obtain									
5	Task 2	Create a frequency distribution table that shows									
6		1. The intervals									
7		2. The absolute frequency of each interval									
8		3. The relative frequency of each interval									
9	Task 3	Repeat task 1 and 2, but this time, use the exact interval width. Don't round up to the nearest whole number, that is.									
10											
11											
12	Dataset		# Intervals	6		=E16+\$F\$13					
13	8		Interval width	46							
14	30										
15	30										
16	50										
17	86										
18	94										
19	102										
20	110										
21	169										
22	170										
23	176										
24	236										
25	240										
26	241		Using exact width								
27	242		# Intervals	6							
28	255		Interval width	45.67							
29	262										
30	276										
31	279										
32	282										
33											
34	count										
35	20										
36											

interval start	interval end	frequency	relative frequency
8	54	4	0.2
54	100	2	0.1
100	146	2	0.1
146	192	3	0.15
192	238	1	0.05
238	284	8	0.4

interval start	interval end	frequency	relative frequency
8.00	53.67	4	0.2
53.67	99.33	2	0.1
99.33	145.00	2	0.1
145.00	190.67	3	0.15
190.67	236.33	1	0.05
236.33	282.00	8	0.4

Frequency distribution table