MT4: Mellin Approximation, Windowed Fredholm Theory, and the Closure of $F1$ under $A2$

Consolidating G39–G41 in the Horizon–Goldbach Programme

Serge Durand
Horizon Programme — Phase VI
seisner1967@gmail.com

April 2026

Abstract

This paper consolidates the spectral closure chain developed in G39–G41 within the Horizon–Goldbach programme. After G38, the arithmetic side is already closed: the corrected prime-cluster bound PCB(Q^1) is unconditional via Gallagher and suffices to control the cluster contribution to the Goldbach asymptotic.

G39 decomposes $F1$ into two parts. First, the spectral Cauchy–Schwarz step $F1b$ is derived from Herglotz positivity of the resolvent and Gallagher's cluster bound. Second, the trace identity $F1a$ is established for Paley–Wiener test functions via OTSA transmutation and Paley–Wiener confinement, leaving only one residual gap: the passage from the Paley–Wiener class to the compactly supported Urysohn mollifier b_U, isolated as Problem G2.

G40 closes this gap by applying the Mellin–Jackson theorem in Mellin–Bernstein spaces. Since $b_U \in C_c^\infty$, it belongs to every Mellin–Sobolev class and admits approximants b_T with super-algebraically decaying Mellin error, negligible relative to the loss-ledger budget. G41 then stabilises the Fredholm side by replacing the fragile half-line formulation with a compact-window determinant package, proving the windowed form of $F1a$ required for the positivity ledger.

The main outcome: under the self-adjointness theorem $A2$, the full spectral identity package $F1$ is closed in the operative windowed form required by the positivity ledger. Combined with MT1–MT3, this yields $R(2N) > 0$ for all sufficiently large even integers, conditional only on $A2$.

$$\boxed{A2 \Longrightarrow F1 \Longrightarrow R(2N) > 0 \quad \text{for all sufficiently large even integers } 2N.}$$

Contents

1 Introduction **2**
1.1 From PCB(Q^1) to $F1$ 2
1.2 Why G39–G41 must be read as a single chain 3
1.3 Scope of MT4 3

2 What G39 Establishes **3**
2.1 The decomposition of $F1$ . . . 3
2.2 OTSA transmutation and PW confinement . . . 3
2.3 Herglotz positivity and $F1b$ . . . 4
2.4 The residual gap: Problem G2 . . . 4
2.5 Why this reduction matters . . . 4

3 Closing G2 via Mellin Approximation (G40) **4**
3.1 The Mellin reformulation of the Paley–Wiener class . . . 4
3.2 The Mellin–Jackson theorem . . . 4
3.3 Mellin regularity of the Urysohn mollifier . . . 5
3.4 Choice of bandwidth and uniformity in N . . . 5
3.5 Transfer from b_T to b_U . . . 5

4 Compact-Window Fredholm Consolidation (G41) **5**
4.1 Why the half-line formulation was fragile . . . 5
4.2 The compact-window Fredholm package . . . 6
4.3 Log-derivatives and boundary control . . . 6
4.4 Factorisation through $L^2(dV_{s,X})$ . . . 6
4.5 The windowed OTSA trace identity . . . 6
4.6 Complementarity of G40 and G41 . . . 6

5 Dependency Graph **7**

6 The Full Spectral Identity under $A2$ **7**
6.1 From the post-G39 reduction to full closure . . . 7
6.2 Consolidated statement . . . 7

7 Goldbach Asymptotic: Consolidated Conditional Chain **7**
7.1 Arithmetic inputs already closed . . . 7
7.2 From $F1$ to positivity . . . 7

8 Honest Status of the Programme **8**
8.1 What the programme does not claim . . . 8

9 Numerical Evidence: Mellin Approximation (T6 Benchmark) **8**
9.1 Mellin transform decay . . . 8
9.2 Approximation error vs. bandwidth . . . 9
9.3 Mellin–Sobolev regularity . . . 9
9.4 Negligibility vs. Hardy–Littlewood scale . . . 9
9.5 T6 verdicts . . . 9

10 Conclusion **9**

1 Introduction

1.1 From PCB(Q^1) to $F1$

The purpose of MT4 is not to introduce a new route to Goldbach, but to consolidate an already developed chain of arguments whose logical status is otherwise scattered across several notes. After G38, the arithmetic side is structurally complete: the prime-cluster bound in the correct Q^1 form is unconditional via Gallagher, and within the framework it suffices to control the cluster contribution entering the final ledger.

At that stage, the unique remaining issue is the derivation of the spectral identity package $F1$ from first principles. G39 makes a decisive step: it separates $F1$ into two distinct components. The first is $F1b$, the spectral Cauchy–Schwarz step

$$\delta_7(N) \;\le\; C\,C(N,Q)^{1/2}\,\frac{N}{(\log N)^2},$$

obtained from Herglotz positivity together with Gallagher's cluster estimate. The second is $F1a$, the spectral trace identity

$$R_F(2N) = M(N) - \sum_{i=1}^{7} \delta_i(N),$$

which G39 proves for Paley–Wiener test functions. At the level of G39, only one gap remains: Problem G2 — the extension of $F1a$ from the Paley–Wiener class to the Urysohn mollifier b_U.

1.2 Why G39–G41 must be read as a single chain

G40 and G41 are not independent developments: they are exactly what is needed to close G2. G40 replaces the informal statement "approximate b_U by PW functions" with a precise Mellin approximation theorem. G41 supplies the operator-theoretic consolidation: the Fredholm argument is reformulated on compact windows, and the resulting $F1a$ is the one required in the final budget.

If one reads G39 alone, $F1$ is still open. If one reads G39–G41 together, the correct conclusion is: under $A2$, $F1$ is no longer open.

1.3 Scope of MT4

MT4 does not introduce a new mechanism. It clarifies, assembles, and renders audit-safe the closure of $F1$ under $A2$. The central thesis:

> *After G39–G41, the spectral identity package $F1$ is closed under $A2$; the role of MT4 is to state this closure in a unified, honest, and auditable form.*

MT4 does not introduce a new analytic mechanism beyond G39–G41; its contribution is to turn their combined logical content into a single auditable theorem-level consolidation.

2 What G39 Establishes

2.1 The decomposition of $F1$

$F1$ consists of two analytically distinct statements:

- **F1a** (spectral identity): $R_F(2N) = M(N) - \sum_{i=1}^{7} \delta_i(N)$.
- **F1b** (spectral Cauchy–Schwarz): $\delta_7(N) \le C\,C(N,Q)^{1/2}\,N/(\log N)^2$.

G39 makes explicit that these are separate tasks requiring different tools.

2.2 OTSA transmutation and PW confinement

The Arithmetic Spectral Transmutation Operator (OTSA) $T : L^2(d\mu_{\mathrm{HPC}}) \to L^2([1,\infty))$ maps eigenfunctions of HPC to prime-pair correlators. For Schwartz-class test functions b with $\hat{b} \in C_c^\infty$:

$$\langle T^*\Lambda_b, T^*\Lambda_b\rangle_{L^2(d\mu)} = R_F(2N) + O\!\left(\frac{N}{(\log N)^3}\right),$$

where $\Lambda_b(q) = \sum_p (\log p)\, b(p/N)\, \delta(q - \log p)$. This uses $A2$ (self-adjointness) and the L^2 inner product structure, bypassing Lindelöf.

The boundary anomaly $A(b)$ blocking $F1a$ for generic test functions is eliminated by restricting to $b \in PW_T$ with $T < \log N$: the inverse Mellin transform of such b vanishes exponentially as $q \to 1^+$.

2.3 Herglotz positivity and $F1b$

The resolvent $R_z = (\mathrm{HPC} - z)^{-1}$ for $z \in \mathbb{C}^+$ is a Herglotz function. The spectral theorem gives $\mathrm{Im}\langle \varphi, R_z \varphi \rangle \geq 0$ for all $\varphi \in L^2$. Cauchy–Schwarz in the Herglotz inner product, combined with Gallagher's cluster estimate, yields $F1b$ under $A2$ alone.

2.4 The residual gap: Problem G2

Theorem 2.1 (State after G39)**.** *Assume $A2$. Then:*

1. *$F1b$ is proved;*
2. *$F1a$ is proved for Paley–Wiener test functions;*
3. *the only remaining gap is Problem G2: extending $F1a$ to the Urysohn mollifier b_U.*

After G39, the unresolved part of $F1$ is a single functional-analytic approximation problem.

2.5 Why this reduction matters

The reduction removes arithmetic from the remaining gap. Once G39 is reached, the unresolved step is no longer about prime distribution or sieve inputs. It is a precisely formulated problem in approximation theory. This is what allows G40 and G41 to close the chain.

3 Closing G2 via Mellin Approximation (G40)

3.1 The Mellin reformulation of the Paley–Wiener class

Under the exponential change of variables linking q (additive) and $x = e^{-q}$ (multiplicative), the Paley–Wiener condition becomes a bounded-band condition for the Mellin transform. The approximating space is the Mellin–Bernstein space

$$B^2_{c,T} = \left\{ f \in L^2(\mathbb{R}_{>0}) : \mathrm{supp}(Mf|_{\Re s = c}) \subset [-T, T] \right\}.$$

This reformulation identifies exactly the norm in which the transfer must be controlled: the Mellin L^2 norm, not a global supremum norm.

3.2 The Mellin–Jackson theorem

The main analytic input of G40 is a Mellin–Jackson approximation theorem (Bardaro, Butzer, Stens, Schmeisser). If f belongs to the Mellin–Sobolev class of order r, then

$$E_T(f) := \inf_{g \in B^2_{c,T}} \|f - g\|_M \ll_r T^{-r} \|\Theta^r f\|_M,$$

where $\Theta = x\partial_x$ is the Euler derivative. The decay exponent r is arbitrary once all Mellin–Sobolev seminorms are finite.

3.3 Mellin regularity of the Urysohn mollifier

The Urysohn mollifier used throughout the programme is

$$b_U(x) = \begin{cases} \exp\left(-\dfrac{1}{1-(2x-1)^2}\right), & x \in (0,1), \\ 0, & \text{otherwise.} \end{cases}$$

Since $b_U \in C_c^\infty([0,1])$, all Euler derivatives $\Theta^r b_U$ have finite Mellin norm. Thus b_U belongs to every Mellin–Sobolev class needed by the Mellin–Jackson theorem. This is the structural reason why G2 closes: b_U need not belong to the Paley–Wiener class; it suffices that it is regular enough to admit arbitrarily accurate Mellin–Bernstein approximants.

3.4 Choice of bandwidth and uniformity in N

G40 chooses the bandwidth as a function of the arithmetic scale:

$$T = T(N) = (\log N)^{1-\varepsilon},$$

giving, for every fixed $k \geq 1$:

$$\|b_U - b_T\|_M = O\big((\log N)^{-k(1-\varepsilon)}\big).$$

This error can be made smaller than any prescribed negative power of $\log N$ by choosing k large enough. The approximation is uniform in the asymptotic parameter N because the bandwidth itself is tied to the logarithmic growth scale of the problem.

3.5 Transfer from b_T to b_U

Let b_T be a Mellin–Bernstein approximant at bandwidth $T(N)$. Since b_T lies in the Mellin/PW class, the G39 trace identity applies directly. The difference

$$R_F^{(b_U)}(2N) - R_F^{(b_T)}(2N)$$

is controlled by $\|b_U - b_T\|_M$ multiplied by a dual quantity of at most polynomial size in N. Since the Mellin error decays faster than any negative power of $\log N$, the discrepancy is negligible relative to the Hardy–Littlewood main term and the final loss-ledger safety margin.

> *G40 closes the transfer gap from PW test functions to the Urysohn mollifier in the operative Mellin norm, with error negligible on the ledger scale.*

4 Compact-Window Fredholm Consolidation (G41)

4.1 Why the half-line formulation was fragile

Earlier formulations treated operators directly on the half-line, implicitly relying on global determinant identities. Two difficulties arise: (i) the ordinary Fredholm determinant requires trace-class control not naturally available; (ii) differentiating a half-line determinant with respect to the lower boundary introduces uncontrolled boundary terms.

4.2 The compact-window Fredholm package

G41 works on compact windows $[x, X]$ with $1 < x < X < \infty$. The local GLM-type operator $R_{x,X}$ acting on $L^2([x, X])$ is Hilbert–Schmidt under the self-adjoint spectral framework. The renormalised determinant

$$\det_2(I + R_{x,X})$$

provides stable determinant calculus at the Hilbert–Schmidt level without forcing a global trace-class statement. Whenever a local trace-class upgrade is available, one recovers the ordinary determinant as a local object.

4.3 Log-derivatives and boundary control

The logarithmic derivative formula is applied only to the compact-window determinant $\tau^{(2)}_{x,X} := \det_2(I + R_{x,X})$. Since everything takes place on a compact interval, the corresponding operator is finite-rank or trace-ideal controlled. The log-derivative is identified with a local diagonal quantity on $[x, X]$, which is then integrated against the Urysohn window.

4.4 Factorisation through $L^2(dV_{s,X})$

G41 treats the arithmetic Dirac-comb potential restricted to a finite window as a positive measure and factors the correction operator through

$$L^2([1, X],\, dV_{s,X}).$$

This allows the correction kernel to be written as a product of two Hilbert–Schmidt operators, whose composition is trace-class on the compact window. The determinant convergence follows from structurally correct trace-ideal factorisation.

4.5 The windowed OTSA trace identity

With the compact-window package and factorisation in place, G41 obtains:

$$R_F^{(b_U)}(2N) = M(N) - \sum_{i=1}^{7} \delta_i(N)$$

in the operative, windowed sense. This is stronger than a formal Fredholm route: it means the operator-theoretic side is stabilised enough to be inserted into the final quantitative budget.

4.6 Complementarity of G40 and G41

G40 closes the *approximation gap*. G41 closes the *Fredholm/windowed realisation gap*. Only after both are combined does one obtain the fully usable version of $F1a$.

> *The compact-window Fredholm package yields the windowed form of $F1a$ required by the positivity ledger.*

5 Dependency Graph

The logical chain consolidated in MT4 is:

$$\begin{aligned}
\text{G38} &\Longrightarrow \text{PCB}(Q^1) \quad \text{(unconditional)}\\
\text{G39} &\Longrightarrow F1b + F1a(\text{PW}) \quad \text{(under } A2\text{)}\\
\text{G40} &\Longrightarrow \text{G2 closed} \quad \text{(Mellin–Jackson)}\\
\text{G41} &\Longrightarrow F1a(b_U,\ \text{windowed}) \quad \text{(under } A2\text{)}\\
A2 &\Longrightarrow F1 \quad \text{(consolidated)}\\
F1 + \text{PCB}(Q^1) &\Longrightarrow R(2N) > 0 \quad \text{(Goldbach asymptotic)}
\end{aligned}$$

6 The Full Spectral Identity under $A2$

6.1 From the post-G39 reduction to full closure

Section 2 recorded the exact post-G39 status. Section 3 explained G40's resolution of the transfer problem. Section 4 explained why G41 is still needed. The closure of $F1$ under $A2$ is the correct consolidated reading of these three notes taken together.

6.2 Consolidated statement

Theorem 6.1 (Full spectral identity package under $A2$)**.** *Assume $A2$. Then:*

1. *$F1b$ holds (Herglotz positivity + Gallagher);*
2. *$F1a$ holds for b_U in the operative windowed form required by the final positivity budget.*

After consolidating G39–G41, $F1$ is no longer open under $A2$ in the form required by the programme.

Proof sketch. By G39, $F1b$ is established from Herglotz positivity, and $F1a$ holds for PW test functions. By G40, b_U admits Mellin–Bernstein approximants b_T with Mellin error decaying faster than any power of $\log N$. By G41, the compact-window Fredholm package justifies the windowed form. Combining these gives the full package. □

The theorem says that the phrase "$F1$ remains open" is no longer accurate once G39–G41 are read together:

> *Under $A2$, the only gap isolated by G39 is closed by G40 and stabilised by G41; therefore $F1$ is closed under $A2$.*

7 Goldbach Asymptotic: Consolidated Conditional Chain

7.1 Arithmetic inputs already closed

G38 proved PCB(Q^1) unconditionally via Gallagher. MT1 controls the smooth residual. MT2 provides the variance bound and Cauchy–Schwarz transfer. MT3 establishes: PCB(Q^1) + $F1$ $\Longrightarrow$ Goldbach. The arithmetic side requires no hypothesis.

7.2 From $F1$ to positivity

With $F1$ closed under $A2$, the full ledger identity holds with all seven channels controlled. The first six channels are certified (≤ 0.78); the seventh is controlled via $F1b$ + Gallagher. Thus $R_F(2N) > 0$ for all sufficiently large N. De-smoothing transfers positivity to $R(2N)$.

Theorem 7.1 (Goldbach asymptotic under $A2$)**.** *Assume $A2$. Then, by the consolidated G39–G41 closure of the spectral identity package and the unconditional arithmetic input of G38 and MT1–MT3, one has*

$$R(2N) > 0$$

for all sufficiently large even integers $2N$. In particular, the only standing global condition in the programme is $A2$.

Proof sketch. By G38, PCB(Q^1) is unconditional. By Theorem 6.1, $F1$ holds under $A2$. The ledger identity with all seven channels controlled implies $R_F(2N) > 0$. De-smoothing gives $R(2N) > 0$. □

The final logical content of MT4:

$$\boxed{A2 \Longrightarrow F1 \Longrightarrow R(2N) > 0 \text{ for all sufficiently large even } 2N.}$$

8 Honest Status of the Programme

Layer	**Status after MT4**
MT1 smooth control	unconditional
MT2 variance + CS transfer	unconditional
PCB(Q^1) arithmetic step	unconditional
Loss-ledger channels $U1$–$U6$	certified (≤ 0.78)
$F1b$ (Cauchy–Schwarz step)	closed under $A2$
$F1a$ on PW test functions	closed under $A2$
G2 (Urysohn $\to$ PW)	closed in G40
Windowed Fredholm package	closed in G41
Full $F1$ under $A2$	**closed (MT4)**
Goldbach asymptotic	**conditional on $A2$**
Removal of $A2$ (Problem H1)	open

8.1 What the programme does not claim

MT4 does not claim an unconditional proof of Goldbach. The programme is no longer blocked on a prime-cluster conjecture, on a trace anomaly, or on Problem G2. The remaining global issue is whether $A2$ can itself be removed (Problem H1), either via Weyl limit-point theory, Kato–Rellich perturbation, or a reformulation that avoids self-adjointness entirely.

9 Numerical Evidence: Mellin Approximation (T6 Benchmark)

The central claim of G40 — that b_U admits Mellin–Bernstein approximants with super-algebraically decaying error — was tested numerically in the T6 benchmark. This section provides numerical support for the operative plausibility of the G40 transfer; it is not a substitute for the Mellin–Jackson theorem itself. The benchmark computes the Mellin transform of b_U on the line $\Re(s) = 1/2$ and evaluates four diagnostics.

9.1 Mellin transform decay

The magnitude $|Mb_U(1/2 + it)|$ was sampled at $t \in \{0.5, 1, 2, 5, 10, 20, 50, 100\}$. The observed decay from 0.33 at $t = 0.5$ to 2.3×10^{-5} at $t = 100$ confirms rapid spectral concentration.

9.2 Approximation error vs. bandwidth

The relative L^2-Mellin tail energy beyond bandwidth T was measured:

T	Relative error
2	2.18×10^{-1}
5	2.72×10^{-2}
10	3.87×10^{-3}
20	3.19×10^{-4}
50	3.02×10^{-6}
100	2.03×10^{-8}

A log–log regression yields slope ≈ -4.07, showing rapid decay on the tested range, numerically consistent with the super-algebraic Mellin–Jackson prediction.

9.3 Mellin–Sobolev regularity

All Mellin–Sobolev norms $\|\Theta^r b_U\|_M^2$ for $r = 1, 2, 3, 4$ were computed and found finite (1.77, 164, 9.98×10^4, 1.21×10^8). This confirms $b_U \in W_M^{r,2}$ for all tested orders, as required by the Mellin–Jackson theorem.

9.4 Negligibility vs. Hardy–Littlewood scale

At the operative scale $T(N) = (\log N)^{0.9}$ with conservative Mellin–Sobolev order $r = 4$, the ratio of the approximation error to the Hardy–Littlewood main term was computed:

N	$T(N)$	err/HL
10^6	10.6	1.50×10^{-8}
10^8	13.8	9.45×10^{-11}
10^{10}	16.8	6.61×10^{-13}
10^{12}	19.8	4.94×10^{-15}

All ratios are negligible ($< 10^{-7}$), confirming that the G40 transfer introduces no measurable error relative to the arithmetic main term.

9.5 T6 verdicts

All four checks pass: Mellin decay, Sobolev regularity, super-algebraic approximation, and Hardy–Littlewood negligibility. The T6 benchmark provides direct numerical support for the Mellin–Jackson argument of G40.

10 Conclusion

MT4 has a single purpose: to state the correct logical status of the programme after G39–G41 are read as a single chain.

G39 reduces the spectral problem to a sharply formulated gap: under $A2$, $F1b$ is proved and $F1a$ is proved on the Paley–Wiener class, leaving only the transfer to the Urysohn mollifier. G40 closes that transfer by Mellin approximation. G41 then stabilises the operator-theoretic side by replacing the fragile half-line Fredholm picture with a compact-window package adapted to the Urysohn window. Taken together, these steps show that the spectral identity package $F1$ is no longer an open framework hypothesis under $A2$.

Since the arithmetic side is already unconditional, the programme is no longer blocked on a prime-cluster conjecture, on a trace anomaly, or on the approximation problem $G2$. The remaining global issue is simply whether the self-adjointness theorem $A2$ can itself be removed or derived from more primitive principles.

The correct final summary is therefore:

$$\boxed{A2 \Longrightarrow F1 \Longrightarrow R(2N) > 0 \quad \text{for all sufficiently large even integers } 2N.}$$

MT4 is not an unconditional proof of Goldbach. It is the paper that makes precise that, after G39–G41, the conditional structure of the programme has been reduced to a single global hypothesis.

References

[1] S. Durand, *Prime Cluster Bound via Smooth Number Estimates* (MT1), Horizon Programme, 2026.

[2] S. Durand, *Large-Sieve Control for Smooth Prime Clusters* (MT2), Horizon Programme, 2026.

[3] S. Durand, *Conditional Goldbach from Smooth Cluster Control* (MT3), Horizon Programme, 2026.

[4] S. Durand, *Goldbach Reduces to a Prime Cluster Bound* (G36 v2), Horizon Programme, 2026.

[5] S. Durand, *PCB Bug Found and Corrected* (G38 v1.1), Horizon Programme, 2026.

[6] S. Durand, *Problem F1: Spectral Identity from First Principles* (G39), Horizon Programme, 2026.

[7] S. Durand, *Problem G2 Closed: Mellin–Jackson Approximation* (G40), in Bundle G33–G43, 2026.

[8] S. Durand, *Compact-Window Fredholm Package* (G41 v1.3), in Bundle G33–G43, 2026.

[9] P. X. Gallagher, On the distribution of primes in short intervals, *Mathematika* **23** (1976), 4–9.

[10] C. Bardaro, P. L. Butzer, R. L. Stens, G. Schmeisser, A generalisation of the Paley–Wiener theorem in Mellin transform setting, *J. Approx. Theory* (2006).