

1.


$$\begin{aligned}
 p(x) &= \int_{-\infty}^{+\infty} p(x, \tau^2) d\tau^2 \\
 &= \int_{-\infty}^{+\infty} p(x|\tau^2) p(\tau^2) d\tau^2 \\
 &= \int_{-\infty}^{+\infty} \frac{1}{\sqrt{2\pi}} (\tau^2)^{1/2} \exp\left\{-\frac{1}{2}\tau^2 x^2\right\} \frac{1}{\Gamma\left(\frac{\nu}{2}\right) \left(\frac{2}{\nu}\right)^{\frac{\nu}{2}}} (\tau^2)^{\frac{\nu}{2}-1} \exp\left\{-\frac{\nu}{2}\tau^2\right\} d\tau^2 \\
 &= \frac{1}{\sqrt{2\pi}} \times \frac{1}{\Gamma\left(\frac{\nu}{2}\right) \left(\frac{2}{\nu}\right)^{\frac{\nu}{2}}} \int_{-\infty}^{+\infty} (\tau^2)^{1/2} \exp\left\{-\frac{1}{2}\tau^2 x^2\right\} (\tau^2)^{\frac{\nu}{2}-1} \exp\left\{-\frac{\nu}{2}\tau^2\right\} d\tau^2 \\
 &= \frac{1}{\sqrt{2\pi}} \times \frac{1}{\Gamma\left(\frac{\nu}{2}\right) \left(\frac{2}{\nu}\right)^{\frac{\nu}{2}}} \int_{-\infty}^{+\infty} (\tau^2)^{\frac{\nu+1}{2}-1} \exp\left\{-\left(\frac{1}{2}x^2 + \frac{\nu}{2}\right)\tau^2\right\} d\tau^2 \\
 &= \frac{1}{\sqrt{2\pi}} \times \frac{1}{\Gamma\left(\frac{\nu}{2}\right) \left(\frac{2}{\nu}\right)^{\frac{\nu}{2}}} \Gamma\left(\frac{\nu+1}{2}\right) \left(\frac{1}{\frac{1}{2}x^2 + \frac{\nu}{2}}\right)^{\frac{\nu+1}{2}} \\
 &= \frac{\left(\frac{2}{\nu}\right)^{\frac{1}{2}}}{\sqrt{2\pi}} \times \frac{\Gamma\left(\frac{\nu+1}{2}\right) \left(\frac{1}{\frac{1}{2}x^2 + \frac{\nu}{2}}\right)^{\frac{\nu+1}{2}}}{\Gamma\left(\frac{\nu}{2}\right) \left(\frac{2}{\nu}\right)^{\frac{\nu+1}{2}}} \\
 &= \frac{1}{\sqrt{\nu\pi}} \times \frac{\Gamma\left(\frac{\nu+1}{2}\right)}{\Gamma\left(\frac{\nu}{2}\right)} \left(\frac{x^2}{\nu} + 1\right)^{-\frac{\nu+1}{2}}
 \end{aligned}$$

2. When $\nu=1$,

$$f(x) = \frac{1}{\sqrt{\pi}} \times \frac{\Gamma(1)}{\Gamma\left(\frac{1}{2}\right)} (x^2 + 1)^{-1} \sim t_1$$

$$\text{or } f(x) = \frac{1}{\pi(x^2 + 1)} \sim \text{Cauchy Dis.}$$

The figure is as below.



Compute 2.5% & 97.5% percentile points of the distribution:

The theoretical values: -12.7062 , 12.7062

The random samples values: -12.71138, 11.33432

They are very close to each other.

3. p-value for ks test is 0.3592. It denotes that x may have a same distribution as $t(1)$ distribution.

4.

The central limit theory does not hold for x when $\nu = 1$, for the

distribution of x will be a Cauchy Distribution, whose first and second moment don't converge.

When $\nu=2$, The central limit theory does not hold either. Because the variance of t_2 distribution does not converge.

The theory holds when $\nu=3$.

#####R Code#####

#2#

```
tau2<-rgamma(10000,shape=.5,rate=.5)
```

```
x.giventau2<-rnorm(10000,0,1/sqrt(tau2))
```

```
hist( x.giventau2)
```

```
qt(c(0.025,0.975),1)
```

```
quantile(x.giventau2,probs=0.0250)
```

```
quantile(x.giventau2,probs=0.0251)
```

```
(quantile(x.giventau2,probs=0.975)+quantile(x.giventau2,probs=  
0.9751))/2
```

```
quantile(x.giventau2,probs=0.9750)
```

```
quantile(x.giventau2,probs=0.9751)
```

```
(quantile(x.giventau2,probs=0.0250)+quantile(x.giventau2,probs  
=0.0251))/2
```

```
#3#
```

```
ks.test(c(x.giventau2),"pt",1)
```