

Definition.

γ

Definition.

$$\gamma : (\alpha, \beta)$$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector*

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

Show that $\dot{\gamma}(t)$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

Show that $\dot{\gamma}(t) = s'(t)$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

Show that $\dot{\gamma}(t) = s'(t)\mathbf{T}(t)$.

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

Show that $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \mathbf{T}(t)$.

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

Show that $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \underbrace{\mathbf{T}(t)}_{\text{”Change in direction”}} .$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

Show that $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \underbrace{\mathbf{T}(t)}_{\text{”Change in direction”}} .$

Solution. $\dot{\gamma}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} \times \square$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

Show that $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \underbrace{\mathbf{T}(t)}_{\text{”Change in direction”}}$.

Solution. $\dot{\gamma}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} \times \|\dot{\gamma}(t)\|$ □

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

Show that $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \underbrace{\mathbf{T}(t)}_{\text{”Change in direction”}}$.

Solution. $\dot{\gamma}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} \times \|\dot{\gamma}(t)\| = \mathbf{T}(t) \times \square$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

Show that $\dot{\gamma}(t) = \underbrace{s'(t)}_{\text{”Change in distance”}} \times \underbrace{\mathbf{T}(t)}_{\text{”Change in direction”}}$.

Solution. $\dot{\gamma}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} \times \|\dot{\gamma}(t)\| = \mathbf{T}(t) \times s'(t) \quad \square$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

γ

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta)$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma}$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta})$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t})$$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$



Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\dot{\tilde{\gamma}}(\tilde{t})$$



Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t}))\end{aligned}$$



Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If $g(f(t)) = t$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If $g(f(t)) = t$, then

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If $g(f(t)) = t$, then

$$g'(f(t))f'(t) = 1,$$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore,



Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

□

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

□

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.
 $\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.
 $\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.
Show that,
 $\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,
If $g(f(t)) = t$, then
 $g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.
 $\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.
 $\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.
Show that,
 $\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned} \dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ?? \end{aligned}$$

□

Recall,
If $g(f(t)) = t$, then
 $g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking $f(t) = s(t)$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking $f(t) = s(t)$ and $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)}$$

□

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

□

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking $f(t) = s(t)$ and $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\gamma'(t)\|}$$

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking $f(t) = s(t)$ and $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\gamma'(t)\|}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))}$$

□

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking $f(t) = s(t)$ and $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\dot{\gamma}(t)\|}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|}$$

□

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned}\dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \times ??\end{aligned}$$

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking $f(t) = s(t)$ and $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\dot{\gamma}(t)\|}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|}$$

□

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned} \dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|} \end{aligned}$$

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking $f(t) = s(t)$ and $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\gamma'(t)\|}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|}$$

□

Definition.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\mathbf{T}(t) = \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$ is the *unit tangent vector* at t .

Exercise.

$\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$ a regular parametrization.

$\tilde{\gamma} : (\tilde{\alpha}, \tilde{\beta}) \rightarrow \mathbb{R}^2$ a unit speed reparametrization.

Show that,

$$\dot{\tilde{\gamma}}(\tilde{t}) = \mathbf{T}(s^{-1}(\tilde{t}))$$

Solution. $\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$

$$\begin{aligned} \dot{\tilde{\gamma}}(\tilde{t}) &= \dot{\gamma}(s^{-1}(\tilde{t}))(s^{-1}(\tilde{t}))' \\ &= \dot{\gamma}(s^{-1}(\tilde{t})) \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|} \\ &= \mathbf{T}(s^{-1}(\tilde{t})) \end{aligned}$$

□

Recall,

If $g(f(t)) = t$, then

$g'(f(t))f'(t) = 1$, therefore, if $f'(t) \neq 0$ for any t ,

$$g'(f(t)) = \frac{1}{f'(t)}$$

Taking $f(t) = s(t)$ and $g(t) = s^{-1}(t)$

$$(s^{-1})'(s(t)) = \frac{1}{s'(t)} = \frac{1}{\|\dot{\gamma}(t)\|}$$

$$(s^{-1})'(\tilde{t}) = \frac{1}{s'(s^{-1}(\tilde{t}))} = \frac{1}{\|\dot{\gamma}(s^{-1}(\tilde{t}))\|}$$

Exercise.

If $\|\dot{\gamma}(t)\|$

Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c ,

Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$

Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant.

Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function

Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$s(t) =$$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$s(t) = \int$$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$s(t) = \int_{t_0}$$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$s(t) = \int_{t_0}^t$$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$s(t) = \int_{t_0}^t \|\dot{\gamma}(t)\|$$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$s(t) = \int_{t_0}^t \|\dot{\gamma}(t)\| dt$$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$\begin{aligned} s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \end{aligned}$$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

So, $s(t)/c$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

So, $s(t)/c = t - t_0$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

So, $s(t)/c = t - t_0$

\implies \square

Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

So, $s(t)/c = t - t_0$

$\implies s(t)/c - t$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

So, $s(t)/c = t - t_0$

$$\implies s(t)/c - t = -t_0$$



Exercise.

If $\|\dot{\gamma}(t)\|$ is a constant c , show that $t - s(t)/c$ is a constant. Here $s(t)$ is the arc-length function “measured from (fixed) t_0 ”.

Solution.

$$\begin{aligned}s(t) &= \int_{t_0}^t \|\dot{\gamma}(t)\| dt \\ &= \int_{t_0}^t c dt \\ &= ct - ct_0\end{aligned}$$

$$\begin{aligned}\text{So, } s(t)/c &= t - t_0 \\ \implies s(t)/c - t &= -t_0\end{aligned}$$



Curvature

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$$\|\dot{\gamma}(t)\|$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$\gamma(t)$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t), r \sin(t))$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t), r \sin(t))$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t)$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (-r/r \sin(t/r), r/r \cos(t/r))$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (-r/r \sin(t/r), r/r \cos(t/r))$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), r/r \cos(t/r))$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t)$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\|\ddot{\gamma}(t)\| = 1/r$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\|\ddot{\gamma}(t)\| = 1/r$$

Curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\kappa(t) = \|\ddot{\gamma}(t)\| = 1/r$$

Curvature

Exercise. Show that the curvature

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), r/r \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \cos(t/r), -1/r \sin(t/r))$$

$$\kappa(t) = \|\ddot{\gamma}(t)\| = 1/r$$

Curvature

Exercise. Show that the curvature at any point of any line

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\kappa(t) = \|\ddot{\gamma}(t)\| = 1/r$$

Curvature

Exercise. Show that the curvature at any point of any line is 0.

Definition. $\gamma : (\alpha, \beta) \rightarrow \mathbb{R}^2$, *unit speed* parametrization

$\|\ddot{\gamma}(t)\|$ is the curvature, denoted $\kappa(t)$, at $\gamma(t)$

Example.

$$\gamma(t) = (r \cos(t/r), r \sin(t/r))$$

now unit speed

$$\dot{\gamma}(t) = (\sin(t/r), \cos(t/r))$$

$$\ddot{\gamma}(t) = (-1/r \sin(t/r), 1/r \cos(t/r))$$

$$\kappa(t) = \|\ddot{\gamma}(t)\| = 1/r$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently,

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition),

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t)$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) =$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t)$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) =$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 =$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) =$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) =$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) =$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2 +$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t) \cdot \dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t) \cdot \ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t) \cdot \ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

$$\kappa(t)$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$
and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$\begin{aligned}s'(t) &= \|\dot{\gamma}(t)\| \\ (s'(t))^2 &= \dot{\gamma}(t) \cdot \dot{\gamma}(t) \\ 2(s'(t))s''(t) &= 2\dot{\gamma}(t) \cdot \ddot{\gamma}(t) \\ 2\|\dot{\gamma}(t)\|s''(t) &= 2\dot{\gamma}(t) \cdot \ddot{\gamma}(t) \\ s''(t) &= \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}\end{aligned}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t)) \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\kappa(t) \; = \; \|\ddot{\tilde{\gamma}}(s(t))\|$$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|} = \ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2$$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$

and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$$\kappa(t) \; = \; \|\ddot{\tilde{\gamma}}(s(t))\|$$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2 = \ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

Deriving a general formula for curvature

$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$
Equivalently, $\gamma(t) = \tilde{\gamma}(s(t))$
and (by definition), $\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$

$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$

$s'(t) = \|\dot{\gamma}(t)\|$
 $(s'(t))^2 = \dot{\gamma}(t).\dot{\gamma}(t)$
 $2(s'(t))s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$
 $2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t).\ddot{\gamma}(t)$
 $s''(t) = \frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$

$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$

$\ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2 = \ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$

$\kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$
 $= \left\| \frac{\ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t))\frac{\dot{\gamma}(t).\ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}}{\|\dot{\gamma}(t)\|^2} \right\|$

Deriving a general formula for curvature

$$\tilde{\gamma}(\tilde{t}) = \gamma(s^{-1}(\tilde{t}))$$

$$\text{Equivalently, } \gamma(t) = \tilde{\gamma}(s(t))$$

$$\text{and (by definition), } \kappa(t) = \|\ddot{\tilde{\gamma}}(s(t))\|$$

$$\dot{\gamma}(t) = \dot{\tilde{\gamma}}(s(t))s'(t) = \dot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|$$

$$s'(t) = \|\dot{\gamma}(t)\|$$

$$(s'(t))^2 = \dot{\gamma}(t) \cdot \dot{\gamma}(t)$$

$$2(s'(t))s''(t) = 2\dot{\gamma}(t) \cdot \ddot{\gamma}(t)$$

$$2\|\dot{\gamma}(t)\|s''(t) = 2\dot{\gamma}(t) \cdot \ddot{\gamma}(t)$$

$$s''(t) = \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\gamma}(t) = \underbrace{\ddot{\tilde{\gamma}}(s(t))}_{\text{to get } \kappa(t)} \|\dot{\gamma}(t)\|^2 + \dot{\tilde{\gamma}}(s(t)) \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\ddot{\tilde{\gamma}}(s(t))\|\dot{\gamma}(t)\|^2 = \ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t)) \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}$$

$$\begin{aligned} \kappa(t) &= \|\ddot{\tilde{\gamma}}(s(t))\| \\ &= \left\| \frac{\ddot{\gamma}(t) - \dot{\tilde{\gamma}}(s(t)) \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}}{\|\dot{\gamma}(t)\|^2} \right\| \\ &= \left\| \frac{\ddot{\gamma}(t) - \frac{\dot{\gamma}(t)}{\|\dot{\gamma}(t)\|} \frac{\dot{\gamma}(t) \cdot \ddot{\gamma}(t)}{\|\dot{\gamma}(t)\|}}{\|\dot{\gamma}(t)\|^2} \right\| \end{aligned}$$