

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,

We will define smooth functions on surfaces

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,

We study the surface using a patch

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$

if,

given a (regular) surface patch  $\sigma : U \rightarrow S$ ,

so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,

Which contains  $p$

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$

if,

given a (regular) surface patch  $\sigma : U \rightarrow S$ ,

so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,

$f \circ \sigma$

We view the surface in terms of a patch

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$

if,

given a (regular) surface patch  $\sigma : U \rightarrow S$ ,

so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,

$f \circ \sigma$

Now its domain is a subset of  $\mathbb{R}^2$

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

so we know what it means to be smooth

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,

Of course, we need to check that it does not depend on the chosen patch

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth,

This will always happen but we will prove it later



# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth,

Let us examine the relationship between  $f \circ \sigma$  and  $f \circ \tilde{\sigma}$

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

We know the relationship between the two patches

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Composing with  $f$

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,

Since the composition of smooth functions is smooth

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth



# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth



# Smooth functions

## Definition.

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)



# Smooth functions

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function**

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

We now similarly study functions between surfaces via their surface patches



# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$



# Smooth functions

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if, given a (regular) surface patch  $\sigma : U \rightarrow S$ , so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

if,

given a (regular) surface patch  $\sigma : U \rightarrow S$ ,

so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,

$f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,

where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse

is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  $f \circ \sigma$  and  $\Phi$  are smooth)

This time there is a surface patch not just for the domain

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\begin{aligned}\tilde{\sigma} &= \sigma \circ \Phi \\ f \circ \tilde{\sigma} &= f \circ \sigma \circ \Phi\end{aligned}$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )



# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )  
and  $\sigma_2 : U \rightarrow S_2$ ,

but also for the co-domain

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )  
and  $\sigma_2 : U \rightarrow S_2$ ,

This time we also compose by  $\sigma_2^{-1}$  so that the the input and output are from  $U_1$  and  $U_2$ , respectively

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\begin{aligned}\tilde{\sigma} &= \sigma \circ \Phi \\ f \circ \tilde{\sigma} &= f \circ \sigma \circ \Phi\end{aligned}$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )  
and  $\sigma_2 : U \rightarrow S_2$ ,  
 $\sigma_2^{-1} \circ f \circ \sigma_1$  is smooth.



# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )  
and  $\sigma_2 : U \rightarrow S_2$ ,  
 $\sigma_2^{-1} \circ f \circ \sigma_1$  is smooth.

**Exercise.** Show that the definition of a smooth map does not depend on the choice of parametrizations.

**Definition.**

This exercise tells us why the definition does not depend on the choice of patches

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )  
and  $\sigma_2 : U \rightarrow S_2$ ,  
 $\sigma_2^{-1} \circ f \circ \sigma_1$  is smooth.

**Exercise.** Show that the definition of a smooth map does not depend on the choice of parametrizations.

**Definition.** Consider a smooth map,  $f : S_1 \rightarrow S_2$

$f$  naturally defines a map on the tangent spaces as we shall now see



# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )  
and  $\sigma_2 : U \rightarrow S_2$ ,  
 $\sigma_2^{-1} \circ f \circ \sigma_1$  is smooth.

**Exercise.** Show that the definition of a smooth map does not depend on the choice of parametrizations.

**Definition.** Consider a smooth map,  $f : S_1 \rightarrow S_2$   
so that  $f(p) = q$  for some  $p \in S_1$  and  $q \in S_2$ .



# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\tilde{\sigma} = \sigma \circ \Phi$$

$$f \circ \tilde{\sigma} = f \circ \sigma \circ \Phi$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )  
and  $\sigma_2 : U \rightarrow S_2$ ,  
 $\sigma_2^{-1} \circ f \circ \sigma_1$  is smooth.

**Exercise.** Show that the definition of a smooth map does not depend on the choice of parametrizations.

**Definition.** Consider a smooth map,  $f : S_1 \rightarrow S_2$   
so that  $f(p) = q$  for some  $p \in S_1$  and  $q \in S_2$ .  
Let  $\mathbf{v} \in T_p(S_1)$  denote a tangent vector at  $p$ .



# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\begin{aligned}\tilde{\sigma} &= \sigma \circ \Phi \\ f \circ \tilde{\sigma} &= f \circ \sigma \circ \Phi\end{aligned}$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )  
and  $\sigma_2 : U \rightarrow S_2$ ,  
 $\sigma_2^{-1} \circ f \circ \sigma_1$  is smooth.

**Exercise.** Show that the definition of a smooth map does not depend on the choice of parametrizations.

**Definition.** Consider a smooth map,  $f : S_1 \rightarrow S_2$   
so that  $f(p) = q$  for some  $p \in S_1$  and  $q \in S_2$ .  
Let  $\mathbf{v} \in T_p(S_1)$  denote a tangent vector at  $p$ .  
i.e.  $\mathbf{v} = \dot{\gamma}(t_0)$  for some  $\gamma : (\alpha, \beta) \rightarrow S_1$  and  $t_0 \in (\alpha, \beta)$ .

As usual, the tangent vector is a velocity vector of some curve on the surface

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\begin{aligned}\tilde{\sigma} &= \sigma \circ \Phi \\ f \circ \tilde{\sigma} &= f \circ \sigma \circ \Phi\end{aligned}$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )  
and  $\sigma_2 : U \rightarrow S_2$ ,  
 $\sigma_2^{-1} \circ f \circ \sigma_1$  is smooth.

**Exercise.** Show that the definition of a smooth map does not depend on the choice of parametrizations.

**Definition.** Consider a smooth map,  $f : S_1 \rightarrow S_2$   
so that  $f(p) = q$  for some  $p \in S_1$  and  $q \in S_2$ .  
Let  $\mathbf{v} \in T_p(S_1)$  denote a tangent vector at  $p$ .  
i.e.  $\mathbf{v} = \dot{\gamma}(t_0)$  for some  $\gamma : (\alpha, \beta) \rightarrow S_1$  and  $t_0 \in (\alpha, \beta)$ .  
Define  $d_p f : T_p(S_1) \rightarrow T_p(S_2)$   
(where  $T_p(S)$  denotes the tangent space of  $S$  at  $p$ )  
by  $d_p f(\mathbf{v}) = \frac{d}{dt} f(\gamma(t)) \in T_p(S_2)$

We simply consider the velocity vector of the image of that curve

# Smooth functions

**Definition.**  $f : S \rightarrow \mathbb{R}$  is called a **smooth map** at  $p$  if,  
given a (regular) surface patch  $\sigma : U \rightarrow S$ ,  
so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ ,  
 $f \circ \sigma$  is smooth at  $(x_0, y_0)$ .

If  $\tilde{\sigma} : \tilde{U} \rightarrow S$  is another surface patch so that,  $\tilde{\sigma} = \sigma \circ \Phi$ ,  
where  $\Phi : \tilde{U} \rightarrow U$  is smooth, invertible, and the inverse  
is smooth, Since,

$$\begin{aligned}\tilde{\sigma} &= \sigma \circ \Phi \\ f \circ \tilde{\sigma} &= f \circ \sigma \circ \Phi\end{aligned}$$

Therefore,  $f \circ \sigma$  smooth  $\implies f \circ \tilde{\sigma}$  is smooth (because,  
 $f \circ \sigma$  and  $\Phi$  are smooth)

**Definition.**  $f : S_1 \rightarrow S_2$  is said to be a **smooth function** at  $p \in S_1$  if, given (regular) surface patches  
 $\sigma_1 : U \rightarrow S_1$   
(so that  $p \in \sigma(U)$ ,  $p = \sigma(x_0, y_0)$ )  
and  $\sigma_2 : U \rightarrow S_2$ ,  
 $\sigma_2^{-1} \circ f \circ \sigma_1$  is smooth.

**Exercise.** Show that the definition of a smooth map does not depend on the choice of parametrizations.

**Definition.** Consider a smooth map,  $f : S_1 \rightarrow S_2$   
so that  $f(p) = q$  for some  $p \in S_1$  and  $q \in S_2$ .  
Let  $\mathbf{v} \in T_p(S_1)$  denote a tangent vector at  $p$ .  
i.e.  $\mathbf{v} = \dot{\gamma}(t_0)$  for some  $\gamma : (\alpha, \beta) \rightarrow S_1$  and  $t_0 \in (\alpha, \beta)$ .  
Define  $d_p f : T_p(S_1) \rightarrow T_p(S_2)$   
(where  $T_p(S)$  denotes the tangent space of  $S$  at  $p$ )  
by  $d_p f(\mathbf{v}) = \frac{d}{dt} f(\gamma(t)) \in T_p(S_2)$

and define that to be the image of  $\mathbf{v}$  under  $d_p f$

$$f : S_1 \rightarrow S_2,$$

We now try to describe  $d_p f$  in terms of the surface patch

$$f : S_1 \rightarrow S_2,$$

$$\sigma_2^{-1}(f(\sigma_1(x,y))) = (g_1(x,y), g_2(x,y))$$

Here is  $f$  in terms of the surface patch

$$f : S_1 \rightarrow S_2,$$

$$\sigma_2^{-1}(f(\sigma_1(x, y))) = (g_1(x, y), g_2(x, y))$$

$$f(\sigma_1(x(t), y(t))) = \sigma_2(g_1(x(t), y(t)), g_2(x(t), y(t)))$$

And this is the image of  $\gamma$  under  $f$



$$f : S_1 \rightarrow S_2,$$

$$\sigma_2^{-1}(f(\sigma_1(x, y))) = (g_1(x, y), g_2(x, y))$$

$$f(\sigma_1(x(t), y(t))) = \sigma_2(g_1(x(t), y(t)), g_2(x(t), y(t)))$$

$$\frac{\mathrm{d}}{\mathrm{d}t}f(\sigma_1(x(t), y(t))) = g_1'(x(t), y(t))\sigma_{2x} + g_2'(x(t), y(t))\sigma_{2y}$$

Written in a form that will allow us to write it in terms of  $\sigma_{2x}$  and  $\sigma_{2y}$

$$f : S_1 \rightarrow S_2,$$

$$\sigma_2^{-1}(f(\sigma_1(x, y))) = (g_1(x, y), g_2(x, y))$$

$$f(\sigma_1(x(t), y(t))) = \sigma_2(g_1(x(t), y(t)), g_2(x(t), y(t)))$$

$$\begin{aligned} \frac{d}{dt}f(\sigma_1(x(t), y(t))) &= g'_1(x(t), y(t))\sigma_{2_x} + g'_2(x(t), y(t))\sigma_{2_y} \\ &= (x'(t)g_{1_x}(x(t), y(t)) + y'(t)g_{1_y}(x(t), y(t))\sigma_x(x(t), y(t)) \\ &\quad + (x'(t)g_{2_x}(x(t), y(t)) + y'(t)g_{2_y}(x(t), y(t))\sigma_y(x(t), y(t)) \end{aligned}$$

Now we apply chain rule to each coefficient

$$f : S_1 \rightarrow S_2,$$

$$\sigma_2^{-1}(f(\sigma_1(x, y))) = (g_1(x, y), g_2(x, y))$$

$$f(\sigma_1(x(t), y(t))) = \sigma_2(g_1(x(t), y(t)), g_2(x(t), y(t)))$$

$$\begin{aligned} \frac{d}{dt}f(\sigma_1(x(t), y(t))) &= g'_1(x(t), y(t))\sigma_{2x} + g'_2(x(t), y(t))\sigma_{2y} \\ &= (x'(t)g_{1x}(x(t), y(t)) + y'(t)g_{1y}(x(t), y(t))\sigma_x(x(t), y(t)) \\ &\quad + (x'(t)g_{2x}(x(t), y(t)) + y'(t)g_{2y}(x(t), y(t))\sigma_y(x(t), y(t)) \end{aligned}$$

In terms of coordinates,

$$= \begin{pmatrix} g_{1x}(t) & g_{1y}(t) \\ g_{2x}(t) & g_{2y}(t) \end{pmatrix}$$

And write it in terms of coordinates

$$f : S_1 \rightarrow S_2,$$

$$\sigma_2^{-1}(f(\sigma_1(x, y))) = (g_1(x, y), g_2(x, y))$$

$$f(\sigma_1(x(t), y(t))) = \sigma_2(g_1(x(t), y(t)), g_2(x(t), y(t)))$$

$$\begin{aligned} \frac{d}{dt}f(\sigma_1(x(t), y(t))) &= g'_1(x(t), y(t))\sigma_{2_x} + g'_2(x(t), y(t))\sigma_{2_y} \\ &= (x'(t)g_{1_x}(x(t), y(t)) + y'(t)g_{1_y}(x(t), y(t))\sigma_x(x(t), y(t)) \\ &\quad + (x'(t)g_{2_x}(x(t), y(t)) + y'(t)g_{2_y}(x(t), y(t))\sigma_y(x(t), y(t)) \end{aligned}$$

In terms of coordinates,

$$= \begin{pmatrix} g_{1_x}(t) & g_{1_y}(t) \\ g_{2_x}(t) & g_{2_y}(t) \end{pmatrix} \begin{pmatrix} x'(t) \\ y'(t) \end{pmatrix}$$



$$f : S_1 \rightarrow S_2,$$

$$\sigma_2^{-1}(f(\sigma_1(x, y))) = (g_1(x, y), g_2(x, y))$$

$$f(\sigma_1(x(t), y(t))) = \sigma_2(g_1(x(t), y(t)), g_2(x(t), y(t)))$$

$$\begin{aligned} \frac{d}{dt}f(\sigma_1(x(t), y(t))) &= g'_1(x(t), y(t))\sigma_{2x} + g'_2(x(t), y(t))\sigma_{2y} \\ &= (x'(t)g_{1x}(x(t), y(t)) + y'(t)g_{1y}(x(t), y(t))\sigma_x(x(t), y(t)) \\ &\quad + (x'(t)g_{2x}(x(t), y(t)) + y'(t)g_{2y}(x(t), y(t))\sigma_y(x(t), y(t)) \end{aligned}$$

In terms of coordinates,

$$\begin{aligned} &= \begin{pmatrix} g_{1x}(t) & g_{1y}(t) \\ g_{2x}(t) & g_{2y}(t) \end{pmatrix} \begin{pmatrix} x'(t) \\ y'(t) \end{pmatrix} \\ &= J(\sigma_2^{-1} \circ f \circ \sigma_1) \begin{pmatrix} x'(t) \\ y'(t) \end{pmatrix} \end{aligned}$$

Notice that the familiar Jacobian matrix shows up again

For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,

The inner product of two tangent vectors is simply the dot product

For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$



For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

The angular bracket notation only emphasizes that  $\mathbf{v}_1$  and  $\mathbf{v}_2$  must be tangent vectors



For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

We will try to express this in terms of the surface patch

For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\mathbf{v}_1 = \dot{\gamma}_1(t_0)$$

First note that by definition they are velocity vectors

For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\mathbf{v}_1 = \dot{\gamma}_1(t_0)$$

$$\mathbf{v}_2 = \dot{\gamma}_2(t_0)$$



For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\mathbf{v}_1 = \dot{\gamma}_1(t_0) = \frac{d}{dt} \sigma(x_1(t_0), y_1(t_0))$$

$$\mathbf{v}_2 = \dot{\gamma}_2(t_0)$$

And now we use chain rule to express them in terms of  $\sigma_x$  and  $\sigma_y$

For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\mathbf{v}_1 = \dot{\gamma}_1(t_0) = \frac{\mathrm{d}}{\mathrm{d}t}\sigma(x_1(t_0), y_1(t_0))$$

$$\mathbf{v}_2 = \dot{\gamma}_2(t_0) = \frac{\mathrm{d}}{\mathrm{d}t}\sigma(x_2(t_0), y_2(t_0))$$



For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,

$$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$$

$$\begin{aligned}\mathbf{v}_1 &= \dot{\gamma}_1(t_0) = \frac{\mathrm{d}}{\mathrm{d}t}\sigma(x_1(t_0), y_1(t_0)) = x_1'(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y_1'(t_0)\sigma_y(x_1(t_0), y_1(t_0)) \\ \mathbf{v}_2 &= \dot{\gamma}_2(t_0) = \frac{\mathrm{d}}{\mathrm{d}t}\sigma(x_2(t_0), y_2(t_0))\end{aligned}$$



For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,

$$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$$

$$\begin{aligned} \mathbf{v}_1 &= \dot{\gamma}_1(t_0) = \frac{d}{dt}\sigma(x_1(t_0), y_1(t_0)) = x_1'(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y_1'(t_0)\sigma_y(x_1(t_0), y_1(t_0)) \\ \mathbf{v}_2 &= \dot{\gamma}_2(t_0) = \frac{d}{dt}\sigma(x_2(t_0), y_2(t_0)) = x_2'(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y_2'(t_0)\sigma_y(x_2(t_0), y_2(t_0)) \end{aligned}$$



For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\begin{aligned}\mathbf{v}_1 &= \dot{\gamma}_1(t_0) = \frac{d}{dt}\sigma(x_1(t_0), y_1(t_0)) = x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0)) \\ \mathbf{v}_2 &= \dot{\gamma}_2(t_0) = \frac{d}{dt}\sigma(x_2(t_0), y_2(t_0)) = x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))\end{aligned}$$

$$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle = \mathbf{v}_1 \cdot \mathbf{v}_2$$

Finally, using them in the dot product



For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,

$$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$$

$$\begin{aligned} \mathbf{v}_1 &= \dot{\gamma}_1(t_0) = \frac{d}{dt}\sigma(x_1(t_0), y_1(t_0)) = x_1'(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y_1'(t_0)\sigma_y(x_1(t_0), y_1(t_0)) \\ \mathbf{v}_2 &= \dot{\gamma}_2(t_0) = \frac{d}{dt}\sigma(x_2(t_0), y_2(t_0)) = x_2'(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y_2'(t_0)\sigma_y(x_2(t_0), y_2(t_0)) \end{aligned}$$

$$\begin{aligned} \langle \mathbf{v}_1, \mathbf{v}_2 \rangle &= \mathbf{v}_1 \cdot \mathbf{v}_2 \\ &= (x_1'(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y_1'(t_0)\sigma_y(x_1(t_0), y_1(t_0))).(x_2'(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y_2'(t_0)\sigma_y(x_2(t_0), y_2(t_0))) \end{aligned}$$



For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\begin{aligned}\mathbf{v}_1 &= \dot{\gamma}_1(t_0) = \frac{d}{dt}\sigma(x_1(t_0), y_1(t_0)) = x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0)) \\ \mathbf{v}_2 &= \dot{\gamma}_2(t_0) = \frac{d}{dt}\sigma(x_2(t_0), y_2(t_0)) = x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))\end{aligned}$$

$$\begin{aligned}\langle \mathbf{v}_1, \mathbf{v}_2 \rangle &= \mathbf{v}_1 \cdot \mathbf{v}_2 \\ &= (x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0))).(x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))) \\ &= x'_1(t_0)x'_2(t_0)E(x(t_0), y(t_0)) + x'_1(t_0)y'_2(t_0)F(x(t_0), y(t_0)) \\ &\quad + y'_1(t_0)x'_2(t_0)F(x(t_0), y(t_0)) + y'_1(t_0)y'_2(t_0)G(x(t_0), y(t_0))\end{aligned}$$

Distributing and recognizing the appearance of  $E$ ,  $F$ , and  $G$

For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\mathbf{v}_1 = \dot{\gamma}_1(t_0) = \frac{d}{dt}\sigma(x_1(t_0), y_1(t_0)) = x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0))$$

$$\mathbf{v}_2 = \dot{\gamma}_2(t_0) = \frac{d}{dt}\sigma(x_2(t_0), y_2(t_0)) = x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))$$

$$\begin{aligned} \langle \mathbf{v}_1, \mathbf{v}_2 \rangle &= \mathbf{v}_1 \cdot \mathbf{v}_2 \\ &= (x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0))).(x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))) \\ &= x'_1(t_0)x'_2(t_0)E(x(t_0), y(t_0)) + x'_1(t_0)y'_2(t_0)F(x(t_0), y(t_0)) \\ &\quad + y'_1(t_0)x'_2(t_0)F(x(t_0), y(t_0)) + y'_1(t_0)y'_2(t_0)G(x(t_0), y(t_0)) \end{aligned}$$

Observe that since  $\mathbf{v}_1$  and  $\mathbf{v}_2$  are based on the same point,  $\gamma_1(t_0) = \gamma_2(t_0)$

For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\begin{aligned}\mathbf{v}_1 &= \dot{\gamma}_1(t_0) = \frac{d}{dt}\sigma(x_1(t_0), y_1(t_0)) = x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0)) \\ \mathbf{v}_2 &= \dot{\gamma}_2(t_0) = \frac{d}{dt}\sigma(x_2(t_0), y_2(t_0)) = x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))\end{aligned}$$

$$\begin{aligned}\langle \mathbf{v}_1, \mathbf{v}_2 \rangle &= \mathbf{v}_1 \cdot \mathbf{v}_2 \\ &= (x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0))).(x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))) \\ &= x'_1(t_0)x'_2(t_0)E(x(t_0), y(t_0)) + x'_1(t_0)y'_2(t_0)F(x(t_0), y(t_0)) \\ &\quad + y'_1(t_0)x'_2(t_0)F(x(t_0), y(t_0)) + y'_1(t_0)y'_2(t_0)G(x(t_0), y(t_0))\end{aligned}$$

So  $(x_1(t_0), y_1(t_0)) = (x_2(t_0), y_2(t_0))$

For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\mathbf{v}_1 = \dot{\gamma}_1(t_0) = \frac{d}{dt}\sigma(x_1(t_0), y_1(t_0)) = x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0))$$

$$\mathbf{v}_2 = \dot{\gamma}_2(t_0) = \frac{d}{dt}\sigma(x_2(t_0), y_2(t_0)) = x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))$$

$$\begin{aligned} \langle \mathbf{v}_1, \mathbf{v}_2 \rangle &= \mathbf{v}_1 \cdot \mathbf{v}_2 \\ &= (x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0))) \cdot (x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))) \\ &= x'_1(t_0)x'_2(t_0)E(x(t_0), y(t_0)) + x'_1(t_0)y'_2(t_0)F(x(t_0), y(t_0)) \\ &\quad + y'_1(t_0)x'_2(t_0)F(x(t_0), y(t_0)) + y'_1(t_0)y'_2(t_0)G(x(t_0), y(t_0)) \\ &= \begin{pmatrix} x'_1(t_0) & y'_1(t_0) \end{pmatrix} \end{aligned}$$

But now observe that this can be expressed in matrix form

For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\begin{aligned}\mathbf{v}_1 &= \dot{\gamma}_1(t_0) = \frac{d}{dt}\sigma(x_1(t_0), y_1(t_0)) = x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0)) \\ \mathbf{v}_2 &= \dot{\gamma}_2(t_0) = \frac{d}{dt}\sigma(x_2(t_0), y_2(t_0)) = x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))\end{aligned}$$

$$\begin{aligned}\langle \mathbf{v}_1, \mathbf{v}_2 \rangle &= \mathbf{v}_1 \cdot \mathbf{v}_2 \\ &= (x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0))).(x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))) \\ &= x'_1(t_0)x'_2(t_0)E(x(t_0), y(t_0)) + x'_1(t_0)y'_2(t_0)F(x(t_0), y(t_0)) \\ &\quad + y'_1(t_0)x'_2(t_0)F(x(t_0), y(t_0)) + y'_1(t_0)y'_2(t_0)G(x(t_0), y(t_0)) \\ &= \begin{pmatrix} x'_1(t_0) & y'_1(t_0) \end{pmatrix} \begin{pmatrix} E(x(t_0), y(t_0)) & F(x(t_0), y(t_0)) \\ F(x(t_0), y(t_0)) & G(x(t_0), y(t_0)) \end{pmatrix}\end{aligned}$$

■

For  $\mathbf{v}_1, \mathbf{v}_2 \in T_p(S)$ ,  
 $\langle \mathbf{v}_1, \mathbf{v}_2 \rangle := \mathbf{v}_1 \cdot \mathbf{v}_2$

$$\begin{aligned}\mathbf{v}_1 &= \dot{\gamma}_1(t_0) = \frac{d}{dt}\sigma(x_1(t_0), y_1(t_0)) = x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0)) \\ \mathbf{v}_2 &= \dot{\gamma}_2(t_0) = \frac{d}{dt}\sigma(x_2(t_0), y_2(t_0)) = x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))\end{aligned}$$

$$\begin{aligned}\langle \mathbf{v}_1, \mathbf{v}_2 \rangle &= \mathbf{v}_1 \cdot \mathbf{v}_2 \\ &= (x'_1(t_0)\sigma_x(x_1(t_0), y_1(t_0)) + y'_1(t_0)\sigma_y(x_1(t_0), y_1(t_0))).(x'_2(t_0)\sigma_x(x_2(t_0), y_2(t_0)) + y'_2(t_0)\sigma_y(x_2(t_0), y_2(t_0))) \\ &= x'_1(t_0)x'_2(t_0)E(x(t_0), y(t_0)) + x'_1(t_0)y'_2(t_0)F(x(t_0), y(t_0)) \\ &\quad + y'_1(t_0)x'_2(t_0)F(x(t_0), y(t_0)) + y'_1(t_0)y'_2(t_0)G(x(t_0), y(t_0)) \\ &= \begin{pmatrix} x'_1(t_0) & y'_1(t_0) \end{pmatrix} \begin{pmatrix} E(x(t_0), y(t_0)) & F(x(t_0), y(t_0)) \\ F(x(t_0), y(t_0)) & G(x(t_0), y(t_0)) \end{pmatrix} \begin{pmatrix} x'_2(t_0) \\ y'_2(t_0) \end{pmatrix}\end{aligned}$$

■

Surface

Surface patch

---

We will summarize how various concepts appear in terms of surface patches



Surface	Surface patch
$p \in S$	

A surface patch gives two coordinates to every point on part of a surface

Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	

To every subset in the patch of  $S$ , it associates a subset in  $U$

Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	

It associates to every curve on that part of the surface, a curve in  $U$

Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	

It provides a basis  $\sigma_x$  and  $\sigma_y$ , and tangent vectors are written in terms of them

Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$





Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	

To a function with domain  $S_1$ , it associates a function with domain  $U$

Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	

To a function with surfaces as both domains and ranges, it associates a function between the domains of their patches.

Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$
$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle$	

To the inner product, it associates the matrix of “first fundamental form”

Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$
$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle$	$\begin{pmatrix} x'_1 & y'_1 \end{pmatrix} \begin{pmatrix} E & F \\ F & G \end{pmatrix} \begin{pmatrix} x'_2 \\ y'_2 \end{pmatrix}$ , where $\mathbf{v}_i = x'_i\sigma_x + y'_i\sigma_y$



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$
$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle$	$\begin{pmatrix} x'_1 & y'_1 \end{pmatrix} \begin{pmatrix} E & F \\ F & G \end{pmatrix} \begin{pmatrix} x'_2 \\ y'_2 \end{pmatrix}$ , where $\mathbf{v}_i = x'_i\sigma_x + y'_i\sigma_y$



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$
$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle$	$\begin{pmatrix} x'_1 & y'_1 \end{pmatrix} \begin{pmatrix} E & F \\ F & G \end{pmatrix} \begin{pmatrix} x'_2 \\ y'_2 \end{pmatrix}$ , where $\mathbf{v}_i = x'_i\sigma_x + y'_i\sigma_y$
$d_p(f) : T_p(S_1) \rightarrow T_{f(p)}(S_2)$	

To a derivative of a function between two surfaces, it associates the “Jacobian” matrix



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$
$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle$	$\begin{pmatrix} x'_1 & y'_1 \end{pmatrix} \begin{pmatrix} E & F \\ F & G \end{pmatrix} \begin{pmatrix} x'_2 \\ y'_2 \end{pmatrix}$ , where $\mathbf{v}_i = x'_i\sigma_x + y'_i\sigma_y$
$d_p(f) : T_p(S_1) \rightarrow T_{f(p)}(S_2)$	$\begin{pmatrix} g_{1x} & g_{1y} \\ g_{2x} & g_{2y} \end{pmatrix}$ , where $(g_1, g_2) = \sigma_2^{-1} \circ f \circ \sigma_1$



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$
$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle$	$\begin{pmatrix} x'_1 & y'_1 \end{pmatrix} \begin{pmatrix} E & F \\ F & G \end{pmatrix} \begin{pmatrix} x'_2 \\ y'_2 \end{pmatrix}$ , where $\mathbf{v}_i = x'_i\sigma_x + y'_i\sigma_y$
$d_p(f) : T_p(S_1) \rightarrow T_{f(p)}(S_2)$	$\begin{pmatrix} g_{1x} & g_{1y} \\ g_{2x} & g_{2y} \end{pmatrix}$ , where $(g_1, g_2) = \sigma_2^{-1} \circ f \circ \sigma_1$
$\ \sigma_x \times \sigma_y\ $	

To the infinitesimal area, it associates the determinant of the first fundamental form matrix

Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$
$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle$	$\begin{pmatrix} x'_1 & y'_1 \end{pmatrix} \begin{pmatrix} E & F \\ F & G \end{pmatrix} \begin{pmatrix} x'_2 \\ y'_2 \end{pmatrix}$ , where $\mathbf{v}_i = x'_i\sigma_x + y'_i\sigma_y$
$d_p(f) : T_p(S_1) \rightarrow T_{f(p)}(S_2)$	$\begin{pmatrix} g_{1x} & g_{1y} \\ g_{2x} & g_{2y} \end{pmatrix}$ , where $(g_1, g_2) = \sigma_2^{-1} \circ f \circ \sigma_1$
$\ \sigma_x \times \sigma_y\ $	$ EG - F^2  = \det \begin{pmatrix} E & F \\ F & G \end{pmatrix}$



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$
$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle$	$\begin{pmatrix} x'_1 & y'_1 \end{pmatrix} \begin{pmatrix} E & F \\ F & G \end{pmatrix} \begin{pmatrix} x'_2 \\ y'_2 \end{pmatrix}$ , where $\mathbf{v}_i = x'_i\sigma_x + y'_i\sigma_y$
$d_p(f) : T_p(S_1) \rightarrow T_{f(p)}(S_2)$	$\begin{pmatrix} g_{1x} & g_{1y} \\ g_{2x} & g_{2y} \end{pmatrix}$ , where $(g_1, g_2) = \sigma_2^{-1} \circ f \circ \sigma_1$
$\ \sigma_x \times \sigma_y\ $	$ EG - F^2  = \det \begin{pmatrix} E & F \\ F & G \end{pmatrix}$
$\text{Area} = \int_{\sigma(U)} \ \sigma_x \times \sigma_y\ $	

And to the area, the integral of the above determinant.

Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$
$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle$	$\begin{pmatrix} x'_1 & y'_1 \end{pmatrix} \begin{pmatrix} E & F \\ F & G \end{pmatrix} \begin{pmatrix} x'_2 \\ y'_2 \end{pmatrix}$ , where $\mathbf{v}_i = x'_i\sigma_x + y'_i\sigma_y$
$d_p(f) : T_p(S_1) \rightarrow T_{f(p)}(S_2)$	$\begin{pmatrix} g_{1x} & g_{1y} \\ g_{2x} & g_{2y} \end{pmatrix}$ , where $(g_1, g_2) = \sigma_2^{-1} \circ f \circ \sigma_1$
$\ \sigma_x \times \sigma_y\ $	$ EG - F^2  = \det \begin{pmatrix} E & F \\ F & G \end{pmatrix}$
$\text{Area} = \int_{\sigma(U)} \ \sigma_x \times \sigma_y\ $	$\int_U  EG - F^2 $



Surface	Surface patch
$p \in S$	$(x, y) \in U$ , where $\sigma(x, y) = p$
$A \subset S$	$B \subset U$ , where $\sigma(B) = A$
$\gamma : (\alpha, \beta) \rightarrow S$	$\delta : (\alpha, \beta) \rightarrow U$ , where $\gamma = \sigma \circ \delta$
$\mathbf{v} = \dot{\gamma}(t_0)$	$\mathbf{v} = x'\sigma_x + y'\sigma_y$ , where $\gamma(t) = \sigma(x(t), y(t))$
$f : S_1 \rightarrow \mathbb{R}$	$g : U \rightarrow \mathbb{R}$ , where $g = f \circ \sigma$
$f : S_1 \rightarrow S_2$	$g : U_1 \rightarrow U_2$ , where $g = \sigma_2^{-1} \circ f \circ \sigma_1$
$\langle \mathbf{v}_1, \mathbf{v}_2 \rangle$	$\begin{pmatrix} x'_1 & y'_1 \end{pmatrix} \begin{pmatrix} E & F \\ F & G \end{pmatrix} \begin{pmatrix} x'_2 \\ y'_2 \end{pmatrix}$ , where $\mathbf{v}_i = x'_i\sigma_x + y'_i\sigma_y$
$d_p(f) : T_p(S_1) \rightarrow T_{f(p)}(S_2)$	$\begin{pmatrix} g_{1x} & g_{1y} \\ g_{2x} & g_{2y} \end{pmatrix}$ , where $(g_1, g_2) = \sigma_2^{-1} \circ f \circ \sigma_1$
$\ \sigma_x \times \sigma_y\ $	$ EG - F^2  = \det \begin{pmatrix} E & F \\ F & G \end{pmatrix}$
$\text{Area} = \int_{\sigma(U)} \ \sigma_x \times \sigma_y\ $	$\int_U  EG - F^2 $

